

# Effect of various level of phosphorus and spacing on growth and yield of black wheat (*Triticum aestivum* L.)

Comment [i-[1]: I think it is fixed on materials and methods part\*\*\*4 level not necessary

## Abstract

The study examined the effects of different phosphorus levels and plant spacings on the growth, yield, and economic viability of black wheat (*Triticum aestivum* L.), a variety known for its nutritional benefits. The experiment was conducted during the 2023–24 rabi season at the Crop Research Farm of Dev Bhoomi Uttarakhand University in Dehradun, India. A factorial randomized block design with three replications was employed. Four phosphorus levels (40, 50, 60, and 70 kg  $P_2O_5 \text{ ha}^{-1}$ ) were applied as diammonium phosphate (DAP) at planting, in combination with three plant spacings (15 cm x 10 cm, 20 cm x 10 cm, and 25 cm x 10 cm). The results showed that 70 kg  $P_2O_5 \text{ ha}^{-1}$  significantly improved growth parameters, yield attributes, and nitrogen uptake, while the optimal plant spacing of 25 cm x 10 cm further enhanced these outcomes. The highest grain and straw yields were obtained with 70 kg  $P_2O_5 \text{ ha}^{-1}$ , though 60 kg  $P_2O_5 \text{ ha}^{-1}$  provided similar yields with greater cost-effectiveness. Economic analysis indicated that the combination of 70 kg  $P_2O_5 \text{ ha}^{-1}$  and optimal spacing resulted in the highest net returns. These findings suggest that strategic phosphorus application, combined with appropriate plant spacing, can improve both the productivity and profitability of black wheat cultivation, offering a sustainable approach to agriculture.

Comment [i-[2]: Write in etalic form

Comment [i-[3]: Which spacing? Put in figures

## Keywords

Black wheat, Growth, Nitrogen uptake, Phosphorus, Yield

## Introduction

Black wheat (*Triticum aestivum* L.) is gaining attention due to its rich anthocyanin content and associated nutritional benefits, such as anti-inflammatory and anti-cancer properties (Nguyen et al., 2023; Sharma et al., 2021). This unique variety presents a nutrient-dense alternative to conventional wheat, prompting increased interest in its cultivation. Phosphorus ( $P_2O_5$ ) management is critical for optimizing growth and yield in black wheat, as phosphorus is essential for energy transfer and root development (Khan et al., 2023). However, phosphorus availability in soil can be limited, leading to the necessity of fertilizer applications. Excessive use of fertilizers raises concerns regarding environmental sustainability (Weeks & Hettiarachchi, 2019; Liu et al., 2023).

While recent studies on colored wheat varieties, such as purple wheat in China, have demonstrated that optimized phosphorus levels can enhance anthocyanin content and overall yield (Liu et al., 2023), research specifically targeting black wheat—particularly within the context of Indian agriculture—is still sparse. Existing literature often emphasizes high-yielding varieties and advanced agronomic practices to improve wheat production in India (Mishra et al., 2005; Joshi et al., 2007), yet the impact of phosphorus levels and plant spacing on black wheat has not been thoroughly investigated.

This study is warranted to fill this knowledge gap and provide insights into the effective management of phosphorus and plant spacing for black wheat cultivation. It aims to assess the combined effects of various phosphorus levels (40, 50, 60, and 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) and plant spacings (15 cm × 10 cm, 20 cm × 10 cm, and 25 cm × 10 cm) on growth and yield parameters. By exploring these factors, the research seeks to establish innovative agronomic practices that can enhance productivity and sustainability in black wheat farming. The objective of this study is to identify optimal phosphorus application rates and plant spacing that maximize growth, yield, and economic viability, thereby supporting farmers in adopting sustainable agricultural practices.

## MATERIAL AND METHODS

### Experimental site

The experiment was conducted during the 2023–2024 rabi season at the Crop Research Farm of Dev Bhoomi Uttarakhand University, Dehradun, Uttarakhand, India. This region experiences a subtropical climate, with summer temperatures reaching 35–39°C and winter lows of around 0.5°C. The average annual rainfall is approximately 1200 mm, primarily between November and April, with some additional winter rainfall.

Before establishing the experimental layout, soil samples were collected from various points within the 0–15 cm depth to assess the site's physical and chemical properties. These samples were pooled into a single representative composite sample for analysis. The soil at the study site was classified as sandy loam, with low organic carbon content but moderate levels of available phosphorus, nitrogen, and potassium. The wheat variety used in the study was Black 76.

### Field preparation and experimental details

The experiment followed a randomized block design with 12 treatment combinations, each replicated three times. Treatments were randomly arranged within each replication, resulting in a total of 36 plots.

Comment [i-4]: Use standard depth for cereal crops\*\*\* (0- 20 cm)

Comment [i-5]: RCBD

Each plot measured 4 m<sup>2</sup> (2 × 2 m). The area was irrigated before initial ploughing with a tractor-pulled disc plow. Additional cultivation involved cross-ploughing and levelling (planking) to achieve a finely textured soil surface. Before cross-ploughing, nitrogen and potassium were uniformly applied across the field and thoroughly incorporated into the soil during ploughing.

**Comment [i-[6]:** Check it, for different spacing similar plot size?

**Table 1. Chemical analysis of soil at pre -experiment stage**

| Parameter            | Value                     | Method                       | References                    |
|----------------------|---------------------------|------------------------------|-------------------------------|
| Organic carbon (%)   | 0.24%                     | Walkley and Black Method     | Walkley and Black Method 1934 |
| Available Nitrogen   | 237 kg ha <sup>-1</sup>   | Alkaline Permanganate Method | Subhaiah and Asija, 1956      |
| Available Phosphorus | 19.60 kg ha <sup>-1</sup> | Olsen's Colorimetric Method  | Olsen <i>et al.</i> , 1954    |
| Available Potassium  | 95.00 kg ha <sup>-1</sup> | Flame Photometer method      | Jackson, 1973                 |
| Soil pH              | 7.6                       | Glass electrode pH meter     | Jackson, 1973                 |

### Details of Treatment

The treatments comprised four levels of phosphorus, specifically 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P1), 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P2), 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P3), and 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P4), combined with three plant spacings: 15 cm × 10 cm (S1), 20 cm × 10 cm (S2), and 25 cm × 10 cm (S3).

**Comment [i-[7]:**

**Comment [i-[8]:**

**Comment [i-[9]:** Repeated

Phosphorus was applied in the form of diammonium phosphate (DAP) at the time of planting, with the rates corresponding to the phosphorus levels (P<sub>2</sub>O<sub>5</sub>). The specific application rates were calculated to ensure that the respective phosphorus levels were accurately met.

**Comment [i-[10]:** For phosphorous fertilizer experiments, TSP preferred rather than DAP

Other agronomic practices followed during the experiment included maintaining optimal soil moisture levels, weed management, and pest control as per standard agricultural guidelines. The crop was sown with a seed rate of approximately 100 kg ha<sup>-1</sup>, utilizing a high-yielding variety of black wheat known for its adaptability and resistance to common diseases. These practices ensured the establishment of a healthy crop, facilitating a comprehensive assessment of the treatments' effects on growth and yield.

**Comment [i-[11]:** Mention major once for black wheat production

**Table 1. Treatment combinations**

The experiment included a variety of treatment combinations, each designed to assess the effects of phosphorus levels and plant spacings on the growth and yield of black wheat. The treatment combinations consisted of the following: T1 combined 15 cm × 10 cm spacing with 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P1), T2 used the same spacing with 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P2), T3 with 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P3), and T4 with 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P4). For the second spacing treatment, T5 combined 20 cm × 10 cm spacing with 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P1), T6 with 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P2), T7 with 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P3), and T8 with 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P4). The third spacing treatment included T9 with 25 cm × 10 cm spacing and 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P1), T10 with 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P2), T11 with 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P3), and T12 with 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P4).

**Comment [i-[12]:** Repeated objective is not necessary

The plot size for each treatment was standardized to ensure consistent measurements and comparisons, with each plot measuring 4 m × 3 m. This design facilitated a thorough examination of the combined effects of phosphorus application and plant spacing on the performance of black wheat. A tube-well served as the irrigation source. In addition to pre-sowing irrigation, five irrigations were applied, each corresponding to a critical stage of plant growth. The crop was harvested once it reached maturity, determined through visual assessments on April 20, 2024. To ensure accuracy, two border rows on both sides of the field were excluded, and half a meter was trimmed from each plot's length. The harvested yield from the designated area was collected to calculate biomass yield. Each participant small plot was threshed using a manual thresher.

**Comment [i-[13]:** This part is already known b/s your experimental design mentioned above

**Comment [i-[14]:** ?

## Growth and yield estimations

### Plant height (cm)

For each plot, five plants were selected at random and marked for measuring their height at different time intervals. Height was measured at 30, 60, and 90 DAS and also at harvest by using a meter scale from the ground to the top leaf pre-heading, and from the ear head base post-heading.

**Comment [i-[15]:** How you manage different heading at these days? Plant height is one of the growth parameter, so you can take their height in physiological maturity

### Number of tillers(m<sup>-2</sup>)

The number of tillers per plant was recorded at 60 and 90 days post-sowing. This was done by counting the total number of tillers on five randomly selected plants per plot.

### Leaf area index (LAI)

The leaf area index was determined by measuring the leaf area at 30, 60, and 90 days post-sowing. Plants were chosen with a row length of 0.25m, and their green leaves were separated to measure surface area with an automatic leaf area meter (LI-3100C Area Meter (LI-COR Biosciences)).

$$\text{Leaf area index} = \frac{\text{Leaf area}}{\text{Ground area}}$$

### 2.6.3. Economics

#### 2.6.3.1. Cost of cultivation (₹ ha<sup>-1</sup>)

The cost of cultivation was worked out on the basis of input rates at the farm. Treatment's cost was calculated separately. The common cost of cultivation (₹ ha<sup>-1</sup>) was worked out by considering all the expenses incurred in the cultivation and added variable cost due to treatments (including interest of working capital) in order to get total cost of cultivation.

Comment [i-[16]: What about other costs like fertilizer, chemicals, .....

#### 2.6.3.2. Gross return (₹ ha<sup>-1</sup>)

The overall income was determined by multiplying the crop and straw production with the prevailing market rate in various conditions. The total income (₹ /ha) was calculated by adding up the earnings from both the grain and straw harvest.

$$\text{Gross return (₹ ha}^{-1}\text{)} = \text{Total income from the grain and straw harvest}$$

#### 2.6.3.3. Net return (₹ ha<sup>-1</sup>)

Net profit is the outcome received by subtracting the cost of cultivation from gross income (₹ ha<sup>-1</sup>). The net return was worked out by using the following formula:

$$\text{Net return (₹ ha}^{-1}\text{)} = \text{Gross return (₹ ha}^{-1}\text{)} - \text{Cost of cultivation (₹ ha}^{-1}\text{)}$$

#### 2.6.3.4. B: C ratio (₹ ha<sup>-1</sup>):

$$\text{B:C} = \frac{\text{Net return (₹ ha}^{-1}\text{)}}{\text{Cost of cultivation (₹ ha}^{-1}\text{)}}$$

### Statistical analysis

The data collected for various characteristics underwent statistical analysis using Fisher's method of analysis of variance (ANOVA). Critical difference (CD) values were calculated when the 'F' test was found significant at the 5% level. Heatmap clustering using SRPLOT

(<http://www.bioinformatics.com.cn/en?keywords=heatmap>) was employed to visually summarize the data, highlighting key features and their interrelationships.

## RESULT AND DISCUSSION

### Growth attributes

Plant height showed significant growth until 90 days after sowing (DAS), primarily influenced by phosphorus levels starting from 60 DAS. The tallest plants were recorded with 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, followed by 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Row spacing also impacted height, with 25x10 cm resulting in taller plants compared to 20x10 cm. Dry matter accumulation peaked at 90 DAS and harvest, with P<sub>4</sub>, P<sub>2</sub>, and P<sub>3</sub> outperforming P<sub>1</sub> (40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>). Increased row spacing correlated with higher dry matter. Both phosphorus levels and row spacing affected shoot density, with the highest densities observed at 90 DAS and harvest.

The leaf area index rose significantly up to 60 DAS due to phosphorus and spacing (**Table 2**). Overall, growth followed a sigmoid curve, characterized by initial slow growth followed by rapid acceleration until 60 DAS. While phosphorus levels did not significantly affect initial plant population, wider row spacing led to increases (**Singh, 2019; Williams et al., 2021**). Growth rates between 30 and 90 DAS were attributed to improved light absorption, photosynthetic activity, and nutrient uptake (**Styles et al., 2018**).

Phosphorus application significantly enhanced tiller number, plant height, leaf area index, dry matter accumulation, and maturity, with peak values at P<sub>4</sub>. Initial plant population and maturity showed no significant differences among phosphorus levels. Maximum dry matter (1097.25 g m<sup>2</sup>) and height (91.18 cm) were recorded at P<sub>4</sub>. Similarly, increasing row spacing improved initial population, tillers, height, leaf area index, and dry matter, with the highest values at 25 cm x 10 cm spacing (S<sub>3</sub>) (**Sendhil et al., 2019; Qin et al., 2013**).

Overall, both phosphorus levels and row spacing significantly influenced the growth parameters of black wheat. Optimal phosphorus levels and appropriate spacing can enhance black wheat productivity and yield, suggesting effective agronomic practices for farmers.

**Table 2. Effect of different phosphorus levels and row spacing on growth parameters of wheat**

| Treatments | Initial plant population (m <sup>-2</sup> ) | Number of tillers (m <sup>-2</sup> ) | Plant height (cm) | Leaf area index (%) | Dry matter accumulation (g m <sup>-2</sup> ) | Maturity (Days) |
|------------|---|--------------------------------------|-------------------|---------------------|--|-----------------|
|            |   |                                      |                   |                     |  |                 |

Comment [i-17]: Some parameters not included in materials and methods

| Phosphorus (P)                                       |               |               |              |             |                |               |
|--|---------------|---------------|--------------|-------------|----------------|---------------|
| 40 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 181.17        | 370.50        | 80.75        | 4.10        | 971.85         | 118.27        |
| 50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 186.93        | 398.47        | 86.87        | 4.40        | 1045.02        | 122.03        |
| 60 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 189.67        | 406.33        | 88.58        | 4.49        | 1065.90        | 124.03        |
| 70 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | <b>189.33</b> | <b>418.37</b> | <b>91.18</b> | <b>4.62</b> | <b>1097.25</b> | <b>125.10</b> |
| SEd (±)  | 4.94          | 10.31         | 2.28         | 0.10        | 22.49          | 2.56          |
| CD (P= 0.05)   | NS            | 30.24         | 6.69         | 0.32        | 65.97          | NS            |
| Row spacing  |               |               |              |             |                |               |
| 15 cm X 10 cm  | 170.95        | 378.50        | 82.50        | 4.18        | 992.75         | 121.83        |
| 20 cm X 10 cm  | 194.95        | 402.40        | 87.74        | 4.45        | 1055.48        | 122.35        |
| 25 cm X 10 cm  | <b>208.93</b> | <b>414.35</b> | <b>90.30</b> | <b>4.58</b> | <b>1086.79</b> | <b>122.90</b> |
| SEd (±)  | 8.7           | 8.93          | 1.97         | 0.09        | 19.48          | 1.47          |
| CD (P= 0.05)   | 25.7          | 26.19         | 5.79         | 0.27        | 57.13          | NS            |




Fig. 1. Effect of different phosphorus levels and row spacing on growth parameters of wheat

### Yield parameters

The impact of phosphorus levels on wheat spike characteristics was significant, with the highest spike number (316.05) observed at 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> equalling the performance of 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> treatments and surpassing 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Similarly, spike length reached its peak (8.93) with P4, matching P2 and P3 but outperforming P1. Grain number per spike (40.12) was also highest with P4, comparable to P2 and P3. Sowing at 25x10 cm spacing resulted in a higher number of grains per spike (39.75), similar to 20x10 cm spacing. Phosphorus levels significantly influenced grain yield, with the highest yield (33.81 q ha<sup>-1</sup>) achieved at 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> (P4), similar to P2 and P3, and superior to P1. Straw yield followed a similar pattern, with P4 producing 42.84 q ha<sup>-1</sup>, comparable to P2 and P3. Nitrogen content in grain (1.88%) and straw (0.37%) was also highest with P4, showing superiority over lower phosphorus levels (Table 3). Row spacing did not significantly affect test weight but influenced nitrogen content, with higher levels observed at 25x10 cm spacing. The findings underscore the role of phosphorus and row spacing in influencing yield attributes and overall wheat productivity, aligning with previous research (Patra, 2024; Kolarić et al., 2021).

**Table 3.** Effect of different phosphorus levels and row spacing on yield parameters of wheat

| Treatment  | Number of spikes/m <sup>2</sup> | Length of spike (cm) | Grain spike <sup>-1</sup> | Grain yield (qha <sup>-1</sup> ) | Straw yield (qha <sup>-1</sup> ) | Test weight (g) 10000 grain weight | Nitrogen content (%) |             |
|--|---------------------------------|----------------------|---------------------------|----------------------------------|----------------------------------|------------------------------------|----------------------|-------------|
|  |                                 |                      |                           |                                  |                                  |                                    | Grain                | Straw       |
| <b>Phosphorus level</b>                              |                                 |                      |                           |                                  |                                  |                                    |                      |             |
| 40 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 279.95                          | 7.92                 | 35.53                     | 29.95                            | 39.75                            | 36.17                              | 1.67                 | 0.29        |
| 50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 301.02                          | 8.52                 | 30.22                     | 32.22                            | 41.27                            | 36.93                              | 1.80                 | 0.33        |
| 60 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 307.03                          | 8.67                 | 38.97                     | 32.85                            | 41.19                            | 37.20                              | 1.81                 | 0.36        |
| 70 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | <b>316.05</b>                   | <b>8.93</b>          | <b>40.12</b>              | <b>33.81</b>                     | <b>42.84</b>                     | <b>37.50</b>                       | <b>1.88</b>          | <b>0.37</b> |
| <b>SEm(±)</b>  | 7.78                            | 0.19                 | 1.07                      | 0.68                             | 4.11                             | 0.92                               | 0.02                 | 0.004       |
| <b>CD(p=0.05)</b>                                    | 22.83                           | 0.56                 | 3.14                      | 2.00                             | 1.06                             | NS                                 | 0.06                 | 0.012       |
| <b>Row spacing(cm)</b>                               |                                 |                      |                           |                                  |                                  |                                    |                      |             |
| 15 cm X 10 cm  | 285.96                          | 8.09                 | 36.29                     | 30.95                            | 39.63                            | 36.65                              | 1.78                 | 0.33        |
| 20 cm X 10 cm  | 304.01                          | 8.59                 | 38.59                     | 32.54                            | 41.90                            | 37.15                              | 1.80                 | 0.34        |
| 25 cm X 10 cm  | <b>313.06</b>                   | <b>8.85</b>          | <b>39.75</b>              | <b>33.49</b>                     | <b>42.26</b>                     | 37.05                              | 1.84                 | 0.34        |
| <b>SEm±</b>  | 6.74                            | 0.16                 | 0.92                      | 0.59                             | 0.92                             | 0.80                               | 0.01                 | 0.004       |

|               |       |      |      |      |      |    |      |       |
|---------------|-------|------|------|------|------|----|------|-------|
| C.D. (p=0.05) | 19.77 | 0.48 | 2.72 | 1.73 | 2.71 | NS | 0.05 | 0.010 |
|---------------|-------|------|------|------|------|----|------|-------|

#### Nitrogen uptake in grain, straw and total Nitrogen uptake:

Higher nitrogen uptake by grain and straw (63.53 kg ha<sup>-1</sup> and 15.98 kg ha<sup>-1</sup>) was recorded with 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, matching the performance of 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, significantly surpassing 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Similarly, higher total nitrogen uptake (79.51 kg ha<sup>-1</sup>) by grain and straw was observed with P4 (125% RDF), lower (61.68 kg ha<sup>-1</sup>) with P1 (50% RDF + 50% N through FYM), showing superiority over P1, P2, and at par with P3 treatments. Row spacings significantly influenced nitrogen uptake in seed and straw, with higher uptake (61.59 kg ha<sup>-1</sup> and 14.73 kg ha<sup>-1</sup>) observed at 25x10 cm spacing, comparable to 20x10 cm. Phosphorus levels also affected phosphorus content in grain, with the highest content (0.36%) under 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, comparable to P2 and P3, superior to P1 (**Table 4**). Nutrient uptake reflects dry matter accumulation and content in plants, with nitrogen and phosphorus uptake correlating with growth and dry matter production. Potassium uptake was notably influenced by phosphorus levels and row spacings, with nutrient uptake increasing with higher nutrient supply systems, enhancing plant growth and soil nutrient absorption. Row spacings, affecting tiller density, also played a significant role in nutrient uptake efficiency, aligning with previous research (Abichouet *et al.*, 2019; Jat *et al.*, 2021).

**Comment [i-[18]:** Is there another rate common application for nitrogen fertilizer or Its fro 18% of DAP? Please justify accordingly source and levels

**Table 4.** Nitrogen uptake in grain and straw and total nitrogen uptake as influenced by Phosphorus level and row spacing on wheat crop.

| Treatment  | Nitrogen uptake (kg ha <sup>-1</sup> ) |              | Total Nitrogen uptake |
|--|--|--------------|-----------------------|
|  | Grain                                  | Straw        |                       |
| <b>Phosphorus level</b>                              |  |              |                       |
| 40 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 49.94                                  | 11.73        | 61.68                 |
| 50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 51.87                                  | 13.86        | 71.74                 |
| 60 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 61.50                                  | 15.19        | 76.68                 |
| 70 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | <b>63.53</b>                           | <b>15.98</b> | <b>79.51</b>          |
| SEm±   | 1.40                                   | 0.46         | 1.73                  |
| C.D. (P=0.05)  | 4.12                                   | 1.35         | 5.08                  |
| <b>Row spacing (cm)</b>                              |  |              |                       |
| 15 cm X 10 cm  | 54.36                                  | 13.42        | 67.77                 |
| 20 cm X 10 cm  | 58.69                                  | 14.43        | 73.12                 |
| 25 cm X 10 cm  | <b>61.59</b>                           | <b>14.73</b> | <b>76.12</b>          |
| SEm±   | 1.21                                   | 0.39         | 1.50                  |
| C.D. (p=0.05)  | 3.57                                   | 1.16         | 4.40                  |

#### Phosphorus uptake in grain, straw and total Phosphorus uptake:

The data presented in **Table 5** show that the highest total phosphorus uptake (15.24 kg ha<sup>-1</sup>) by grain and straw was recorded with the treatment of 75 kg P/ha (P4), which was at par with the 50 kg P/ha (P2) and 60 kg P/ha (P3) treatments. These treatments demonstrated significant superiority over

the 40 kg P/ha (P1) treatment. Additionally, different row spacings had a significant effect on total phosphorus uptake in both grain and straw. The highest total phosphorus uptake (14.62 kg ha<sup>-1</sup>) was observed with the 25x10 cm row spacing, which was at par with the 20x10 cm spacing.

**Table 5.** Phosphorus uptake in grain and straw and total nitrogen uptake as influenced by Phosphorus level and row spacing on wheat crop.

| Treatment  | Phosphorus uptake (kg ha <sup>-1</sup> ) |       | Total Phosphorus uptake |
|--|--|-------|-------------------------|
|  | Grain                                    | Straw |                         |
| <b>Phosphorus level</b>                              |  |       |                         |
| 40 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 8.94                                     | 2.51  | 11.45                   |
| 50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 11.23                                    | 2.71  | 13.93                   |
| 60 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 12.00                                    | 2.82  | 14.82                   |
| 70 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> | 12.28                                    | 2.95  | 15.24                   |
| SEm±   | 0.25                                     | 0.07  | 0.30                    |
| C.D. (P=0.05)  | 0.65                                     | 0.21  | 0.89                    |
| <b>Row spacing (cm)</b>                              |  |       |                         |
| 15 cm X 10 cm  | 10.28                                    | 2.60  | 12.88                   |
| 20 cm X 10 cm  | 11.27                                    | 2.80  | 14.07                   |
| 25 cm X 10 cm  | 11.78                                    | 2.84  | 14.62                   |
| SEm±   | 0.22                                     | 0.06  | 0.26                    |
| C.D. (p=0.05)  | 0.65                                     | 0.18  | 0.77                    |

### Economics:

The data clearly demonstrates that cultivation costs increased linearly with higher phosphorus levels across all row spacings. The highest cultivation cost (Rs 33,710.09₹ ha<sup>-1</sup>) was incurred with 70 kg P<sub>2</sub> O<sub>5</sub> ha<sup>-1</sup> across all row spacings. Conversely, the treatment combination of 70 kg P<sub>2</sub> O<sub>5</sub> ha<sup>-1</sup> with 25x10 cm row spacing yielded the highest gross return (75,894.50₹ ha<sup>-1</sup>) and net return (45,064.00₹ ha<sup>-1</sup>)(Table 6). This combination also achieved the highest benefit-cost ratio (1.46). In contrast, the combination of higher phosphorus level (70 kg P<sub>2</sub> O<sub>5</sub> ha<sup>-1</sup>) and wider row spacing (25x10 cm) incurred the highest cultivation cost (33,710₹ ha<sup>-1</sup>) due to additional expenses associated with increased phosphorus application and row spacing (Qin *et al.*, 2013).


**Table6.**Economics of various treatment combinations of wheat

| Treatment combination         | Cost of cultivation<br>(₹ ha <sup>-1</sup> ) | Gross return<br>(₹ ha <sup>-1</sup> ) | Net return<br>(₹ ha <sup>-1</sup> ) | B:C ratio |
|-------------------------------|--|---------------------------------------|-------------------------------------|-----------|
| P <sub>1</sub> S <sub>1</sub> | 28864.60                                     | 48416.00                              | 19551.40                            | 0.67      |
| P <sub>2</sub> S <sub>1</sub> | 29255.62                                     | 53650.00                              | 24394.38                            | 0.83      |
| P <sub>3</sub> S <sub>1</sub> | 29121.09                                     | 66901.00                              | 37780.00                            | 1.29      |
| P <sub>4</sub> S <sub>1</sub> | 29580.22                                     | 68260.00                              | 38679.78                            | 1.30      |
| P <sub>1</sub> S <sub>2</sub> | 29489.60                                     | 46555.00                              | 17065.40                            | 0.57      |
| P <sub>2</sub> S <sub>2</sub> | 29880.62                                     | 46955.00                              | 17074.38                            | 0.57      |
| P <sub>3</sub> S <sub>2</sub> | 29746.09                                     | 55507.00                              | 25760.91                            | 0.86      |
| P <sub>4</sub> S <sub>2</sub> | 30105.22                                     | 71425.00                              | 41319.78                            | 1.37      |
| P <sub>1</sub> S <sub>3</sub> | 30114.60                                     | 49127.00                              | 19012.40                            | 0.63      |
| P <sub>2</sub> S <sub>3</sub> | 30505.62                                     | 52489.00                              | 21983.38                            | 0.75      |
| P <sub>3</sub> S <sub>3</sub> | 30830.22                                     | 63805.00                              | 30094.91                            | 0.89      |
| P <sub>4</sub> S <sub>3</sub> | 33710.09                                     | 75894.50                              | 45064.00                            | 1.46      |

Comment [i-[19]: You should incorporate un adjusted and adjusted grain yield and straw yield in your table before cost of cultivation

Heat map clustering

UNDER PEER REVIEW



**Fig. 2.** Clustered heat map of the effect of different treatment combination on growth, yield and economic parameters of wheat; DM= Dry matter accumulation; NS= No. of shoots, NSP=No. of spike/m<sup>2</sup>; IPP= Initial plant population; LAI= Leaf area index; NCG= Nitrogen content in grain; NCS= Nitrogen content in spike; LS= Length of spike; NUS= Nitrogen uptake in spike; NUG= Nitrogen uptake in grain; GY= Grain yield; SY= Straw yield; GS= Grain spike<sup>-1</sup>; TW= Test weight; M= Maturity; PH= Plant height; TNU= Total Nitrogen uptake.

To observe the variation in black wheat parameters and treatment effects, a heat map clustering analysis was performed (Arya *et al.*, 2022). The results illustrate three distinct data groups: plant growth, yield, and economics. This classification is represented by two types of dendrograms: a horizontally oriented plant parameters dendrogram and a vertically oriented treatments dendrogram (Figure 2). In the heat map, traits are visually represented by colors, with blue indicating the maximum value and yellow the minimum.

The plant parameter dendrogram shows two primary clusters (Cluster 1 and Cluster 2). Cluster 1 contains only the trait DM, while Cluster 2 further divides into two sub-clusters (Cluster 3 and Cluster 4). Cluster 3 contains two traits, NS and NSP. Cluster 4 further divides into two sub-clusters (Cluster 5 and Cluster 6). Cluster 5 contains only the trait IPP, while Cluster 6 encompasses all remaining plant parameters.

The treatment dendrogram, positioned vertically, distinctly separates all samples into two main clusters (Cluster 1 and Cluster 2) based on their treatments. The first main cluster further divides into two sub-clusters (Cluster 3 and Cluster 4). Similarly, the second main cluster splits into two sub-clusters (Cluster 5 and Cluster 6). Cluster 3, there are only single treatment (Treatment 1) and another with treatments (Treatment 5). Cluster 5 also divides into two sub-clusters (Cluster 7 and Cluster 8). Cluster 7 includes only Treatment 2, while Cluster 8 includes Treatments 3 and 6. Cluster 6 also divides into two sub-clusters (Cluster 9 and Cluster 10). Cluster 9 includes only Treatment 4, while Cluster 10 includes only Treatment 7.

The analysis of heat maps and dendrograms offers valuable insights into the relationships between plant characteristics and treatment effects. The plant parameter dendrogram groups traits like No. of shoots (NS) and No. of spikes (NSP), highlighting their similarities and importance for the number of shoots and spikes. The treatment dendrogram clusters treatments based on their effects, identifying the most effective and redundant options, which aids in optimizing agricultural practices and reducing costs. Temporal data comparison, distinguishing pre-harvest from post-harvest, provides a deeper understanding of how traits evolve over time, guiding the timing of interventions for maximum benefit. By revealing which traits are influenced similarly by various treatments, we can pinpoint key genetic or physiological pathways, aiding in breeding for desirable traits and stress resistance. This precise visualization simplifies complex data, enabling farmers, agronomists, and researchers to make informed decisions that promote sustainable and effective agricultural practices.

## **Conclusion**

The study highlighted the significant effects of phosphorus levels and plant spacing on black wheat (*Triticum aestivum* L.) growth, yield, nitrogen uptake and economic viability. Results showed that applying 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> notably improved plant growth, yield parameters, and nitrogen uptake. The 25 cm x 10 cm spacing further enhanced these attributes by reducing competition for resources. The highest grain and straw yields, as well as nitrogen uptake, were observed at 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, though 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> proved almost equally effective, suggesting it as a cost-effective strategy. Economically, the combination of 70 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and optimal spacing (15 cm × 10 cm (S1), 20

cm × 10 cm (S2), and 25 cm × 10 cm (S3)) yielded the highest returns, but 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> offered similar benefits, making it a recommended practice for maximizing yield and profit. Thus, strategic phosphorus application and appropriate plant spacing were crucial for sustainable black wheat cultivation, providing a balanced approach to improving crop productivity and economic returns.

## References

1. Abichou, M., De Solan, B. and Andrieu, B. 2019. Architectural response of wheat cultivars to row spacing reveals altered perception of plant density. *Frontiers in Plant Science*, 10: 999.
2. Arya, S., Kumar, R., Prakash, O., Kumar, S., Mahawer, S.K., Chamoli, S., de Oliveira, M.S. 2022. Chemical composition and biological activities of *Hedychium coccineum* Buch.- Ham. ex Sm. essential oils from Kumaun hills of Uttarakhand. *Molecules*, 27(15): 4833.
3. Jackson, M.L. 1973. *Soil Chemical Analysis*, Prentice Hall of India Private Limited, New Delhi,
4. Jat, R. K., Bijarniya, D., Kakraliya, S. K., Sapkota, T. B., Kakraliya, M. and Jat, M. L. 2021. Precision nutrient rates and placement in conservation maize-wheat system: effects on crop productivity, profitability, nutrient-use efficiency, and environmental footprints. *Agronomy*, 11(11): 2320.
5. Joshi, A.K., Mishra, B., Chatrath, R., Ortiz Ferrara, G., Singh, R.P. 2007. Wheat improvement in India: present status, emerging challenges and future prospects. *Euphytica*, 157: 431-446.
6. Khan, F., Siddique, A. B., Shabala, S., Zhou, M. and Zhao, C. 2023. Phosphorus plays key roles in regulating plants physiological responses to abiotic stresses. *Plants*, 12(15): 2861.
7. Kolarić, L., Popović, V., Živanović, L., Ljubičić, N., Stevanović, P., Šarčević, T. and Todosijević, L., and Ikanović, J. 2021. Buckwheat yield traits response as influenced by row spacing, nitrogen, phosphorus, and potassium management. *Agronomy*, 11(12): 2371.
8. Liu, Y., Chen, X. and Wu, J. 2023. Optimizing phosphorus levels in purple wheat cultivation for enhanced anthocyanin retention and yield. *Crop Science Journal*, 65(3): 311-319.
9. Mishra, B., Shorean, J., Chatrath, R., Sharma, A.K., Gupta, R.K., Sharma, R.K., Singh, R., Rane, J., Kumar, A. 2005. Cost effective and sustainable wheat production technology. Directorate of wheat research, Karnal, Technical Bulletin. 8(1).
10. Nguyen, D. H., El-Ramady, H., Llanaj, X., Törös, G., Hajdú, P. and Prokisch, J. 2023. Chemical composition and health attributes of agri-foods: A scientific overview on black foods. *Sustainability*, 15(4): 3852.

11. Olsen, S.R., Cole, C.V., Watanable, F.S., Dean, L.A. 1954. Estimation of available phosphorous in soil by extraction with sodium bicarbonate. *USDA Cric.* 930:19-23.
12. Patra, P. S., Saha, R., Ahmed, A. S., Kanjilal, B., Debnath, M. K., Paramanik, B. and Biswas, A. 2024. Enhancing aromatic rice production through agronomic and nutritional management for improved yield and quality. *Scientific Reports*, 14(1): 15555.
13. Qin, X. L., Weiner, J., Qi, L., Xiong, Y. C. and Li, F. M. 2013. Allometric analysis of the effects of density on reproductive allocation and Harvest Index in 6 varieties of wheat (*Triticum*). *Field Crops Research*, 144: 162-166.
14. Sendhil, R., Kiran Kumar, T.M., Singh, G.P. 2019. *Wheat Production in India: Trends and Prospects. Recent Advances in Grain Crop.*
15. Sharma, P., Sharma, R. K. and Kaur, N. 2021. Phosphorus and nutrient management in quinoa: Effects on growth, yield, and nutrient uptake. *Journal of Plant Nutrition*, 44(12):1783-1793.
16. Singh, A., Brar, S.K., Gandhi, N. 2019. Effect of spacing and different sowing methods on yield of wheat. *Journal of pharmacognosy and phytochemistry*,4:42-44.
17. Styles, D., Adams, P., Thelin, G., Vaneeckhaute, C., Chadwick, D. and Withers, P. J. 2018. Life cycle assessment of biofertilizer production and use compared with conventional liquid digestate management. *Environmental Science and Technology*, 52(13): 7468-7476.
18. Subbiah, B.V., Asija, C.L. 1956. A rapid procedure for the estimation of available N in Soil. *Current Science*, 25:259-60.
19. Walkley, A. and Black, I.A. 1934. Old piper, S.S. soil and plant analysis. *Soil Science*,37(1): 29-38.
20. Weeks, Jr, J. J. and Hettiarachchi, G. M. 2019. A review of the latest in phosphorus fertilizer technology: possibilities and pragmatism. *Journal of Environmental Quality*, 48(5): 1300-1313.
21. Williams, J. J., Dodds, D. M., Buehring, N. W., Dhillon, J. S. and Henry, W. B. 2021. Corn response to row spacing and plant population in the Mid-South United States. *Agronomy Journal*, 113(5): 4132-4141.