

Original Research Article

Assessment of glyphosate Roundup Turbo (450 g/l) on haemato-biochemical indices of freshwater African catfish, *Clarias gariepinus* fingerlings

Abstract

The escalating use of glyphosate-based herbicides in agricultural practices has sparked concerns regarding their potential ecological impact. This study undertakes a comprehensive investigation into the acute ecotoxicological effects of a brand-new glyphosate (Roundup Turbo 450 g/l) on haematology and biochemical indices of *Clarias gariepinus* a commercially vital fish species in Nigeria. Fish were exposed to 0.0 as control, 0.5, 0.7, 0.9 and 1.2 mg/l glyphosate for 96 hours in the laboratory. Water quality parameters monitored were temperature, pH, dissolved oxygen electrical conductivity, nitrate and nitrite. The blood, liver and gills were extracted and analyzed. Haematological and biochemical analyses revealed significant changes ($P < 0.05$) as concentration increases. The decrease in red blood cell, pack cell volume, haemoglobin and white blood cell was indicative of glyphosate-induced hematotoxicity. Biochemical analyses disclosed alterations in enzymatic activities within liver and gill tissues, including significant increase in alanine aminotransaminases, aspartate aminotransferase and alkaline phosphatase, suggesting hepatic and branchial dysfunction and decrease in total protein, globulin and albumin,

showing enhanced disruption in protein synthesis. Glucose and albumin in liver were not detected. Behavioral observations indicated the fish heightened stress response, characterized by erratic swimming patterns, increased agitation, equilibrium loss, hyperactivity and skin erosion. The 96-hour LC₅₀ value for this study was 2.50 mg/l and it highlights glyphosate roundup turbo's potential to exert detrimental effects on *Clarias gariepinus* populations. These findings underscore the imperative for sustainable agricultural practices, judicious pesticide management, and rigorous environmental monitoring to mitigate the ecological risks associated with glyphosate use.

Keywords: Acute study; aquatic toxicology; biochemistry; blood indices; *Clarias gariepinus*; water quality

1. INTRODUCTION

The increase in aquatic pollution caused by industrialization, urbanization and agricultural activities is a serious threat to human and environmental health (Amaeze *et al.*, 2020). Chemicals used by agriculture and industries that find their way to the aquatic ecosystem has lately gained more attentions on the agendas of regulating bodies (Mensah *et al.*, 2014; Brack *et al.*, 2022; Groh *et al.*, 2022). Despite the widespread use of numerous chemicals, insufficient information and understanding exist about these chemical with respect to human toxicity and environmental health risk thus hindering effective management and the establishment of environmental safety guidelines.

Pesticides represent a crucial group of chemicals that can be subject to regulation, and a more precise evaluation of their risk nature could result to major improvements in environmental safety. Pesticides are mixtures of substances used to regulate, prevent or eliminate pests, including weed, nematodes, insects and fungi. Its usage in agricultural fields to control pests

is extremely toxic to non-target organisms like fish and affect fish health through impairment of metabolism, sometimes leading to mortality (Shankar *et al.*, 2013). The increased use of pesticides and other agrochemical to boost crop yield and quality is closely linked to the advancement of agricultural mechanization. Thus, pesticides have become an indispensable tool in large-scale agricultural development in Nigeria (Opute and Oboh, 2021). In addition, drainage and irrigation systems may become polluted through the application of pesticides during agricultural and pest control activities, and this may negatively impact the living and non-living members of the contaminated water (Mohamed *et al.*, 2012). Pesticides includes fungicides, insecticides and herbicides.

The widespread adoption of herbicides in agricultural sectors worldwide serves as a primary mechanism for controlling unwanted plant growth. However, the uncontrolled use of herbicides intended to increase agricultural output can have a **far-reaching** consequences for aquatic ecosystems and organisms that inhabit them. As such, it may become hazardous to the health of man when these aquatic organisms are harvested and consumed (Williams, 2011). There is a growing public concern about the amount of herbicides that have been introduced into the environment by leachate and runoff, not to mention that the contaminations of the aquatic environments generally occur by a mixture of these compounds and not by isolated substances (He *et al.*, 2012). Herbicide employed in this study is glyphosate Roundup turbo. Glyphosate (N-phosphonomethyl glycine) is the active ingredient in *Roundup turbo*, a broad-spectrum, post-emergence, non-selective herbicides used for controlling annual and perennial grasses, broad-based leafed weeds, trees and other species (Okayi *et al.*, 2010). Glyphosate is considered a probable human carcinogen based on scientific evaluation in humans and other laboratory animals (Portier, 2016).

In aquatic organisms, including fish, any kind of waterborne pollution is easily reflected in the circulatory system Ismail *et al.* (2017); therefore, blood parameters are one of the commonest and most important biomarkers for diagnosing the structural and functional status of fish exposed to effluents and pollutants (Faggio *et al.*, 2014; Burgos Aceves *et al.*, 2019). Changes in hematological parameters depend on the magnitude of the impact of contaminant (concentration), the duration of exposure, fish species, age and health status (Alimba *et al.*, 2019). Alterations in white blood cell (WBC) numbers might be regarded as a prognostic tool or an early-warning signal of disturbance in homeostatic defense abilities of fish (Oladokun *et al.* 2020). Udume *et al.* (2022a) noted that increase in ammonia (NH₃) levels was principally responsible for the haematological modification of fish. Alterations in blood biochemical parameters as important diagnostic tool can be used for the detection of abnormalities in the liver and other tissues both in the laboratory and field studies (Banaee *et al.*, 2011; Javed *et al.*, 2016; Adam *et al.*, 2019; Shah and Parveen, 2020)). According to Prabesh *et al.* (2021) Biochemical parameters of golden mahseer were affected by sub-lethal exposure of chlorpyrifos and dichlorvos.

The toxicity of herbicides to fish **relies** on size and species (Noga, 2012). This study employed African catfish, *Clarias gariepinus* as the test species because of its significant economic importance and widespread cultivation in Nigeria and other developing country. *Clarias gariepinus* also referred to as mudfish, is very hardy and tasty, tolerate adverse aquatic conditions where other cultivable fish species cannot survive (NASS, 2010). The choice of fish as a model in ecotoxicological research could be valued as fish serves as a totally sensitive bio-indicator of aquatic infection in tropical regions (Mdegela *et al.*, 2010). The potentiality of

application of the findings from those researches on humans and different environmental health problems has made fish a greater appealing model organism in toxicology research (Govind, 2011).

Toxicity testing of chemicals on animals has been used for a long time to detect the potential hazards posed by chemicals to environment and human (OECD, 2021). The study of these chemical is essential especially in the early life stages of fish because the continuous use could contribute to a decline in the fishery of water. Aquatic Bioassays are necessary in water pollution control to determine whether a potential toxicant is dangerous to aquatic life and if so, to find the relationship between the toxicant concentration and its effect on aquatic animals (USEPA, 2012; Omoniyi, 2018).

In this study, glyphosate a common herbicide with a brand name “*Roundup Turbo*” used by farmers especially in the South-Eastern part of Nigeria is evaluated for its effect of acute exposure in *Clarias gariepinus*.

2. MATERIALS AND METHODS

2.1 Experimental Site

The experiment was conducted in wet laboratory Fisheries Department, College of Natural Resources and Environmental Management, Michael Okpara University of Agriculture Umudike, Abia State in South Eastern Nigeria. The institute is situated in the tropical rainforest zone on Latitude $05^{\circ}26^1 - 5^{\circ}25^1\text{N}$ and Longitude $07^{\circ}34^1 - 7^{\circ}36^1\text{E}$ with total annual rainfall ranging from 1800 to 2100mm and at altitude of 122 m above sea level (NRCRI, 2010). The major activities carried out in this location is farming and pesticides commonly used by farmers is organophosphate chemical such as glyphosate hence the choice of the study area. The primary route of glyphosate entry into the research area is *via* human induced activities specifically indiscriminate use of agrochemical by farmers and other users consequently resulting in soil contamination that eventually drains into nearby water bodies.

2.2 Fish sampling and Acclimation

For this study, 150 fingerlings of *Clarias gariepinus* were sourced from Treasure fish farm in Umuahia, Nigeria. The fingerlings had an average weight of 2.9 ± 1.90 g, a total length of 6.08 ± 0.07 cm, and a standard length of 5.40 ± 0.20 cm. They were transported to the experimental site in plastic containers. Prior to the experiment, the specimens were treated with a prophylactic dose of 0.02% potassium permanganate solution to ensure they were disease-free. The fish were then acclimated in a 500-liter tank at a temperature of 26.5°C , with a pH of 7.4 and dissolved oxygen (DO) of 5.9, for a duration of 14 days. The tank was filled with continuously aerated tap water and maintained in a static system. During the acclimation period, the fish were fed twice daily with a commercial feed that contained 35% crude protein.

2.3 Ethical Approval

Approval was granted by Michael Okpara University of Agriculture Umudike Ethical Committee at the College of Veterinary Medicine to conduct research on pesticides using animal model (MOU/AVM/REC/2024/18). All procedures involving test fishes were carried out with utmost care, adhering to the established ethical principles for animal use in scientific research, as outlined in the EU directive 2010/63/EU.

2.4 Assay chemical

Glyphosate with the trade name “*Roundup turbo*” was used as a test chemical for the study. It was procured as a commercial formulated herbicide at a concentration of 450g/L in a one liter

container in Agro-based shop within Umuahia, Abia State, Nigeria with a production date, 9th January, 2024 and expired date, 10th February, 2027.

2.5 Range finding test (LC₅₀) and Preparation of stock solution

This study used a static renewal bioassay, following ASTM standards (1990) to evaluate acute toxicity. Preliminary studies were carried out to evaluate the definitive concentration range for testing chemical. This was determined following the methods of Sogbanmu *et al.* (2018). Thirty (30) fishes divided into triplicate groups of ten (10), were subjected to varying levels of glyphosate treatment, consisting of a control group (0.0 mg/l) and four treated groups (0.5, 0.7, 0.9 and 1.2 mg/l) dissolved in 40-liter water volume. The fish were not fed during the 96-hour duration (Krishna and Hayashi 2000). Fish mortality was monitored at 24-hour time intervals (24, 48, 72 and 96 hours) and dead fish removed to maintain water quality. The test solutions were replaced every 48 hours with freshly prepared glyphosate solutions to ensure optimal exposure conditions. The median lethal concentration LC₅₀ at 96 hours was calculated and recorded.

2.6 Behavioral responses

Behavioral studies were carried out in the order of U.S.EPA, (2008) and guideline No. 249 of OECD, (2021). Behavioral and formative changes observed were equilibrium status, swimming rate, fin movement and hyperactivity. Experimental tanks were closely monitored for 10 to 12 minutes to allow enough time to accurately assess each specimen.

2.7 Water quality analysis

Water quality monitoring was performed daily (every 24 hours) throughout the study period in the aquaria. Key parameters including dissolved oxygen, pH, temperature, total dissolved solids and electrical conductivity were assessed using a HI-769828 multi-parameter probe. Ammonia, nitrate and nitrite analysis were conducted using NT LABS test kit.

2.8 Haematological Analysis

2.8.1 Blood sampling

Blood samples were extracted from the caudal peduncle of the test organism using 2.5ml syringes and hypodermal needle treated with an anticoagulant ethylenediaminetetraacetic acid (Lewbert 2001; Ada *et al.*, 2012). Composite samples of 3-5 fishes were taken from each replicate in order to obtain sufficient blood for haematological analysis. All samples were properly labeled. The haematological indices were evaluated to determine changes and parameters assessed include: red blood cell (RBC), packed cell volume (PCV), haemoglobin content (HBC), white blood cell (WBC). These haematological indices were assessed simultaneously for each blood specimen using an Automated Haematology Analyser BC-2800 following standard procedures outlined by the manufacturers Shenzhen MINDRAY Bio-Medical Electronics Co., Ltd, China.

The derived hematological indices consisting mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHC) were obtained using the calculation method described by Dacie and Lewis (1984).

$$MCV (fl) = \frac{PCV (\%) \times 10}{RBC \text{ Count in Millions/mm}^3}$$

$$MCH (pg) = \frac{Hb \times \frac{g}{dl} \times 10}{RBC \text{ Count in Millions/mm}^3}$$

$$MCHC (g/dl) = \frac{Hb \times \left(\frac{g}{dl}\right) \times 10}{PCV (\%)}$$

2.9 Determination of serum biochemical indices

The serum was isolated from the fish blood samples via centrifugation at 3500 rpm for 15 minutes, after which the extracted serum was preserved at -20 °C. After 96-hour exposure to glyphosate, serum biochemical parameters were evaluated in control and test fish. Liver and gill tissues were assessed for alanine amino transferase (ALT) and aspartate amino transferase (AST) activity using standardized kits from Randox Laboratories United Kingdom following recommended procedures. Alkaline phosphatase (ALP) was measured according to the method proposed by Bergmeyer *et al.* (1980). This underlying principle depend on the enzymatic action of serum alkaline phosphatase which hydrolyzes phenolphthalein monophosphate to produce phosphoric acid and phenolphthalein. The phenolphthalein then undergoes a colorimetric change to pink at alkaline pH values enabling photometric determination. The glucose levels in liver and gills were measured according to the protocols of Schmidt (1961). Serum total protein was determined according to the protocols of Lowry *et al.* (1951) as modified by Spector (1978), taking Bovine serum albumin (BSA) as a standard. A colorimeter was used to measure the absorbance of the sample relative to a standard at 540 nm. The globulin concentration was determined through subtraction of albumin values from the total protein values. Activities of ALT, AST and ALP in the serum were expressed in international unit per litre (IU/L).

2.10 Data analysis

Data were analyzed using Statistical Packages for Social Sciences (SPSS) version 23 (IBM Corporation, Armonk, NY, USA). The study employed Analysis of Variance (ANOVA) to examine the mean values of hematological and biochemical parameters aiming to detect significances among various chemical exposures and concentrations. Probit analysis was used to calculate the LC₅₀. To further investigate significant differences, Least Square Difference (LSD) and Duncan test was used. The threshold for statistical significance was set at p<0.05 following the guidelines established by (Steel and Torries, 2003).

3. RESULTS

3.1. Acute toxicity effect of glyphosate Roundup Turbo on the experimental fish

Table 1 presented the result of an experiment where fish were exposed to different concentrations of glyphosate in separate tanks. The result revealed a clear dose-response relationship between mortality in *Clarias gariepinus* and exposure of test chemical. The control experiment yielded no mortality during the exposure duration (24-96 hours) demonstrating no toxic effect. However, in 24 hours 2% mortality was observed and at 96 hours 20% mortality occurred at concentration of 0.5 mg/l whereas at concentration of 1.2 mg/l, 22% mortality were observed under 24 hours and 100% mortality occurred under 96 hours. The highest percentage mortality (100%) was seen at concentration 1.2 mg/l of glyphosate followed by 0.9 mg/l which had 80% at 96 hour exposure time. This table therefore suggest that as the concentration of glyphosate increases mortality rate also increases with time.

Table 1: Total mortality of *Clarias gariepinus* exposed to varying glyphosate doses

Conc. (Mg/l)	No of Fish	Exposure		Duration (hours)		Mortality (%)		Mortality (%)	
		24	Mortality (%)	48	Mortality (%)	72	Mortality (%)	96	Mortality (%)
0.0	30	0	0	0	0	0	0	0	0
0.5	30	2	6	2	6	4	10	6	20
0.7	30	5	16	6	20	8	26	10	30
0.9	30	12	40	15	53	20	66	25	80
1.2	30	22	73	26	86	30	100	30	100

3.2 Mean physicochemical parameters of the test concentrations glyphosate Roundup Turbo on *Clarias gariepinus*

Table 2 presented the result of water quality parameters at different concentrations of glyphosate. The temperature of the study remained relatively constant across all concentrations with a slight increase at 0.7 mg/l. The temperature ranged between 25.51 ± 0.51 and 25.85 ± 0.52 showing that the temperature was closely monitored and kept within a range, providing a stable environment for the research. pH decreases gradually as concentration increases but remained within a narrow range (5.24-6.57). Electrical conductivity increases as concentration increases indicating potential increase in ionic substances. The level of Total dissolved solids, ammonia, nitrate and nitrite remained relatively constant cross all concentrations. The result also showed that Dissolved oxygen decreased significantly as concentration increases indicating potential toxic effect to the fish. However, all other parameters showed no significant difference and were within the range of standard quality water (APHA, 2005).

Table 2: Mean water quality parameters of culture medium of *Clarias gariepinus* observed during 96-hour exposure

Parameters	Concentrations (Mg/l)				
	0.0	0.5	0.7	0.9	1.2
Temp (⁰ C)	25.51±0.51	25.60±0.19	25.85±0.52	25.62±0.10	25.49±0.18
pH	6.57±0.18 ^a	6.29±0.15 ^a	6.22±0.12 ^a	5.50±0.20 ^b	5.24±0.40 ^b
DO (mg/l)	5.11±0.21 ^c	4.08±0.18 ^d	3.95±0.26 ^c	3.98±0.20 ^b	3.85±0.35 ^a
TDS	95.83±1.10 ^a	96.00±1.60 ^a	95.54±2.31 ^a	95.17±1.70 ^a	95.08±1.90 ^a
EC (μs)	45.7±0.58 ^c	46.3±0.58 ^c	47.0±2.00 ^c	49.7±0.58 ^b	56.0±1.00 ^a
NH ₃ (mg/l)	0.5±0.17	0.34±0.16	0.32±0.20	0.36±0.18	0.40±0.16
NO ₃ ⁻ (mg/l)	7.10±2.62	7.50±2.47	6.68±2.46	7.50±2.47	7.08±2.58
NO ₂ ⁻ (mg/l)	1.41±0.61	1.41±0.47	1.45±0.41	1.35±0.50	1.41±0.61

Remark: Mean values with different alphabetical superscripts are significantly different (P<0.05)
 Temp – Temperature, pH – Potential Hydrogen, DO – Dissolved oxygen, TDS – Total dissolved solids, EC – Electrical conductivity, NH₃ – Ammonia, NO₃⁻ – Nitrate, NO₂⁻ – Nitrite

3.3 Behavioral changes of test specimen to glyphosate Roundup Turbo exposure

Table 3 presented the behavioral toxicity of glyphosate Roundup to *Clarias gariepinus* at various concentrations based on a 96-hour exposure time. The observed aberrant behavior demonstrated a strong temporal and dose-dependent relationship glyphosate exposure. In the control group (0.0 mg/l), no anomaly was observed from 24-96 hour of exposure. The fish in this group exhibited normal swimming patterns, body and fin movement and normal equilibrium status whereas fish in exposed groups exhibited a range of abnormal behaviors when exposed to test chemical ranging from swimming rapidly and erratically indicating agitation to increased mouth and opercula movement, decline in swimming rate, suggesting respiratory impairment and loss of equilibrium. As exposure progressed, fish produced excess mucus and became increasingly lethargic, settling at the bottom of the container. The severity of these anomalies was high in 0.9 and 1.2 mg/l during the 72 and 96-hour exposure to glyphosate.

Table 3: Behavioral changes of *Clarias gariepinus* subjected to glyphosate Roundup turbo at varying levels

Duration/ Concentration (mg/l)	Swimming rate	Fin movement	Hyperactivity	Skin erosion	Equilibrium status	Uncoordinated movement
24 hours						
0.0	-	-	-	-	-	-
0.5	+	+	++	-	-	+
0.7	+	+	++	-	-	+
0.9	++	++	+++	-	+	+
1.2	++	++	++	+	++	+
48 hours						
0.0	-	-	-	-	-	-
0.5	+	++	+++	+	+	+
0.7	++	++	+++	+	+	++
0.9	++	++	++	+	++	++
1.2	++	++	+	++	++	++
72 hours						
0.0	-	-	-	-	-	-
0.5	++	++	++	+	++	++
0.7	++	++	++	+	+++	++
0.9	+++	+	+	++	+++	++
1.2	+++	-	-	+++	+++	+++
96 hours						
0.0	-	-	-	-	-	-
0.5	++	+++	++	+	+++	++
0.7	++	++	++	+	+++	++
0.9	+++	+++	+	+++	+++	+++
1.2	-	-	-	+	-	-

Key: None = (-), Low = (+), Medium = (++) , High = (+++)

3.4 Haematological parameters of *Clarias gariepinus* during 96-hour exposure to glyphosate Roundup Turbo

Table 4 showed mean value of the haematology of the test fish exposed to acute concentrations of glyphosate for 96-hour. Packed cell volumes (PCV) of the control group (0.0 mg/l) were significantly higher (48.35 ± 0.49) than the exposed group. Toxicant exposure in PCV decrease with increase in concentration of glyphosate except for 0.5 mg/l which was slightly higher than 0.7 mg/l but showed no significant differences. The white blood cell (WBC) of the exposed group were significantly lower than the control. There was no significant difference between 0.5 mg/l and 0.7 mg/l. Also, no significant difference between 0.9 mg/l and 1.2 mg/l. The haemoglobin was highest (25.84) in the control group while it was least (4.54) in the group exposed to highest (1.2 mg/l) concentration of glyphosate roundup turbo. The treated group showed gradual decrease in response to increased level of the toxicant while the ungroup group had the highest red blood cell count (RBC). The platelet (PLT) count exhibited significant difference between the control group and those exposed to glyphosate roundup turbo. A notable uptrend in platelet values was observed when comparing the control to the toxicant-exposed groups. In contrast, significant difference was observed in the erythrocyte indices, specifically mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH) and mean corpuscular haemoglobin concentration (MCHC). Notably, MCV values at concentrations of 0.9 and 1.2 mg/l were markedly higher compared to the control group with a statistical significance at ($P < 0.05$). The highest mean corpuscular haemoglobin value was recorded at concentration of 1.2 mg/l while the control group exhibited higher mean corpuscular haemoglobin concentration (53.32 ± 1.05) compared to the group exposed to glyphosate. The highest value of neutrophils (26.67 ± 0.48) was recorded in the control group followed by 0.5 mg/l (24.54 ± 0.57) and least value (16.45 ± 0.45) at 1.2 mg/l of glyphosate. Lymphocyte values showed a dose dependent increase with increase in concentration. The highest value (89.40 ± 1.02) was notable at 1.2 mg/l and the least value (75.00 ± 0.10) was recorded at 0.0 mg/l (control). The result also showed that there was no detectable presence of monocytes, basophils and eosinophils in the haematological studies. Generally, there were notable differences in the values between the control group and other concentrations indicating a distinct effect of glyphosate on the parameters measured. The haematological profile showed reduction in mean values of PCV, RBC, HB and neutrophils with increasing concentrations while the mean values of WBC, MCV, PLT, MCH, MCHC and Lymphocyte demonstrated an increase suggesting a notable change in the blood parameters.

Table 4: Mean haematological parameters of *Clarias gariepinus* exposed to varying concentrations of glyphosate Roundup Turbo during 96-hour

Parameters	Concentrations Mg/l				
	0.0	0.5	0.7	0.9	1.2
PCV (%V)	48.35±0.49 ^c	22.59±1.20 ^d	19.10±0.37 ^c	17.20±0.91 ^b	12.02±0.60 ^a
WBC (x10 ³ mm ³)	2489.00±26.47 ^c	1816.68±75.39 ^b	1779.32±28.86 ^b	1450.00±68.27 ^a	1398.67±76.10 ^a
HB (g/dl)	25.84±0.40 ^d	7.95±0.52 ^c	7.38±0.07 ^c	6.32±0.12 ^b	4.54±0.72 ^a
RBC (x10 ⁷ /μl)	6.30±0.08 ^e	2.96±0.18 ^d	2.49±0.12 ^c	1.61±0.05 ^b	1.04±0.02 ^a
MCV (fl/cell)	77.00±0.01 ^a	76.60±0.48 ^a	76.97±0.01 ^a	128.08±2.59 ^c	123.50±2.90 ^b
MCH (pg/cell)	40.83±0.78 ^b	27.80±1.03 ^a	26.24±0.68 ^a	37.80±3.49 ^b	47.51±8.26 ^c
MCHC (g/dl)	53.32±1.05 ^{cd}	36.08±1.29 ^{bc}	33.82±1.24 ^{ab}	29.75±2.20 ^a	39.71±6.25 ^c
PLT	101.4±5.04 ^d	105.2±6.10 ^c	109.5±7.04 ^b	112.4±8.20 ^a	113.4±5.04 ^a
Neutrophils	26.67±0.48 ^a	24.54±0.57 ^b	22.24±0.67 ^c	19.45±0.50 ^d	16.45±0.45 ^e
Lymphocyte	75.00±0.10 ^d	76.40±0.50 ^d	79.20±0.85 ^c	84.10±1.00 ^b	89.40±1.02 ^a

Remark: Mean values with different alphabetical superscripts along a column for a parameter are significantly different (P < 0.05)

PCV - Pack Cell Volume, WBC - White Blood Cell, HB - Haemoglobin, RBC - Red Blood Cell, MCV - Mean Corpuscular Volume, MCH - Mean Corpuscular Haemoglobin, MCHC - Mean Corpuscular Haemoglobin Concentration, PLT - Platelet

3.5 Liver and gill biochemical changes in *Clarias gariepinus* exposed to varying concentration of glyphosate Roundup turbo

Table 5 presents the result of glyphosate exposure on various biochemical parameters in liver and gill tissues of fish. The result of the biochemical properties indicated a significant increase in all group exposed to the chemical compared to the control. The highest value (31.53±1.07) of Aspartate amino transferase (AST) enzyme in liver and in gill (23.10±1.72) was recorded at concentration 1.2 mg/l with lowest value in liver (22.33±2.89) and gill (18.50±1.45) observed in the control group. Alanine amino transferase (ALT) also recorded an increase with the highest value in the liver (53.85±0.30) and gills (40.85±1.30) at 1.2mg/l. Alkaline phosphatase (ALP) also showed a significant increase in both gill and tissue of the fish with an increase in glyphosate concentration. The highest value in liver (87.20±0.38) and gill (52.00±0.28) was recorded at 1.2 mg/l and lowest value in liver (57.20±0.08) and gill (44.00±1.20) was recorded at 0.0 mg/l (control group). Total protein, globulin and albumin decreased in values with increase in the levels of glyphosate roundup turbo. Non-significant difference (P>0.05) was observed in gills of glucose and albumin as it was not detected in the study. In general, these results showed significant differences in all the biochemical parameters except glucose and albumin in gills among different concentrations of glyphosate (P < 0.05).

Table 5: Mean liver and gill biochemical changes in *Clarias gariepinus* exposed to varying concentration of glyphosate Roundup during 96-hour

Parameters	Tissue	Concentration of glyphosate Mg/l				
		0.0	0.5	0.7	0.9	1.2
AST (u/l)	Liver	22.33±2.89 ^d	25.23±1.50 ^c	25.90±0.28 ^c	28.03±1.89 ^b	31.53±1.07 ^a
	Gill	18.50±1.45 ^d	19.05±1.20 ^d	21.50±2.18 ^c	25.50±1.90 ^a	23.10±1.72 ^b
ALT (u/l)	Liver	40.20±0.45 ^c	44.51±0.30 ^b	49.90±1.00 ^a	52.10±0.85 ^a	53.85±0.30 ^a
	Gill	25.00±1.50 ^b	38.55±2.04 ^a	39.40±1.05 ^a	40.10±2.15 ^a	40.85±1.30 ^a
ALP (u/l)	Liver	57.20±0.08 ^b	85.50±0.40 ^a	86.20±0.08 ^a	86.90±0.36 ^a	87.20±0.38 ^a
	Gill	44.00±1.20 ^b	50.60±1.31 ^a	51.01±1.24 ^a	51.82±1.47 ^a	52.00±0.28 ^a
Total protein	Liver	25.14±0.02 ^a	18.20±0.02 ^b	16.14±0.02 ^b	15.41±0.04 ^c	11.55±0.05 ^d
	Gill	4.90±0.45 ^a	3.00±0.90 ^b	2.45±0.04 ^c	1.80±0.30 ^d	1.05±0.43 ^d
Globulin (g/dl)	Liver	5.00±0.02 ^a	3.51±1.22 ^b	2.1±0.02 ^c	2.20±0.05 ^c	1.00±0.09 ^d
	Gill	4.20±1.12 ^a	2.40±1.02 ^b	2.10±1.14 ^b	1.20±1.50 ^c	1.00±1.40 ^c
Glucose (ug/l)	Liver	28.10±0.59 ^c	30.70±0.28 ^c	35.50±0.28 ^a	34.00±0.28 ^b	36.70±0.28 ^a
	Gill	0.00	0.00	0.00	0.00	0.00
Albumin (g/dl)	Liver	21.01±0.03 ^a	19.10±0.02 ^b	18.40±0.09 ^b	15.09±0.12 ^c	12.52±0.20 ^d
	Gill	0.00	0.00	0.00	0.00	0.00

Remark: Mean values with different alphabetical superscripts along a column for a parameter are significantly different ($P < 0.05$) AST – Aspartate amino transferase, ALT – Alanine amino transferase, ALP – Alkaline phosphatase

4. Discussion

Evaluation of acute toxicity provides preliminary understanding of the adverse effects of emerging contaminants, facilitating the establishment of concentration limits for further research into sub-lethal effects (Majumder *et al.*, 2019). Based on our findings the 96-hour LC₅₀ of glyphosate Roundup turbo recorded in this study was 2.50 mg/l. However, the 96-hour LC₅₀ of glyphosate “Forceup” was 1.5 mg/l in *Clarias* species (Ifeoma *et al.*, 2023). The observed differences could be as a result of formulation of the toxicants and age of the fish used. The calculated 96-hour LC₅₀ value reported in this study was higher than the value previously documented for 0.76 mg/l for *Lates calcarifer* Thanomsit *et al.* (2016) and 0.0025 mg/L in zebra fish Wang *et al.* (2023).

In contrast, LC₅₀ of the present study was lower when compared to the findings of Nwani *et al.* (2013) who reported 96-hour LC₅₀ for *Tilapia zilli* at 211.80 mg/l of glyphosate, *Cyprinus carpio* at 86 mg/l (Deivasigamani, 2015) and *C. punctatus* at 5.93 mg/l Kumar *et al.* (2009). The observed variations in the recorded values may be influenced by a range of factors, ambient water temperature, species-specific traits and the developmental stage and size of the fish

(Somdare, 2016). Furthermore, Findik *et al.* (2001) and Osioma *et al.* (2021) observed that differences in LC₅₀ values can be attributed to variations in fish physiology, environmental parameters, water chemistry and the testing methodologies employed.

Acute toxicity data provides valuable insights into understanding the mechanism of action of substance and its potential harm to non-target species. This information is particularly useful for comparing the effects of different chemicals at various doses (Santos *et al.*, 2010). During this study, herbicide dosage increase resulted to corresponding increase in mortality. Our findings is also in agreement with Fishel *et al.* (2013) and Ali and Muhammad (2016) who reported that percentage mortality of *Clarias gariepinus* juvenile increases with increase in concentration of glyphosate. The introduction of pollutants into aquatic environments alters various water parameters such as temperature, pH, alkalinity, and dissolved oxygen (Fagbenro, 2002; Olufayo, 2009). Water temperature of the study ranged from 25.49 to 25.60°C. This is in line with the report from Britz and Hecht (1987) for *Clarias gariepinus* juveniles with optimum temperature range 25°C to 33°C. In this study, pH level and dissolved oxygen in the treated groups test media were significantly reduced ($P < 0.05$), possibly due to glyphosate's oxidative effects on oxygen molecules. This result is constant with the findings of Ololade and Oginni (2010) who investigated the impact of nickel on *Clarias gariepinus*. Ammonia (NH₄), nitrate (NO₃) and nitrite (NO₂) measured during the study could be as a result of uneaten feed, faecal matter and the test chemical used. This finding is in agreement with (keremi *et al.*, 2014).

Observations of fish behaviour following exposure to toxic substances serve as a good indicator for their internal physiological condition, making it a valuable tool for assessing the ecological effects of aquatic pollutant (Simakani *et al.*, 2018). In this study, *Clarias gariepinus* exposed to acute concentrations at 0.5, 0.7, 0.9 and 1.2 mg/l of glyphosate for 96-hours showed increased anomalous behaviors like loss of equilibrium, abnormal swimming, uncoordinated movement, skin erosion and hyperactivity with decreased aggressive behavior when compared to the control group. These signs manifested before death occurred because of physiological reaction from acute glyphosate toxicity. Similar observations have been found in Nile tilapia Fouad *et al.* (2022) and common carp Fallah (2022) after acute exposure to fungicides. Also, Ayanda *et al.* (2017) and Lanzarin *et al.* (2020) who exposed *Clarias gariepinus* and Zebra fish models respectively to acute concentration of glyphosate. Fish exhibit an increased secretion of mucus which is a characteristics response to stress whether induced by environmental factor or diseases. The production of excess mucus in this study was possibly due to the toxicant introduced in the aquatic media as the skin is one of the primary organs to come into contact with aquatic contaminants. The increased mucus secretion can be seen as a natural defensive mechanism. This protective layer forms a barrier between the fish's body and the contaminated environment, mitigating the effects and ultimately leading to the removal of excess epidermal mucus (Ibrahem 2012). More so, the abnormal behaviors caused by glyphosate exposure could be related to the release of its toxic metabolites aminomethylphosphonic acide (AMPA) which may contribute to disrupting brain functions causing neurotoxicity (Chung and Wong, 2022; Campanale *et al.*, 2023). In this regard, loss of equilibrium and lateral swimming are most likely caused by nervous system dysfunction (Sharma, 2019). This investigation also noted fin erosion in fish exposed to elevated acute levels of glyphosate-based Roundup turbo. This phenomenon may be attributed to COX-1 inhibition, triggering the release of endothelin-1, a strong vasoconstrictor potentially causing epithelial cell degeneration and erosion. Studies by various researchers Matozzo *et al.* (2020) and Peillex and Pelletier (2020) reported similar findings in fish models exposed to xenobiotics.

The analysis of blood serves as a vital tool for evaluating animal health, as haematological markers reflect the body's response to environmental change, influencing energy level, respiratory function and immune defense (Iheanacho and Odo, 2020; Melefa, 2020, Onah, 2020). The presence of toxic chemicals in aquatic environments results in significant hematopoietic disturbances (Ahmed and Zakiya, 2022). In this present study, RBC, HB, PCV and WBC decreased significantly $P < 0.05$ during acute exposures to glyphosate roundup when compared to the control. This decline suggests that glyphosate exposure in water may have triggered anaemia in the experimental fish. The observed decrease in PCV may be linked to gill dysfunction and osmoregulatory imbalance, and haemodilution caused by toxic effects. Our findings are similar with previous studies by Udume *et al.* (2022b) who reported anemic condition on sub-lethal dose of glyphosate 360 g/l exposed to African catfish, Maurya *et al.* (2019) in *Heteropneustes fossilis* juveniles exposed to industrial wastewater pesticides and Udume *et al.*, (2022a) *Clarias gariepinus* exposed to varying levels of acute ammonia (NH₃). Furthermore, the decrease in haemoglobin concentration implies impaired oxygen delivery to tissues leading to respiratory stress and reduced physical performance (Amaeze *et al.*, 2020). Research suggests that pesticide exposure can disrupt erythropoiesis, leading to decreased red blood cell production and haemoglobin synthesis, ultimately resulting in lower red blood cell counts and Hb levels (Mostakim *et al.*, 2015). More so, pesticide-derived reactive oxygen species (ROS) can cause oxidative damage to red blood cells and haemoglobin, impairing oxygen (Lutnicka *et al.*, 2016). The observed decrease in white blood cell count among treatment groups aligns with Adeyemo *et al.* (2007) findings, where *Clarias gariepinus* exposure yielded similar result. This phenomenon was further supported by Olanike (2007) who attributed the decline to stress-induced epinephrine release, leading to leucocyte depletion and impaired immune function. Blood cell indices such as mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH) and mean corpuscular haemoglobin concentration (MCHC) values provides useful information on the cause of anemic condition. Hence the fluctuations in MCV, MCH and MCHC are highly dependent on the values from RBC count, haemoglobin concentration and Pack Cell Volume. This study showed fluctuations in MCV, MCH and MCHC. The observed effect is linked to glyphosate exposure in fish as the data does not exhibit a specific pattern. These fluctuations could be due to deficiency in iron, B₁₂, and folic acid. Our findings are in agreement with Rao (2010), Hossain *et al.* (2015) and Salam *et al.* (2015) who reported that the values of blood cell indices were fluctuated in common carp and some other freshwater fish when they were exposed to acute toxic level of pesticides. Similar findings have been reported in studies involving Nile tilapia and *Clarias gariepinus*, where exposure to mancozeb and atrazine yielded comparable result (Ibrahim *et al.*, 2023; Kanu *et al.*, 2023)

Amino transferases (AST and ALT) play a major role in protein metabolism due to their capacity to facilitate transamination reactions between amino acids and keto acids, ultimately contributing to protein synthesis (Ugbomeh *et al.*, 2019). Moreover, AST and ALT integrate protein, carbohydrate, and fat metabolism with the tricarboxylic acid cycle, particularly in response to physiological, pathological, or environmentally induced stress (Ronda *et al.*, 2019). Alterations in AST and ALT activities have been widely employed as biomarkers for detecting aquatic contamination and toxicant exposure in fish (Hedayati *et al.*, 2010; Ibrahim, 2019; Nayak *et al.*, 2019). Research by Tchounwou *et al.* (2015) and Mostakim *et al.* (2015) highlights the significance of AST and ALT enzymes as sensitive indicators of hepatotoxicity and tissue damage in fish. Specifically, these enzymes are crucial markers for liver, kidney, muscle, and gill damage. The present study evaluated ALT, AST, and ALP enzymatic activities (liver and gills)

and noted an increase in response to acute glyphosate roundup turbo exposure. Our findings also suggested that exposure to the toxicant severely impacted these tissues by its reactive oxygen system (ROS) resulting in cellular dysfunction and the subsequent release of biomarkers into the circulatory system. This finding is consistent with previous research (Pesce *et al.*, 2008), which demonstrated a direct relationship between toxicant concentration and enzymatic variation in fish. Similar findings by Sakr and Al-amoudi (2012); Sakr *et al.* (2013) were observed following melathionin exposure in albino mice and rats. The gills, essential for fish respiration, exhibit early signs of aquatic toxicity, including respiratory distress (Ortiz *et al.*, 2003). In this study, *Clarias gariepinus* exposed to glyphosate roundup turbo showed AST and ALT activities increased significantly in their gills compared to controls. This **increases** likely results from stress caused by the absorption of anthropogenic substances through epithelial cells on the gill surface (Monferran *et al.*, 2005). Additionally, the increase in liver during the study could suggest the generation of keto acids such as keto glutarate and oxaloacetate for energy production necessary to meet the energy demand in response to the stress induced by the herbicide (Varadharajan, 2010). This investigation revealed a marked increase in alkaline phosphatase (ALP) activity in both hepatic and branchial tissues of fish exposed to glyphosate roundup turbo, relative to the control group. These findings align with previous research Shaalan *et al.* (2024); Miri and Khandan (2017), which similarly documented increased ALP activity. The observed upsurge in ALP activity may be attributed to cellular damage or alterations (Khattab, 2007). Furthermore, the heightened ALP levels in the liver could be a response to stress triggered by hepatobiliary cells during detoxification processes. The primary energy source for organisms is carbohydrates, which are converted into glucose to fuel brain and muscle function (Holcombe *et al.*, 1982). However, stress exposure disrupts carbohydrate metabolism to accommodate shifting energy requirements (Shivaraj and Asiya, 2018). This study's findings suggest that exposure to glyphosate roundup turbo impairs carbohydrate metabolism in *Clarias gariepinus*. Similar to other animals, fish store glucose as glycogen in various tissues, including the liver, skeletal muscle, myocardium, and brain (Raje, 2018). Notably, this study observed a significant elevation in hepatic glucose levels (28.10-36.70 µg/l), potentially indicating stress triggered by glyphosate exposure. This glucose increase aligns with established responses of fish to acute and sub-lethal pollutant effects, characterized by increased glucose levels (Luskova *et al.*, 2002). However, glucose was not detected in the gill. An organism's physiological state can be inferred from its protein metabolic status (Magar and Shaikh, 2012). This study observed a significant decline in total protein levels (25.14 – 11.55) in both gill and liver tissues. This finding aligns with previous research documenting protein depletion in *Clarias gariepinus* exposed to primextra (Nwani *et al.*, 2014) and diazepam (Ogueji *et al.*, 2017). The decreased protein content suggests disrupted protein synthesis or enhanced degradation into amino acids, potentially triggered by glyphosate roundup turbo-induced stress. Stress-induced protein catabolism may funnel amino acids into the tricarboxylic acid cycle for energy production (Naveed *et al.*, 2010; Ganeshwade, 2012). Hepatic impairment due to toxicant stress can also lead to protein depletion (Dogan and Can, 2011). Furthermore, cellular necrosis may have contributed to the decline in protein levels, disrupting protein synthesis mechanisms (Rajput *et al.*, 2012).

Conclusion

The study used a brand-new chemical glyphosate *Roundup* turbo 450g/l. The 96-hour LC₅₀ of the study was 2.50 mg/l. Our findings showed increased mortality rate of *Clarias gariepinus* was dose-dependent. It revealed that the chemical affected the liver and gill biochemistry, behavior and haematological parameters of *Clarias gariepinus*. Blood indices showed significant decline in erythrocyte count, hematocrit, hemoglobin levels, and leukocyte count suggesting that the chemical was toxic to the fish. Biochemical analyses indicated alterations in enzymatic activities within liver and gill tissues, including elevations in transaminases (ALT and AST) and alkaline phosphatase (ALP), suggestive of branchial and hepatic malfunction. The current investigation revealed a range of morphological and behavioral alterations in exposed specimens, characterized by erratic swimming, loss of balance, mucus secretions, hanging on the surface and inactivity. Water quality was affected due to the increase in the concentration of the toxicant. Generally, the result suggested that the presence of glyphosate roundup turbo is toxic to the health of *Clarias gariepinus*. Hence, we recommend integration of eco-friendly agricultural practices to minimize pesticide application and development of effective pesticide remediation technologies such as glyphosate remediation using bacteria and enzymes.

Ethical Approval

The authors hereby declare that the experimental procedures were approved by Michael Okpara University of Agriculture Umudike Ethical Committee (MOUAU/CVM/REC/202418). All procedures involving test fishes were carried out with utmost care, adhering to the established ethical principles for animal use in scientific research, as outlined in the EU directive 2010/63/EU on the protection of animals against cruelty.

Declaration of Competing Interest

The authors declare no competing interest.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

References

1. Ada, F.B., Ayotunde, O. and Offem, B. O. Surface and ground waters concentration of metal elements in central Cross River State, Nigeria and their suitability for fish culture, *Int. J. Environ. Sustain.* 2012; 1 (2) 9–20, <http://sciencetarget.com/journal/index.php/IJES>.
2. Adam, M. A., Maftuch, M., Kilawati, Y. and Risjani, Y. The effect of cadmium exposure on the cytoskeleton and morphology of the gill chloride cells in juvenile mosquito fish (*Gambusia affinis*). *Egy. J. Aquat. Res.* 2019; 45, 337–343. <https://doi.org/10.1016/j.ejar.2019.11.011>
3. Adeyemo, O., Ajani, F., and Ajiboye, O. Acute toxicity and blood profile of adult *Clarias gariepinus* exposed to lead nitrate. *The Internet Journal of Hematology*, 2007; 4 (2)

4. Al-Akel, A. S., Alkahem-Al-Balawi, H. F., Al-Misned, F., Mahboob, S., Ahmad, Z. and Suliman, E. M. Effects of dietary copper exposure on accumulation, growth, and 95 haematological parameters in *Cyprinus carpio*. *Toxicology and Environmental Chemistry*, 2010; 92: 1865 – 1878.
5. Ali, S. and Muhammad, K. I. Acute toxicity of herbicide (glyphosate) in *Clarias gariepinus* juveniles. *Toxicology Reports* 2016; (3) 513–515
<http://dx.doi.org/10.1016/j.toxrep2016.05.004>
6. Alimba, C. G., Adekoya, K. O. and Soyinka, O. O. Exposure to effluent from pharmaceutical industry induced cytogenotoxicity, hematological and histopathological alterations in *Clarias gariepinus* (Burchell, 1822). *EXCLI J.* 2019; 18, 63–78.
7. Alkahem Al-Balawi, H. F., Ahmad, Z., Al-Akel, A. S., Al-Misned, F., Suliman, E. M. and Al-Ghanim, K. A. Toxicity bioassay of lead acetate and effects of sub-lethal exposure on growth, haematological parameters and reproduction in *Clarias gariepinus*. *African Journal Biotechnology*, 2011; 10: 11039 – 11047.
8. Amaeze, N. H., Komolafe, B. O., Salako, A. F., Akagha, K. K., Briggs, T. D., Olatinwo, O. O., and Abiodun, M. F. Comparative assessment of the acute toxicity, Haematological and Genotoxic effects of ten commonly used pesticides on the African Catfish, *Clarias gariepinus*. *Heliyon Journal*, 2020; 6(8):47-68
9. American Public Health Association. Standard methods for the examination of water and wastewater, 21st edition. American Public Health Association, Washington, DC 2005.
10. American Society for Testing of Materials. Method: Guide for conducting acute toxicity test with fishes, macro invertebrates and amphibians. 1990; 729-90
11. Ayanda, O. I., Oniyeb, S. J., Auta, J. A. and Ajibola, V. O. Behavioural and some physiological assessment of glyphosate and paraquat toxicity of juveniles of African catfish *Clarias gariepinus*. *Pak. J. Zool.* 2017; 49(1):183-190.
12. Ayoola, S. O and Ajani, E. K. Histopathological effect of cypermethrin on juvenile nile tilapia (*Oreochromis niloticus*). *African Journal of Livestock Extension*, 2007; 5: 1– 13.
13. Banaee, M., Sureda, A., Mirvaghefi, A. R. and Ahmadi, K. Effects of diazinon on biochemical parameters of blood in rainbow trout (*Oncorhynchus mykiss*). *Pesticide Biochemistry and Physiology*, 2011; 99, 1-6. doi:10.1016/j.pestbp.2010.09.001
14. Bergmeyer, H. U., Horder, M. and Rej, R. International federation of clinical chemistry (IFCC) scientific committee *Journal of Clinical Chemistry*, 1985; 24 pp. 481-495
15. Brack, W., Culleres, D. B., Boxall, A., Budzinski, H., Castiglioni, S., Covaci, A., Dulio, V., Escher, B. I., Fantke, P., Kandje, F., Fatta-Kassinos, D., Hernández, F. J., Hilscherová, K., Hollender, J., Hollert, H., Jahnke, A., Kasprzyk-Hordern, B., Khan, S. J., Kortenkamp, A., Kümmerer, K., Lalonde, B., Lamoree, M. H., Levi, Y., Martín, P. A. L., Montagner, C. C., Mougín, C., Msagati, T., Oehlmann, J., Posthuma, L., Reid, M., Reinhard, M., Richardson, S. D., Rostkowski, P., Schymanski, E., Schneider, F., Slobodnik, J., Shibata, Y., Snyder, S. A., Sodr , F. F., Teodorovic, I., Thomas, K. V., Umbuzeiro, G. A., Viet, P. H., Yew-Hoong, K. G., Zhang, X., Zuccato, E. One Planet:

- one health. A call to support the initiative on a global science – policy body on chemicals and waste. *Springer Journal of Environmental Sciences Europe*. vol. 2022; (32) 1, 1-10
16. Britz, P. J. and Hecht, T. Temperature Preference and optimum temperature for growth of African catfish (*Clarias gariepinus*) larvae and post larvae. *Aquaculture*, 63: 205 – 214.
 17. Burgos Aceves, M. A., Lionetti, L. and Faggio, C. 2019. Multidisciplinary hematology as prognostic device in environmental and xenobiotic stress-induced response in fish. *Sci. Total Environ.* 1987; 670, 1170–1183.
 18. Campanale, C., Triozzi, M., Ragonese, A., Losacco, D. and Massarelli, C. Dithiocarbamates: Properties, methodological approaches and challenges to their control. *Toxics*. 2023; 11(10):85
 19. Chung, S. W. C. and Wong, W. W. K. Chromatographic analysis of dithiocarbamate residues and their metabolites in foods employed in dietary exposure studies—A review. *Food Addit Contam-Part A*. 2022; 39:1731–43.
 20. Dacie, J. V., and Lewis, S. M. Practical hematology. 6th ed. New York, London, 1984.
 21. Deivasigamani, S. Effects of herbicides on fish and histological evaluation of common carp (*Cyprinus carpio*). *Int J Appl Res*. 2015; 1(7):437-440.
 22. Dogan, D. and Can, C. Hematological, biochemical and behavioral responses of *Oncorhynchus mykiss* to dimethoate. *Fish Physiology and Biochemistry*. 2011; 37: 951 – 958.
 23. Ezike, C. O. Acute toxicity and haematology of *Clarias gariepinus* exposed to 2, 2 dichlorovinyl dimethyl phosphate (dichlorvos). *International Journal of Fisheries and Aquaculture Studies*, 2017; 5(5):100–105.
 24. Fagbenro, O. A. Tilapia: Fish for Thought, Inaugural Lecture Series 32. Delivered at Federal University of Technology, Akure, 2002.
 25. Faggio, C., Fedele, G., Arfuso, F., Panzera, M. and Fazio, F. Haematological and biochemical response of *Mugil cephalus* after acclimation to captivity. *Cah. Biol. Mar.* 2014; 55, 31–36.
 26. Fallah, F. and Omidzahir, S. Investigation of lethal range concentration and median lethal concentration of hexaflumuron in common carp fishes (*Cyprinus carpio*). *Aquac Sci*. 2022; 10:165–77
 27. Fındık, Ö., Taşdemir, M., Çevik, F. and Göksu, M. Z. L. 2-4 D'nin *C. carpio* (L., 1758) *O. niloticus* (L., 1754) Yavruları Üzerine Akut Toksik Etkisinin Araştırılması. XI. Ul. Su Ür. Semp. Bil., 4-6 Eylül. Sinop: 2001; 492 – 497.
 28. Fishel, F., Ferrell, J., MacDonald, G. and Sellers, B. Herbicides: how toxic are they? (PI-133) 2013.
 29. Fouad, M. R., El-Aswad, A. F. and Aly, M. I. Acute toxicity, biochemical and histological of fenitrothion and thiobencarb on fish Nile tilapia (*Oreochromis niloticus*). *Nus Biosci*. 2022; 14-54.
 30. Ganeshwade, R. M. Biochemical changes induced by dimethoate (Rogo 30% EC) in the gills of freshwater fish *Puntius ticto* (Hamilton). *Journal of Ecology and the Natural Environment*, 2012; 4: 181 – 185.
 31. Gallardo, K., Verbel, J. and Freeman, J. Toxicogenomics to evaluate endocrine disrupting effects of environmental chemicals using the zebra fish model. *Current Genomics*. 2016; 17: 515 527.

32. Geber, G. and Schweizer, D. Cytochemical heterochromatin differentiation in *Sinapis alba* (Cruciferae) using a simple air-drying technique for producing chromosome spreads. 1988.
33. Glusczak, L., Loro, V. L., Pretto, A., Moraes, B. S., Raabe, A., Duarte, M. F., da Fonseca, M. B., de Menezes, C. C. and Valladao, D. M. D. Acute exposure to glyphosate herbicide affects oxidative parameters in Piava (*Leporinus obtusidens*). *Arch. Environ. Contam. Toxicol.* 2011; 61: 624–630.
34. Govind and Pandey. A review on fish model in experimental pharmacology. *International Resources Journal of Pharmacology*, 2011; 2 (9), pp. 33-36.
35. Groh, k., Colette, V. B., Kristin, S. and Ahmed, T. Anthropogenic Chemical as Underestimated Drivers of Biodiversity Loss: Scientific and Societal Implications. *Journal of Environmental Science Technology*, 2022; Vol. (56), 2, 707-710. <https://doi.org/10.1021/acs.est.1c08399>
36. Grzegorz, G., Magdalena, S., Elzbieta, N., Jacek, K. and Mirosława, S. M. Influence of short and long term exposition to Round up on the spontaneous and hormonally stimulated LH secretion and gills condition in prussian carp (*Carassius gibelio*). *Advances in Environmental Sciences International Journal of the Bioflux Society*, 2012; 4(3): 146-152.
37. He, H., Yu, J., Chen, G., Li, W., He, J., and Li, H. Acute toxicity of Butachlor and Atrazine to freshwater green alga *Scenedesmus obliquus* and cladoceran *Daphnia carinata*. *Ecotoxicology Environmental Safety*, 2012; 80:91-96.
38. Hedayati, A., Safahieh, A., Savari, A. and Ghofleh, M. J. Assessment of aminotransferase enzymes in Yellow fin sea bream under experimental condition as biomarkers of mercury pollution. *World Journal of Fish and Marine Sciences*, 2010; 2(3): 186 – 192.
39. Holcombe, G. W., Phipps, G. L. and Tanner, D. K. The acute toxicity of Kelthane, dursban, disulfoton, pydrin and permethrin to fathead minnows, *Pimephales promelas* and rainbow trout, *Salmo gairdneri*. *Environmental Pollution*, 1982; 29: 167 – 178.
40. Hossain, S., Khatun, M. H., Rahman, M. K. and Shahjahan, M. Impacts of Sumithion on blood glucose and some hematological parameters in common carp, *Int. J. Environ.* 2015; (5)8–13.
41. Hued, A. C., Oberhofer, S. and Bistoni, M. D. Exposure to a commercial glyphosate formulation (Roundup®) alters normal gill and liver histology and affects male sexual activity of *Jenynsia multidentata* (Anablepidae, Cyprinodontiformes). *Arch. Environ. Contam. Toxicol.* 2012; 62 (1): 107–117.
42. Ibrahim, M. D. Experimental exposure of African catfish *Clarias Gariepinus* (Burchell, 1822) to phenol: Clinical evaluation, tissue alterations and residue assessment. *Journal of Advanced Research*. 2012; 3: 177 – 183.
43. Ibrahim, R. E., Elbealy, M. A., Salem, G. A., Abdelwarith, A. A., Younis, E. M., Wagih, E., Elkady, A. A., Davies, S. J. and Abdel Rahman, A. N. Acute mancozeb-fungicide exposure induces neuro-ethology disruption, health disorders, and immune oxidative dysfunction in Nile tilapia (*Oreochromis niloticus*). *Aquat Toxicol*, 2023; 261:106630.
44. Ibrahim, R. E., El-Houseiny, W., Behairy, A., Mansour, M. F. and Abd-Elhakim, Y. M. Ameliorative effects of *Moringa oleifera* seeds and leaves on chlorpyrifos-induced growth retardation, immune suppression, oxidative stress, and DNA damage in *Oreochromis niloticus*. *Aquaculture*. 2019; 505:225–34.

45. Ifeoma, C Edeh., Charity, I. Nsofo., Stanley, C. Iheanacho., Chukwumaije, C. Ikechukwu., Ijem, A. Nnachi and Chidimma, G. Nwankwo. Behavioural and Haematological Alterations in the African Catfish (*Clarias gariepinus*) exposed to varying Concentrations of glyphosate. *Asian J. Fish. Aqu. Res.*, 2023; vol. 21, no. 1, pp. 32-42
46. Iheanacho, S. C. and Odo, G. E. Neurotoxicity, oxidative stress biomarkers and haematological responses in African catfish (*Clarias gariepinus*) exposed to polyvinyl chloride microparticles. *Comp Biochem Physiol.* 2020; 232:108741.
47. Ismail, N. M., Ali, S. E. and Mohamed, I. K. Biochemical and histological biomarkers approaches in the assessment of the water pollution in some lined and unlined watercourses of Egypt. *International Journal of Fisheries and Aquatic Studies*, 2017; 5(3): 288-296.
48. Javed, M., Ahmad, I., Usmani, N. and Ahmad, M. Studies on biomarkers of oxidative stress and associated genotoxicity and histopathology in *Channa punctatus* from heavy metal polluted canal. *Chemosphere* 2016; 151, 210–219. <https://doi.org/10.1016/j.chemosphere.2016.02.080>
49. Kanu, K. C., Okoboshi, A. C. and Otitolaju, A. A. Haematological and biochemical toxicity in freshwater fish *Clarias gariepinus* and *Oreochromis niloticus* following pulse exposure to atrazine, mancozeb, chlorpyrifos, lambda-cyhalothrin, and their combination. *Comp Biochem Physiol C Toxicol Pharmacol.* 2023; 1:109643.
50. Keremi, R. I., Davies, O. A. and Abazi, I. D. Physico-chemical analysis of fish pond water in freshwater areas of Bayelsa State, Nigeria. *Greener Journal of Biological Science*, 2014; 4(2): 033 – 038.
51. Khattab, F. K. Histological and ultrastructural studies on the testis of rat after treatment with aluminium chloride. *Journal of Basic and Applied Scientific*, 2007; 1: 63 – 72.
52. Krishna, G. and Hayashi, M. In vivo rodent micronucleus assay: protocol, conduct and data interpretation. *Mutation research*, 2000; 455 (1-2), 155–166.
53. Kumar, R., Nagpure, N. S., Kushwaha, B. and Srivastava, S. K. Investigation of the genotoxicity of malathion to freshwater teleost fish *Channa punctatus* (Bloch) using the micronucleus test and comet assay. *Arch Environ. Con. Tox*, 2009; 58 (1) pp. 123-130
54. Lanzarin, G. A. B., Venancio, C. A. S., Monteiro, S. M, and Felix, L. M. Behavioural toxicity of relevant concentrations of glyphosate commercial formulation UltraMax- During Zebra fish embryogenesis. *Chemosphere.* 2020; 253:126636-126636.
55. Lewbert, G. A. Diagnostic techniques for fish. Proceedings of the atlantic coast veterinary conference. 2001; <http://www.din.com/viNDBPub/searchpb/proceedings/PRO5000/CR00335.HTM> 8/19/2008
56. Lowry, O. H., Rosenbrough, N. J., Farr, A. L. and Randall, B. J. Protein measurements with the Folin phenol reagent. *Journal of Biology Chemistry*, 1951; 193 pp. 265-275
57. Luskova V., Svoboda, M. and Koarova, J. The effect of diazinon on blood plasma biochemistry in carp (*Cyprinus carpio L.*). *Acta Veterinaria Brno*, 2002; 71, 117 – 123.
58. Lutnicka, H., Bojarski, B., Ludwikowska, A., Wrońska, D., Kamińska, T., Szczygieł, J., Troszok, A., Szambelan, K. and Formicki, G. Hematological alterations as a response to exposure to selected fungicides in common carp (*Cyprinus carpio L.*). *Folia Biol.* 2016; 64:235–44.

59. Magar, R. S. and Shaikh, A. Biochemical changes in proteins and amino acids in *Channa punctatus* in responses to sub lethal treatment with the insecticide Malathion. *Trends in Life Sciences*, 2012; 1(3): 19 – 23.
60. Majumder, R. and Kaviraj, A. Acute and sublethal effects of organophosphate insecticide chlorpyrifos on freshwater fish *Oreochromis niloticus*. *Drug Chem Toxicol.* 2019; 42:487–95
61. Matozzo, V., Fabrello, J. and Marin, M. G. The Effects of glyphosate and its commercial formulations to marine invertebrates: A Review. *J Mar Sci Eng.* 2020; 8:399.
62. Maurya, P. K., Malik, D. S., Yadav, K. K., Gupta, N. and Kumar, S. Haematological and histological changes in fish *Heteropneustes fossilis* exposed to pesticides from industrial waste water. *Hum Ecol Risk Assess.* ISSN: 2019; 1080-7039
63. Mdegela, R. H., Braathen, M., Mosha, R. D., SKaare, J. U. and Sandvik, M. Assessment of pollution in sewage ponds using biomarker responses in wild African sharptooth catfish (*Clarias gariepinus*) in Tanzania. *Ecotoxicology.* 2010; 19, 722–734.
64. Melefa, T. D. Morphological, hematological and biochemical changes in African catfish *Clarias gariepinus* (Burchell 1822) juveniles exposed to clotrimazole. *Comparative Biochemistry and Physiology, Part C*, 2020; 236, 108815.
65. Menezes, C. C., Fonseca, M. B., Loro, V. L., Santi, A., Cattaneo, R., Clasen, B., Pretto, A. and Morsch, V. M. Roundup effects on oxidative stress parameters and recovery pattern of *Rhamdia quelen*. *Arch. Environ. Contam. Toxicol.* 2011; 60: 665–671.
66. Mensah, P. K., Palmer, C. G. and Muller, W. J. *Pesticides: Toxic Aspects.* Lethal and Sub lethal Effects of Pesticides on Aquatic Organisms: the Case of a Freshwater Shrimp Exposure to Roundup® InTech Publications Rijeka, Croatia. 2014; pp. 163–185.
67. Miri, M., and Khandan, B. H. Effects of aluminum on some tissue enzymes of gills, liver and muscles in common carp (*Cyprinus carpio*). *Iranian Journal of Fisheries Sciences*, 2017; 16 (2), 742–752.
68. Modesto, K. A. and Martinez, C. B. R. Effects of Roundup Transorb on fish: Hematology, antioxidant defenses and acetylcholinesterase activity. *Chemosphere* 2010; 81: 781–787.
69. Mohamed, A. M., El-Emam, M. A., Osman, G. Y., Abdel-Hamid, H. and Ali, R. E. Effect of basudin, selecron and the phytoalkaloid colchicine (pesticides) on biological and molecular parameters of *Biomphalaria alexandrina* snails. *Pesticide Biochem. Physiol.*, 2012; 102:68–78.
70. Monferran M. V., Pesce S. F., Cazenave J. and Wundelin D. A. Detoxification and antioxidant responses in diverse organs of fish experimentally exposed to 1, 2 and 1, 4 atichlorobenzene. *Environmental Research*, 2005; 23: 184 – 92.
71. Mostakim, M. G., Zahangir, M. M., Monir, Mishu M., Rahman, M. K and Islam, M. S. Alterations of blood parameters and histoarchitecture of liver and kidney of silver barb after chronic exposure to quinalphos. *J. Toxicol.* 2015; 41:59-84.
72. Moustafa, G., Shaaban, F., Hadeed, A. and Elhady, W. Immunotoxicological, biochemical, and histopathological studies on Roundup and Stomp herbicides in Nile catfish (*Clarias gariepinus*). *Vet. World.* 2016; 9: 638-647.
73. National Agricultural Statistics Services. Catfish Processing. United States Department of Agriculture. 2010; Available (Accessed May 13, 2010). <http://usda.mannlib.cornell.edu/usda/current/catfroc/catproc-04-19-2010.pdf>.

74. National Root Crops Research Institute. Annual Report. North Carolina Cooperative Extension Service (NCCS). Soil facts; poultry manure as a fertilizer source. Publication AG-439-5. 2010.
75. Nayak, S., Raut, D. and Patnaik, L. Naphthalene Induced Enzymatic Alterations In The Liver Of Climbing Perch, ANABAS TESTUDINEUS. *Journal of Aquatic Biology and Fisheries*, 2019; 7, 134–141.
76. Naveed, A., Janaiah, C. and Venkateshwarlu, P. The effects of lihocin toxicity on protein metabolism of the fresh water edible fish, *Channa punctatus* (Bloch). *Journal of Toxicology and Environmental Health Sciences*, 2010; 3(1): 018 – 023.
77. Noga, E. J. *Fish Diseases: Diagnosis and Treatment*. Iowa, USA, Wiley Blackwell. 2012; 367pp.
78. Nwani, C. D., Ifo, C., Nwamba, H. O., Ejere, V. C., Onyishi, G. C., Oluah, S. N., Ikwuagwu, O. E. and Odo, G. E. Oxidative stress and biochemical responses in the tissues of African catfish *Clarias gariepinus* juvenile following exposure to primextra herbicide. *Drug and chemical Toxicology*. 2014. DOI: 10.3109/01480545.2014.947503.
79. Nwani, C. D, Ibiam, U. A, Ibiam, O. U., Nworie, O., Onyishi, G. and Atama, C. Investigation on acute toxicity and behavioural changes in *Tilapia zilli* due to glyphosate-based herbicide, forceup. *J Anim and Plant Sci*. 2013; 23(3):888-892.
80. Odo, G. E., Agwu, E., Ivoke, N., Ejere, V. C., Atama, C. I., Ezea, C. O. and Aguoru, G. C. Effect of short-term exposure to cyperdicot on behavioural and haematological responses in African catfish *Clarias gariepinus* *Turkish J Fish Aquat Sci*. 2015; 17:61-70
81. Ogueji, E. O., Iheanacho, S. C., Nwani, C. D., Mbah, C. E, Okeke, O. G. and Usman, I. B. Toxicity of diazepam on lipid peroxidation, biochemical and oxidative stress indicators on liver and gill tissues of African catfish *Clarias gariepinus* (Burchell, 1822). *International Journal of Fisheries and Aquatic Studies*, 2017; 5(3): 114 – 123.
82. Okayi, R. G., Annune, P. A., Tachia, M. U. and Oshoke, O. J. Acute toxicity of glyphosate on *Clarias gariepinus* fingerlings, *Journal Resources of Wildlife and Environment*. 2010; 2 (2).
83. Okpala-Ezennia, K. P., Nlewadim, A. A. and Ikeogu, C. I. Impact of sublethal Atrazine exposure on haematology, serum biochemistry, and muscular accumulation on juveniles of *Clarias gariepinus* and its hybrid with *Heterobranchus longifilis*. *Journal of Marine Biology Environmental Science* 2024; 6 (1):1 22. DOI: <https://doi.org/10.36266/JMBES/131>
84. Oladokun, E. L., Sogbanmu, T. O. and Anikwe, J. C. Sublethal concentration of dichlorvos and paraquat induce genotoxic and histological effects in *Clarias gariepinus*. *Environmental Analysis, Health and Toxicology*. 2020; 35(3).
85. Olanike, K. A. Haematological Profile of *Clarias gariepinus* (Burchell, 1822) Exposed to Lead. *Turkish Journal of Fisheries and Aquatic Sciences* 2007; 7: 163-169
86. Ololade, I. A. and Oginni, O. Toxic stress and hematological effects of nickel on African catfish, *Clarias gariepinus*, fingerlings. *Journal of Environmental Chemistry and Ecotoxicology*, 2010; 22: 14-19
87. Olufayo, M. O. Haematological Characteristics of *C. gariepinus* juveniles exposed to Derris root powder, *African Journal of Food Agricultural and Nutrient Development*, 2009; 9(3):921-933 <https://doi.org/10.4314/ajfand.v9i3.43115>

88. Omoniyi, M. P. Toxicity effect of Atrazine on histology, haematology and biochemical indices of *Clarias gariepinus*. *International Journal of Fisheries and Aquatic Studies*; 2018; 6(3): 87-92.
89. Onah, A. I. Acute oral toxicity studies of the crude extract of endophytic fungi isolated from *Annona senegalensis Pers.* *Journal of Drug Delivery and Therapeutics*, 2020; 10 (3), 207–216.
90. Opute, P. A. and Oboh, I. P. Effects of Sub-Lethal Atrazine Concentrations on Embryogenesis, Larval Survival and Growth of African catfish, *Clarias gariepinus* (Burchell, 1822). *Aquaculture Studies*, 2021; 21, pp. 143-152
91. Organization of Economic Cooperation Development. Guidelines for testing chemicals, Guidelines No. 249: Fish Cell Line Acute Toxicity: The RTgill-W1 cell line assay. Adopted July, 14. 2021.
92. Ortiz, J. B., De Canales, M. I. G. and Sarasqutte C. Histopathological changes induced by lindane in various organs of fish. *Scientia Marina*, 2003; 67(1): 53 – 81.
93. Osioma, E. I. and Ejoh, A. S. Activities of acetylcholinesterase, oxidative and nitrosative stress markers in *Clarias gariepinus* exposed to ‘uproot’, a glyphosate–based herbicide. *Int. J. Zool Investig.* 2021; 7:2454-3055
94. Pesce, S. F., Cazenave, J., Monferra, M. N., Frede, S. and Wunderlin, D. A. Integrated survey on toxic effects of lindane on neotropical fish; *Corydoras paleatus* and *Jenynsia multidentata*. *Environmental Pollution*, 2008; 156: 775 – 783.
95. Peillex, C. and Pelletier M. The impact and toxicity of glyphosate and glyphosate based herbicides on health and immunity. *J. Immunotoxicol.* 2020; 17(1):163-174.
96. Portier, C. J. Differences in the carcinogenic evaluation of glyphosate between the International Agency for Research on Cancer (IARC) and the European Food Safety Authority (EFSA), *Journal of Epidemiology. Community Health*, 2016; <http://dx.doi.org/10.1136/jech-2015-207005>
97. Prabesh, S. K., Amit, K. S., Gudrun, De Boeck and Kumar, S. Modulation of blood biochemical parameters of golden mahseer, *Tor putitora* following exposures to single and mixed organophosphate. *Comparative Biochemistry and Physiology Part C Toxicology and Pharmacology.* 2021; 251:109207. <http://doi.org/10.1016/j.cbpc.2021.109207>.
98. Pruszyński, T. and Pistelok, F. Biological and economical evaluation of African and European catfish rearing in water recirculatory systems. *Archives of Polish Fisheries*, 1999; 7(2): 343 – 352
99. Raje, V. Leptin-the central regulator of starvation. *Journal of Postdoctoral Research*, 2018; 1 (2), 21–34.
100. Rajput, V., Singh, S. K. and Arpita, K. Comparative toxicity of Butachlor, Imidacloprid and Sodium fluoride on protein profile of the walking cat fish *Clarias batrachus*. *Journal of Applied Pharmaceutical Science*, 2012; 2(06): 121 – 124
101. Rao, D. S. Carbaryl induced changes in the haematological, serum biochemical and immunological responses of common carp, *Cyprinus carpio*, (L.) with special emphasis on herbal extracts as immune modulators, Ph.D Thesis Andhra University, India, 2010.
102. Sakr, S. A. and Al-amoudi, W. M. Protective effect of silymarin on metiram fungicide-induced hepatotoxicity in albino rats. *Res. J. Pharm. Biol. Chem Sci.* 2012; 3:691–9. 75.

103. Sakr, S., El-Kenawy, A. and El-Sahra, D. Metiram-induced nephrotoxicity in albino mice: effect of licorice aqueous extract. *Environ Toxicol.* 2013; 28(7):372–9
104. Salam, M. A., Shahjahan, M., Sharmin, S., Haque, F. and Rahman, M. K. Effects of sub-lethal doses of an organo-phosphorous insecticide sumithion on some hematological parameters in common carp, *Cyprinus carpio*, *Pakistan J. Zool.* 2015; (47)1487–1491.
105. Santos, E., Ceron, J. J. and Ferrando, M. D. Cholinesterase activity and hematological parameters as biomarkers of sublethal molinate exposure in *Anguilla anguilla*. *Ecotoxicological and Environmental Safety*, 2010; 46: 81 – 86.
106. Saravanan, M., Kumar, P. K. and Ramesh, M. Haematological and biochemical responses of freshwater teleost, *Cyprinus carpio* (*Actinoperigii: Cypriniformes*) during acute and sublethal exposure to lindane. *Pestic. Biochem. Physiol.* 2011; 100, 206– 211.
107. Schmidt, F. H. 1961. Enzymatic determination of glucose and fructose simultaneously *Klin. Wochenschr.*, 1961; 39 pp. 1244-1250
108. Schwarzenbach, R. P., Escher, B. I., Fenner, K., Hofstetter, T. B., Johnson, C. A., von Gunten, U. and Wehrli, B. The challenge of micropollutants in aquatic system. *Science*, 2006; 313: 1072 – 1077.
109. Shah, Z. U. and Parveen, S. A. Review on pesticides pollution in aquatic ecosystem and probable adverse effects on fish. *Pollut. Res.* 2020; 39 (2), 309–321.
110. Shaalan, M., Mohamed, A. E., Mahmoud, I. M. D., Elsayed, M. Y., Abdelwahab, A. A., Asmaa, I. A., Simon, J. D., Rowida, E. I. and Afaf, N. A. R. Toxicological insight of metiram: immunoxidative, neuro-behavioral, and hematobiochemical changes during acute exposure of Nile tilapia (*Oreochromis niloticus*). *BMC Veterinary Research* 2024; 20:303 <https://doi.org/10.1186/s12917-024-04126-4>
111. Shankar, K. M., Kiran, B. R. and Venkateshwarlu, M. “A review on toxicity of pesticides in fish”, *International Journal of Open Scientific Research*, 2013; 1(1), 15-36.
112. Sharma, M. Behavioural responses in effect to chemical stress in fish: a review. *Int J Fish Aquat Stud.* 2019; 7:01–5.
113. Shivaraj, Y. and Asiya, N. F. B Toxic effect of carbohydrate metabolism changes on carp (*Cyprinus carpio*) exposed to cadmium chloride. *International Journal of Fisheries and Aquatic Studies*, 2018; 6(2): 364 – 371.
114. Simakani, P., Abolhasani, M. H. and Hoseini, S. M. Determination of mancozeb toxicity and biochemical effects in common carp (*Cyprinus carpio*). *Int. J. Aquat. Biol.* 2018; 6:157–61
115. Sogbamu, T. O., Osibona, A. O., Oguntunde, A. O. and Otitolaju, A. A. Biomarkers of toxicity in *Clarias gariepinus* exposed to sublethal concentrations of polycyclic aromatic hydrocarbons. *African Journal of Aquatic Science*, 2018; 43:281-92
116. Somdare, P. O., Nwani, C. D., Nwadinigwe, A. O., Nwani, J. C., Odo, G. E., Ugbor, O. N., Ukonze, J. A. and Ezeibe, A. B. C. Fenthion induced toxicity and histopathological changes in gill tissue of *Clarias gariepinus* (Burchell, 1822). *African Journal of Biotechnology*, 2015; 14: 2103 – 2113.
117. Spector, T. Refinement of the Coomassie blue method of protein quantitation. A simple and linear spectrophotometric assay for less than or equal to 0.5 to 50 micrograms of protein. *Analytical biochemistry*, 1978; 86 (1), 142–146.
118. Steel, R. G. and Torrie, J. H. Principles and Procedures of Statistics. MacMillan Publishing Company. New York, 748. 2003.

119. Tchounwou, P. B., Patlolla, A. K., Yedjou, C. G. and Moore, P. D. Environmental Exposure and Health Effects Associated with Malathion toxicity, Toxicity and Hazard of Agrochemicals Marcelo L. Larramendy and Sonia Soloneski, Intech Open, 2015.
120. Thanomsit, C., Wattanakornsiri, A. and Nanthanawat, P. Effect of glyphosate on fish behaviour and histological alteration of gills in Asian sea bass (*Lates calcarifer*). *Burapha Sci J.* 2016; 21:204-215.
121. Udume, B. U., Ota, H. O., Zhifeng, Yan and Onyewuchi, C. C. Evaluating the effect of acute ammonia toxicity in a freshwater ecological environment on the growth and haematological functioning of *Clarias gariepinus*. *Journal of Water Supply: Research and Technology*, 2002a; Vol.71 No 3, 464. <https://doi.org/10.2166/aqua.2022.138>
122. Udume, B. U., Anyaele, U. and Ota, H. O. The impact of sub-lethal concentrations of glyphosate on growth and haematology of African catfish under aquatic ecological micro-climate. *Journal of Environmental Chemistry and Ecotoxicology*, 2022b; 4:164-170. <http://dx.doi.org/10.1016/J.enceco.2022.06.001>
123. Ugbomeh, A. P., Bob-manuel, K. N. O., Green, A. and Taylorharry, O. Biochemical toxicity of Corexit 9500 dispersant on the gills, liver and kidney of juvenile *Clarias gariepinus*. *Fisheries and Aquatic Sciences*, 2019; 22(15): 1 – 8
124. United States Environmental Protection Agency. Sewage sludge (biosolids): frequently asked questions. 2012; [Accessed January 2014]; <http://water.epa.gov/polwaste/wastewater/treatment/biosolids/genqa.cfm>.
125. Uren Webster, T. M., Laing, L.V., Florance, H. and Santos, E. M. Effects of glyphosate and its formulation, Roundup, on reproduction in zebra fish (*Danio rerio*). *Environmental Science & Technology*, 2013; 48(2):1271-1279. <http://pubs.acs.org/doi/abs/10.1021/es404258h>.
126. Varadharajan, R. Biochemical effects of different phenolic compounds on *Oreochromis mossambicus* Peters. Thesis submitted to the Cochin University of Science and Technology, India. 2010.
127. Wang, Y., Gao, Z., Liu, C., Mao, L., Liu, X., Ren, J., Lu, Z., Yao, J. and Liu, X. Mixture toxicity of pyraclostrobine and metiram to the zebra fish (*Danio rerio*) and its potential mechanism. *Environ Sci Pollut Res Int.* 2023; 30(15):44400–14.
128. Williams, A. K. Organochlorine Pesticides Residues in Shellfishes and Finfishes from Lagos Lagoon. *Ph.D. Thesis*, Covenant University, Ota, Nigeria. 2011; 3(3):38-45. (ISSN: 1553-9865). <http://www.sciencepub.net>