

## Review Article

# Development Of Bioprocess Technology For Parasite Control In Aquaculture Commodities

Commented [MkC1]: Small letters

### ABSTRACT

Controlling parasites in cultivated commodities is a crucial **issue** in the fisheries industry that can impact fish health and production quality. The use of conventional methods such as chemical drugs often results in negative side effects on the environment and human health. Therefore, the development of bioprocess technology offers a more environmentally friendly and sustainable solution for parasite management. In bioprocess technology, probiotic microorganisms are used to make fish stronger, enzymes made by microorganisms are used to lower the number of parasites, and natural predators are used for biological control. Additionally, research also focuses on the development of vaccines to target specific parasites, providing a more permanent control alternative. These approaches are expected to reduce dependence on synthetic chemicals, improve the sustainability of aquaculture, and minimize the risk of parasite resistance. The development and application of this bioprocess technology have the potential to enhance the efficiency and sustainability of aquaculture production.

Commented [MkC2]: Replace 'issue' with 'challenge'

*Keywords: Aquaculture Commodities, Bioprocess, Parasites, Probiotics, Sustainable Aquaculture*

### 1. INTRODUCTION

Cultivation activities currently remain the main choice for the community because of its great potential in supporting the economy and meeting food needs. The high interest in aquaculture shows that the sector has bright prospects and continues to grow, both on a small and large scale (Little et al. 2016). However, in practice, there are various obstacles that are often faced, such as low production efficiency, fluctuations in crop quality, and environmental impacts that are not well managed. To overcome these various obstacles, innovations and technologies that are able to increase productivity and sustainability of aquaculture are needed (Mustafa et al. 2021). One approach that can be applied is bioprocessing, which is the use of biological-based technology to optimize cultivation yields. With the application of bioprocessing, it is hoped that cultivation can become more efficient, produce high-quality products, and remain environmentally friendly.

Commented [MkC3]: Replace "Cultivation activities currently remain the main choice for the community..." with "Aquaculture remains a primary method for supporting economic development and meeting global food demands."

Commented [MkC4]:

The development of bioprocess technology for parasite control in farmed fish is very important to increase productivity and sustainability of aquaculture (Kassim et al. 2014). The application

of this bioprocess technology can produce healthier fish populations, reduce dependence on chemical processing and encourage more environmentally friendly aquaculture practices. By integrating natural biocontrol agents and optimizing growth conditions, aquaculture can achieve a balance that improves fish health while minimizing ecological impacts (Assefa & Abunna, 2018). Innovative approaches such as the use of microbes can further enhance efforts to create ecosystems that support sustainable farming practices. Additionally, it can increase resistance to disease and improve overall fish health.

Microbes used in bioprocessing can come from a variety of sources, including fish bodies. As is known, the body of fish, especially the intestines, is a habitat for various types of bacteria. Aisyah et al. (2019) reported that the bacteria with the highest abundance in the intestines of fish include *Cetobacterium*, *Clostridium sensu stricto 1*, *Bacteroides*, *Enterovibrio*, *Plesiomonas*, *Lactococcus*, *Romboutsia*, *Stenotrophomonas*, *Turicibacter*, and *Edwardsiella*, among other types. These various bacteria have important roles in the metabolic process of fish, including digestion, vitamin synthesis, and protection against pathogens. In addition, these bacteria also have the potential to be utilized in bioprocesses, such as enzyme production, probiotics, and waste treatment, making them an important component in the development of sustainable fisheries biotechnology.

The many potentials possessed by these different types of bacteria show great opportunities in the development of bioprocess techniques. With the right application, this bacterium can be used to support various innovations in the fisheries sector. In order to maximize these benefits, it is important to pay attention to the characteristics of fish parasites, as the interaction between microbes, fish, and parasites can affect the effectiveness of bioprocesses as well as the overall health of the fish. These parasites can be categorized into different types, each affecting fish differently based on their habitat and environmental conditions. [Understanding the nature of these parasites is essential for effective management and control strategies.](#)

## 2. PARASITIC PROBLEMS IN CULTIVATED FISH

[These parasites include protozoan and metazoan groups, such as Microsporidia and Cestoda \(tapeworms\)](#), which are often found in both freshwater and marine fish. These parasitic infections not only impact fish growth and reproduction, but can also cause mass deaths, especially in intensive aquaculture systems (Kanwal et al., 2023; Scholz et al., 2021).

The type of parasitic infection is often specific depending on the habitat of the fish. In freshwater fish, infections such as Renal Sphaerosporosis caused by protozoa of the genus *Sphaerospora* can impair kidney function, disrupt metabolism, and cause high mortality rates. In contrast, marine fish are more susceptible to diseases such as Cryptocaryonosis (sea white spot disease) caused by *Cryptocaryon irritans* and Brooklynelliosis which attacks the skin and gills of marine fish, often causing severe tissue damage (Kanwal et al., 2023).

Some fish parasites not only cause health problems in their hosts but also act as vectors for other pathogens. An example is the copepod *Lernanthropus kroyeri*, which is known to carry and transmit pathogenic bacteria such as *Vibrio anguillarum*. This facilitates the spread of disease in fish populations, especially in high-density environments such as aquaculture (Yildiz & Otcucuoglu, 2021).

## 3. FISH PARASITE CONTROL AND MANAGEMENT

Fish parasite control and management is an important aspect in ensuring the health of farmed fish and the quality of the products produced. Parasitic infections can reduce fish growth, increase mortality rates, and reduce the quality of fish meat which can have an impact on the

**Commented [MkC5]:** Include a brief mention of specific challenges in parasite management in aquaculture to set up the discussion.

**Commented [MkC6]:** Rephrase "These parasites include protozoan and metazoan groups, such as Microsporidia and Cestoda..." to "Parasitic infections in aquaculture are caused by protozoans (e.g., Microsporidia) and metazoans (e.g., Cestoda)."

**Commented [MkC7]:** Can you include a table or chart summarizing common parasites, their effects, and affected species for clarity.

profits of fish farmers. Therefore, an effective parasite control approach is indispensable to prevent greater losses.

1. **One way to control parasites is through good environmental management.** Optimal environmental conditions, such as the right temperature, salinity, and pH, can help reduce the conditions that favor the parasite's life. In addition, maintaining pond cleanliness and using a good water circulation system can reduce the accumulation of parasites in the cultivation environment. These measures can help reduce stress in fish, which is often a trigger factor for the high prevalence of parasites. Effective control strategies include improving fish farming practices and monitoring environmental conditions to reduce the prevalence of parasites (Kanwal et al., 2023).

**Commented [MkC8]:** Replace "One way to control parasites is through good environmental management" with "Effective environmental management is a fundamental approach to parasite control."

2. In addition to environmental management, the use of biotechnology methods can help in the control of parasites in fish farming. One approach is the use of probiotics, such as good bacteria and microalgae, which can increase the fish's immune system to parasitic infections. Probiotics work by improving the balance of the fish's gut microflora, thus inhibiting the growth of parasites naturally.

3. Bioprocess technologies, such as the use of enzymes produced by microorganisms, also have great potential in reducing the number of parasites in aquaculture systems. For example, praziquantel has been shown to be effective against adult tapeworms, although the management of the parasitic larval stage is still a challenge (Scholz et al., 2021).

4. In addition, biological control by utilizing predators or natural competitors to manage parasite populations is an alternative that can reduce dependence on chemicals (Buchmann, 2022). Another promising approach is immunological methods, such as the development of vaccines that target specific parasites, including protozoa and metazoa (Buchmann, 2015).

5. The use of antimicrobial drugs, such as praziquantel, is one option in parasite control in fish. However, its use must be done with caution and in accordance with the right dosage to prevent parasite resistance to the drug. Therefore, a combination of environmental management, the application of biotechnology, and measurable treatment becomes an effective strategy in parasite management in farmed fish. Although dependence on drugs such as praziquantel is still common, the emergence of drug resistance is an increasingly worrying issue and requires more attention (Widdicombe et al., 2024).

#### 4. CONCEPT OF BIOPROCESS TECHNOLOGY IN PARASITE CONTROL

Bioprocess technology plays an important role in developing sustainable methods for controlling parasites, utilizing biological systems and processes to produce effective biopesticides and other control agents. This innovative approach not only minimizes dependence on chemical pesticides but also improves ecological balance, encouraging healthier agricultural practices. and reducing environmental impacts (Vurro & Gressel 2007).

By utilizing natural microorganisms and compounds, bioprocess technology offers a promising alternative that can target specific parasites while preserving beneficial organisms in the ecosystem. This targeted strategy not only increases crop yields but also contributes to overall crop resilience. agricultural system, encouraging a more sustainable and environmentally friendly pest management approach (Kamal et al. 2015). There are several applications of bioprocess technology in disease control, which can be seen in the following table.

[Table 1. Application of Bioprocess in Parasite Control](#)

**Commented [MkC9]:** The table on bioprocess applications is informative but lacks consistent formatting. Ensure all columns are labeled clearly, and references are standardized.

No.	Bioprocess Technology	Description	Advantage	Reference
1	Probiotics	Induction of Bacillus CgM22 Probiotic in Feed	Increases the resistance of fish	(Haetami <i>et al.</i> 2022)
2	Vaccine Manufacturing	Administration of Anti-helminthic vaccines in fish	Effective treatment strategies to prevent getting worms/helminths	Widdicomb <i>et al.</i> , 2024
3	Probiotics	Induction of <i>Staphylococcus</i> sp. JC20 from Octopus Digestive Tract ( <i>Octopus</i> sp.)	Increases the immune system of fish	Istiqomah <i>et al.</i> 2019
4	Probiotics	Induction of bacteria from the intestines of fish to fight <i>Aeromonas</i> .	Increases the resistance of fish from parasite attacks	Mulyani <i>et al.</i> 2018
5	Vaccination creation	Vaccination of Tiger Grouper Fish ( <i>Epinephelus fuscoguttatus</i> ) With Three pathogenic bacteria that have been inactivated	Increases the resistance of fish to parasitic attacks	Zafran, 2015
6	Addition of organic matter	Addition of miana leaf extract ( <i>Coleus scutellarioides</i> ) to feed given to tiger shrimp ( <i>Penaeus monodon</i> )	Increases tiger shrimp's immunity	Febriani <i>et al.</i> 2023
7	Enzyme Inhibitors Cause Parasite Development	Administration of protease inhibitors to inhibit the development of parasites,	This can increase the effectiveness of treatment against various parasitic infections,	Rascon and McKerrow, 2013

No.	Bioprocess Technology	Description	Advantage	Reference
		virulence and pathogenesis	potentially leading to a more efficient and targeted therapeutic approach.	
8	Probiotics	<i>Acetobacter</i> sp., <i>Lactobacillus</i> sp., and <i>Saccharomyces cerevisiae</i> (yeast)	Using natural ingredients that are easy to obtain, safe, and have the potential as an alternative to chemicals to increase fish's immunity	Safir <i>et al.</i> 2023
9	Biofiltration	Administration of 3 probiotics in the fish filtration system. Novozymes pond plus (Novozymes Biological, Boston), Zhongshui BIO-AQUA (Zhongshui Fish Medicine, Wuxi, China) and Effective Microorganisms (Jiangsu Suwei Microbiology Research, Wuxi, China) at Fengqiao farm (Zhuji, China)	A water filtration system that incorporates decomposing microorganisms to reduce contaminants, including parasites, in the aquaculture water.	2016 Donation

The development of bioprocess technology for parasite control in fish farming is increasingly important to support the sustainability of the fisheries sector. One promising approach is the use of microorganisms or natural compounds that can inhibit or kill parasites without damaging the balance of the ecosystem. For example, the use of probiotics or decomposing bacteria that can reduce the number of parasites in fish by affecting the microflora in the fish's body or in their surrounding environment. Application of probiotic bacteria in fish (Aisyah *et al.* 2022)

In addition, enzyme-based technologies are also being developed to break down the body walls of parasites or disrupt their life cycles (Rascon and McKerrow, 2013). Enzymes extracted

from certain microorganisms or plants can be used as an environmentally friendly alternative to parasite control. The application of this technology can also reduce dependence on synthetic chemicals that have the potential to cause adverse effects on the health of fish and the surrounding environment.

On the other hand, the application of biotechnology to improve water quality in aquaculture systems also has the potential to reduce the prevalence of parasites. One example is the development of a biofiltration system that incorporates decomposing microorganisms to clean water from feed residues and feces that can be a source of parasite development (ranjan and bavitha 2014). With the combination of these various bioprocess technologies, parasite control in fish farming can be carried out more efficiently and environmentally friendly.

## REFERENCES

1. Aisyah, Andriani, Y., & Mulyani, Y. (2022). Application of Probiotic Bacteria in Fish Feed. *Journal of Ruaya*, 10(1), 1-7.
2. Aisyah., Mulyani, Y., Dewanti, L.P., and Agung, M. U. K., (2019). Community Structure Of Fish Gut Bacteria Which Have Differences Of Feeding Habits. *GSJ*, 7(12).
3. Assefa, A., & Abunna, F. (2018). Maintenance of Fish Health in Aquaculture: Review of Epidemiological Approaches for Prevention and Control of Infectious Disease of Fish. *Veterinary Medicine International*. <https://doi.org/10.1155/2018/5432497>
4. Buchmann, K. (2015). Impact and control of protozoan parasites in maricultured fishes. *Parasitology*, 142(1), 168-177.
5. Buchmann, K. (2022). Neutrophils and aquatic pathogens. *Parasite Immunology*, 44(6), e12915.
6. Febriani, K., Basir, B., & Heriansah, H. (2023). Survival rate of tiger shrimp (*Penaeus monodon*) pre and post infected with White Spot Syndrome Virus (WSSV) fed with the addition of miana leaf extract (*Coleus scutellarioides*). *Aquaatikisla: Journal of Aquaculture, Coastal and Small Islands*, 7(2), 151-158.
7. Haetami, K., Aisyah, Andriani, Y., & Mulyani, Y. (2022). Application of Probiotic Bacteria in Fish Feed. *Journal of Ruaya*, 10(1), 1-7.
8. Istiqomah, I., Isnansetyo, A., Atitus, I. N., & Rohman, A. F. (2019). Isolation of cellulolytic bacterium *Staphylococcus* sp. JC20 from the Intestine of octopus (*Octopus* sp.) for fish probiotic candidate. *Journal of Fisheries, Gadjah Mada University*, 21(2), 93-98.
9. Jerônimo, G. T., Cruz, M. G. da, Bertaglia, E. de A., Furtado, W. E., & Martins, M. L. (2022). Fish parasites can reflect environmental quality in fish farms. *Reviews in Aquaculture*. <https://doi.org/10.1111/raq.12662>
10. Kamal, R., Gusain, Y. S., Kumar, V., & Sharma, A. K. (2015). Disease management through biological control agents: An eco-friendly and cost effective approach for sustainable agriculture-A Review. *Agricultural Reviews*, 36(1), 37-45.

11. Kassim, Z., John, A., Chin, L. K., Zakaria, N. F., & Asgnari, N. H. (2014). Sustainable Technique for Selected Live Feed Culture. <https://doi.org/10.5772/57212>
12. Little, D. C., Newton, R., & Beveridge, M. (2016). Aquaculture: a rapidly growing and significant source of sustainable food? Status, transitions and potential. <https://doi.org/10.1017/S0029665116000665>
13. Mulyani, Y., Aryantha, I. N. P., Suhandono, S., & Pancoro, A. (2018). Intestinal Bacteria of Common Carp (*Cyprinus carpio* L.) as a Biological Control Agent for *Aeromonas*. *Journal of Pure & Applied Microbiology*, 12(2).
14. Mustafa, S., Estim, A., Shapawi, R., Shalehand, M. J., & Sidik, S. R. M. (2021). Technological applications and adaptations in aquaculture for progress towards sustainable development and seafood security. <https://doi.org/10.1088/1755-1315/718/1/012041>
15. Ranjan, R., & Bavitha, M. (2014). Bioremediation - A potential tool for management of aquatic pollution. *International Journal of Multidisciplinary Research and Development*.
16. Rascón, A., & McKerrow, J. (2013). Synthetic and natural protease inhibitors provide insights into parasite development, virulence and pathogenesis. *Current Medicinal Chemistry*. <https://doi.org/10.2174/0929867311320250005>
17. Safir, M., Armansyah, M., Hasanah, N., & Mangitung, S. F. (2023). Growth, and Feed Conversion Ratio of *Pangasius hypophthalmus* Fish; Sauvage, 1878) Fermented Feed with Different Doses of Probiotics. *Journal of Fishery Science and Innovation*, 7(1), 28-34.
18. Tang, J., Dai, Y., Li, Y., Qin, J., & Wang, Y. (2016). Can Application of Commercial Microbial Products Improve Fish Growth and Water Quality in Freshwater Polyculture. *North American Journal of Aquaculture*. <https://doi.org/10.1080/15222055.2015.1116474>
19. Vurro, M., & Gressel, J. (2007). Novel biotechnologies for biocontrol agent enhancement and management.
20. Widdicombe, M., Coff, L., Nowak, B. F., Ramsland, P. A., & Bott, N. J. (2024). Understanding the host response of farmed fish to blood flukes (Trematoda: Aporocotylidae) for developing new treatment strategies. *Fish & Shellfish Immunology*, 109613.
21. Yildiz, H. Y., & Otcuoglu, Ö. (2021). Marine fish parasite, *Lernanthropus kroyeri* (Copepoda) is a repository of *Vibrio anguillarum* as evidenced by Loop-Mediated Isothermal Amplification method. *Ankara Üniversitesi Veteriner Fakültesi Dergisi*.
22. Zafran, Z. (2015). Vaccination of Juvenile Tiger Grouper with Three Inactivated Bacterial Pathogens. *Journal of Tropical Marine Science and Technology*, 7(2), 98505.

**Commented [MkC10]:** Cross-check references for accuracy, particularly the citation years and authors (e.g., "Widdicombe et al., 2024" seems to reference an unpublished work because you didn't insert the journal/publisher).