

DUS DESCRIPTORS BASED CHARACTERIZATION OF MAIZE (*Zea mays* L.) INBRED LINES

ABSTRACT:

DUS testing is one of the important criteria to test inbred lines for distinctness, uniformity and stability. Effective plant breeding programme requires a basic understanding of the existence of genetic variability in a population. Characterization and evaluation of variability is prerequisite for the efficient use of genetic material. Maize is one of the economic crops of global importance. Hence, in the present investigation, morphological characterization of thirty-two maize inbred lines was carried out as per the DUS descriptors. Out of the twenty-seven traits under study, fourteen were found dimorphic and six characters were with trimorphic expression, indicating that all the traits were informative with respect to trait expression cum characterization. Presence of wide variation among the thirty-two maize inbreds would aid in selection of inbreds with desirable characters for further breeding programme. [The present investigation was undertaken to characterize the thirty-two maize inbreds based on DUS descriptors](#)

Keywords: Maize, DUS, -Characterization, Morphological, Inbred lines

Introduction:

Maize (*Zea mays* L.) is the world's third most widely grown cereal crop and is known as "queen of cereals". It is commercially valued economic crop, majorly used in poultry and cereal food industries, provides raw materials for starch, gluten, corn oil, corn syrup, sugar, corn meal and com flour and occupies an important place in Indian agriculture. This emphasizes the demand of maize in India and there is an urging need to develop high yielding single cross hybrids. Therefore, knowledge on genetic diversity of inbred lines would help the breeder in planning crosses for superior hybrid development.

Morphological data plays an important role in management of genetic resources that are conserved in *ex-situ* gene-banks. Many tools are available to study the relationship among the cultivars, including various types of molecular markers; however, morphological characterization is the first step in the description and classification of germplasm.

Characterization of morphological variability facilitates identification of accessions with desirable characteristics such as earliness, disease resistance and improved ear trait etc.

Formatted: Font: Italic

Hence, grouping of lines aids in avoidance of duplication in sampling populations for identification of varieties and hybrids.

The morphological characters are very important indicators to determine the off types in the seed production programme and guides in maintainance of the genetic purity. Protection of Plant Varieties and Farmers Right Authority insists on characterization and registration of extant, farmers and new varieties as a part of national and botanical asset. DUS testing is one of the important criteria to test inbred lines for distinctness, uniformity and stability. DUS Testing of cultivars is one of the requirements for granting Plant Breeders Rights (PBR) and it is conducted according to national guidelines prepared on the basis of UPOV guidelines. Maize inbred lines represent a fundamental resource for studies in genetics and breeding and are used extensively in hybrid corn production.

Keeping in view the importance of aforesaid aspects, the present investigation was undertaken to characterize the thirty-two maize inbreds based on DUS descriptors.

Material and methods:

The field experiment was conducted during *kharif*, 2020 at Agricultural Research Station, Tornala, Siddipet district of Telangana state. The seed material was selected during field day at Winter Nursery Centre, ICAR-IIMR, Rajendranagar, Hyderabad during 2019-20 and purified & used for DUS characterization during *kharif*, 2020. A total of ~~thirty-two~~thirty-two maize inbred lines were sown in 4 rows of 6 m length by adopting a spacing of 75 cm between rows and 20 cm between plants with in a row in Randomized Block Design replicated thrice. All the necessary precautions were taken to maintain uniform plant population of each genotype per replication. All the recommended package of practices were adopted besides providing necessary prophylactic plant protection measures to raise a good crop.

Morphological characterization of thirty-two maize inbred lines was carried out using twenty-seven DUS characters. Visual observations were recorded on single plant basis on thirty randomly selected plants in each genotype at appropriate growth stages (Rakshith et al., 2005) on eighteen qualitative characters *viz.*, leaf: angle between blade and stem, altitude of blade and anthocyanin colouration of sheath, stem: anthocyanin colouration of brace roots, tassel: anthocyanin colouration at base of glume, anthocyanin colouration of glumes, anthocyanin colouration of anthers, density of spikelets, angle between main axis and lateral branches, altitude of lateral branches, ear: time of silk emergence, anthocyanin colouration of silks, shape, type of grain, colour of top of grain and anthocyanin colouration of glumes of

Commented [NK1]: In the introduction you posed the problem without referring to any similar work in the literature review. Are you the first person to work on this topic? In my opinion you must refer to at least one author who has worked on a similar topic

Formatted: Font: Italic

cob and kernel: row arrangement and shape. While the data on time of anthesis (50 % plants) and time of silking (50 % plants) was noted on plot basis.

Data on seven quantitative traits such as, leaf: width of blade, ~~tassel~~tassel: length of main axis above lowest side branch (cm), plant: length upto flag leaf (cm), ear: length without husk (cm), ear: diameter without husk (cm), number of rows of grains (no.) and 1000 kernel weight (g) were recorded on five randomly selected plants in each plot.

Results and Discussion:

In the present study, ~~thirty two~~thirty-two maize inbreds were characterized by using twenty seven DUS characters including twenty qualitative and seven quantitative traits as per the norms of national DUS test guide lines. The maize inbreds under study showed wide range of variability with respect to different traits studied. The frequency distribution of all the characters under study is presented (Table 1).

For leaf characters like angle between blade and stem, maximum frequency of 81% showed small angle and rest (19%) was observed for “wide” state of expression indicating that these inbreds are good water harvesters and could be best used under rainfed conditions. This result is in confirmed with the findings of ~~Gull~~Gull *et al.*, whereas attitude of leaf blade showed maximum frequency (66%) for “straight” state of expression, this is very useful in increasing the plant population per acre, ~~there by~~thereby results in higher yield levels. Anthocyanin coloration of brace roots and anthocyanin coloration of silks was observed at a frequency of 69% and 53 % respectively remaining 31 % showed absence of anthocyanin coloration of brace roots and 47 % showed absence of ~~anthocyanin~~of anthocyanin coloration of silks (Fig 2).

Anthocyanin colouration of tassels of different character showed maximum frequency for “absent” state, similar results were observed by ~~Sahaby~~Saha *et al.* This presence of coloration is a useful tool for identification of parental lines in the crossing programme. The density of spikelets on tassels showed maximum frequency (84%) for “dense” state indicated more pollen count on the tassels for maximum inbreds. This is very useful parameter of inbred to ~~used~~use as parental ~~line as~~line as it provides pollen for ~~more~~a greater number of days. Time of ~~anthesis~~of anthesis and silk emergence was found late for all genotypes having frequency of 100 % for each character. This parameter will indirectly ~~indicates~~indicate higher yield levels. The anthesis silking ~~interval~~interval (ASI) was found short for maximum genotypes revealed less pollen loss and effective pollination.

Commented [NK2]: Year?

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

All the germplasm lines were registered absence of anthocyanin coloration of leaf sheath. Angle between main axis & lateral branches and attitude of lateral branches of tassel were observed 100 % ~~narrow~~ narrow and straight state of expression. All the inbred lines were grouped into three categories based on length of main axis of tassel, 13 % were showed short, 84 % were showed medium and 3 % were observed long state of expression. This high medium state of expression will ~~indicates~~ indicate more of pollen production. Regarding plant length of inbred lines maximum 88 % showed short height, it is amicable for used as female parent in the crossing programme. ~~Twenty-nine~~ Twenty-nine inbred lines (91% ~~were~~) were having narrow leaf blade which were less than 8cm and three inbred line (9% ~~were~~) were having medium width of leaf blade which is 8-9cm. This is very useful for cultivation of these lines in the dryland agriculture as transpirational are very less.

The trait ear length without husk recorded with maximum frequency of 69% for “medium”state and the diameter of cobs without husk showed 100% for large state of expression. As length and diameter is directly proportional ~~to production~~ to production and productivity. The shape of ear was registered wide variability for states of expression like conical (6%), conico-cylindrical (85 %) and cylindrical of 9 % state of expression. Number of rows of grains of ear was also observed wide variability for states of expression with maximum of medium state of expression with 56-%. This parameter is also in directly proportional ~~to production~~ to production and productivity of crop.

Type of grain of ear showed two categories with 97% of flint type and 3% of semi flint type. The color of top ~~grain was~~ grain was registered wide variability for states of expression like orange (91%), yellow with cap (3-%) and yellow of 6-% state of expression. This orange grain type is more preferable by the consumers and at industrial level also. Anthocyanin coloration of glumes of cob was noticed maximum of 97-% of white color and 3 % of light purple color.

All inbred lines were classified into two categories with 31 inbred lines were having straight row arrangement of kernels and only 1 inbred line is having irregular row arrangement of kernels. The shape of kernel was showed wide variability for states of expression like maximum of round (56-%), indented (16-%) and toothed (28-%). The results are in close conformity with the findings of Gull et *al.* (2020). Thousand kernel weight also showed variability for all states with highest frequency (75%) recorded in small state.

Conclusion:

Formatted: Font: Italic

Frequency distribution of traits for DUS characterization of corn inbreds indicated that wide variation existed among different genotypes which could be better utilized in the selection of inbreds based on their specific requirement for further breeding programmes. The diversity among the different inbreds could be utilized for cultivar improvement and germplasm conservation programs aimed at improving productivity of corn.

REFERENCES:

- Abu-Alru, I., Christiansen J.L., Madsen S., Sevilla R. and Rtiz R. (2006). Assessing tassel, kernel and ear variation in Peruvian highland maize. *Plant Genome Res.*, 137: 34-38.
- Agarwal, R.L. and Pawar, A. 1990. Identification of soybean varieties on the basis of seed and seedling characters. *Seed Research*. 18: 77-81.
- Baishan, L.U., Wang R.H., Wang F.G., Zhang H.S. and Zhao J.R. (2010). Phenotypic diversity of maize inbred lines based on DUS testing traits. *Plant Genetic Resources*. 11(1): 103-107.
- Burbridge, M.J., Batey, J.L., Campbell, W.P., Skerit, J.H. and Wrigley, C.W. 1986. Distinction between barley varieties by grain characteristics, electrophoresis, chromatography and antibody reaction. *Seed Science and Technology*. 14: 619-629.
- Camussi, A. Spagnoletti Zeuli, P.L. and Melchiorre, P. 1983. Numerical taxonomy of Indian Maize populations: genetic distances on the basis of heterotic effects. *Maydica*, 28: 411-424.
- Dowswell, C., Paliwal R. and Ronald (1996). Maize is the third important crop in world. *Maize Journal*: 10-16.
- Gull, A., Lone A.A., Bhat M.A., Sofi P.A., Khan Z.H., Dar Z.A., Swaranlatha V., Wani M.A., Iqbal A.M., Bhat T.A. and Nazir A. (2020). DUS characterization of sweet corn inbreds under temperate conditions. *Plant Archives*. 20 (1): 2357-2362.
- Gupta, A., Amrapali S., Kumar M., Khati P., LalB., Agrawal P.K., Mahajan V. and Bhatt J.C. (2016). Distinctness, uniformity and stability testing in maize inbreds. *Nat. Acad. Sci. Lett* 39(1), 5-9.
- Rakshith, S., Rao, M.V., Ravi, S.B., Seetharam, A., Makhija, O.P., Sharma, S.P., Dhillon, B.S., Singh, R.V., Tikoo, J.L. 2005. Guidelines for the conduct of test for Distinctiveness, Uniformity and Stability on maize (*Zea mays L.*). [New Delhi](#) 110 012.
- Saha, S., Rana, V.B., Sanghavi, B., Pawan, V., Mahadevu, P. 2024. Morphological characterization and dendrogram analysis of DUS (Distinctiveness, Uniformity, Stability) traits in maize inbred lines (*Zea mays L.*). *Plant Archives*, [24:24](#): 431-438.

Yadav, V.K. and Singh I.S. (2010). Comparative evaluation of Maize inbred lines (*Zea mays*. L) according to DUS testing using morphological, physiological and molecular markers. *Agricult. Sci.*, 1(3), 131-142.

UNDER PEER REVIEW

Fig. 1: DUS characterization of maize inbred lines



Anthocyanin coloration of anthers: Absent



Anthocyanin coloration of anthers: Present



Anthocyanin coloration at base of glume: Absent



Anthocyanin coloration at base of glume: Present



Anthocyanin colouration of brace roots: Absent



Anthocyanin colouration of brace roots: Present









Colouration of glumes of cob: White



Colouration of glumes of cob: Purple

UNDER PEER REVIEW

Fig.2. Graphical representation of Morphological characteristics of maize inbred lines inbred line










Table 1. Frequency distribution of germplasm of maize for various DUS characters

| S. No | Character | State of expression | Number of genotypes | Frequency distribution (%) |
|-------|--|---------------------|---------------------|----------------------------|
| 1 | Leaf: Angle between blade and stem | Small | 26 | 81 |
| | | Wide | 6 | 19 |
| 2 | Leaf: Attitude of blade | Straight | 21 | 66 |
| | | Drooping | 11 | 34 |
| 3 | Stem: Anthocyanin colouration of brace roots | Present | 22 | 69 |
| | | Absent | 10 | 31 |
| 4 | Tassel: Time of anthesis | Late | 32 | 100 |
| 5 | Tassel: Anthocyanin colouration at base of glume | Yes | 13 | 41 |
| | | No | 19 | 59 |
| 6 | Tassel: Anthocyanin colouration of glumes excluding base | Yes | 14 | 44 |
| | | No | 18 | 56 |
| 7 | Tassel: Anthocyanin colouration of anthers | Yes | 6 | 19 |
| | | No | 26 | 81 |
| 8 | Tassel: Density of spikelets | Dense | 27 | 84 |
| | | Sparse | 5 | 16 |
| 9 | Tassel: Angle between main axis and lateral branches | Narrow | 32 | 100 |
| 10 | Tassel: Attitude of lateral branches | Straight | 32 | 100 |
| 11 | Ear: Time of silk emergence | Late | 32 | 100 |
| 12 | Ear: Anthocyanin colouration of silks | Present | 17 | 53 |
| | | Absent | 15 | 47 |
| 13 | Leaf: Anthocyanin colouration of sheath | Absent | 32 | 100 |
| 14 | Tassel: Length of main axis | Short | 4 | 13 |
| | | Medium | 27 | 84 |
| | | Long | 1 | 3 |
| 15 | Plant: Length | Short | 28 | 88 |
| | | Medium | 3 | 9 |
| | | Long | 1 | 3 |
| 17 | Leaf: Width of blade | Narrow | 29 | 91 |
| | | Medium | 3 | 9 |
| 18 | Ear: Length without husk (cm) | Short | 10 | 31 |
| | | Medium | 22 | 69 |
| 19 | Ear: Diameter without husk (cm) | Large | 32 | 100 |
| 20 | Ear: Shape | Conical | 2 | 6 |
| | | Conico-cylindrical | 27 | 85 |

| | | | | |
|----|--|-----------------|----|----|
| | | Cylindrical | 3 | 9 |
| 21 | Ear: Number of rows of grains | Few | 4 | 13 |
| | | Medium | 18 | 56 |
| | | Many | 10 | 31 |
| 22 | Ear: Type of grain | Flint | 31 | 97 |
| | | Semi flint | 1 | 3 |
| 23 | Ear: Color of top grain | Orange | 29 | 91 |
| | | Yellow with cap | 1 | 3 |
| | | Yellow | 2 | 6 |
| 24 | Ear: Anthocyanin coloration of glumes of cob | White | 31 | 97 |
| | | Light purple | 1 | 3 |
| 25 | Kernel: Row arrangement | Straight | 31 | 97 |
| | | Irregular | 1 | 3 |
| 26 | Kernel: Shape | Round | 18 | 56 |
| | | Indented | 5 | 16 |
| | | Toothed | 9 | 28 |
| 27 | Kernel: 1000 kernel weight (g.) | Small | 24 | 75 |
| | | Medium | 8 | 25 |

UNDER PEER REVIEW

UNDER PEER REVIEW

