

Prevalence of resistant Enterobacterales, in particular ESBL-producing *Escherichia coli* in hospital wastewater discharged into the environment: Study of the Abidjan district, Côte d'Ivoire

ABSTRACT

Aims: Hospital effluent constitutes a reservoir for antibiotic-resistant bacteria. In particular, Gram-negative bacteria carrying several antibiotic resistance genes. This study was to evaluate the prevalence of antibiotic-resistant Enterobacterales and ESBL *Escherichia coli* in hospital effluents.

Methodology: In total, 112 wastewater samples from 4 hospital sites were collected and examined during 2020 at the Pasteur Institute of Côte d'Ivoire. Enumeration of total Enterobacterales was carried out on McConkey agar. McConkey agar was supplemented with 4 mg/l ceftazidime to detect resistant Enterobacterales. Antibiotic-resistant *Escherichia coli* were identified using MALDI-TOF MS. The production of ESBL was investigated by synergy tests on Mueller-Hinton agar.

Results: The results indicate a prevalence of resistant Enterobacterales varied between 4.6% and 2.6% with an average of 3.9%. A proportion of 69.2% of resistant Enterobacterales were strains of *Escherichia coli*. ESBL *Escherichia coli* represented a rate of 32.7% among resistant Enterobacterales. A rate of 47.3% of resistant *Escherichia coli* strains produced an ESBL. It appears from this study that the phenomenon of multi-resistance is undoubtedly present in bacteria isolated from hospital effluents, because the Enterobacterales isolated from these waters are mainly strains of ESBL-producing *Escherichia coli*.

Conclusion: Resistant Enterobacterales and particularly ESBL-producing *Escherichia coli* present in hospital effluents released without treatment into the environment are likely to cause public health problems. These bacteria should be considered environmental pollutants.

Keywords: Antibiotics, Enterobacteriaceae, *Escherichia coli*, resistance, ESBL, hospital effluent.

1. INTRODUCTION

Hospital wastewater is a highly complex effluent containing antibiotic compounds, metabolized drugs, disinfectants (Emmanuel et al., 2005), patient feces and microorganisms, potentially containing multidrug-resistant genes (Galvin et al., 2010) (Chagas et al., 2011). As such, hospital wastewater is considered a reservoir for antibiotic resistance, generating an environment conducive to the exchange of antibiotic resistance

genes. Gram-negative bacteria carrying multiple resistance genes are increasingly found in hospital wastewater (Zhang et al., 2012). Hospital effluents are discharged as common community wastewater to the wastewater treatment plant (WWTP), without any pre-treatment or cleaning process. After treatment, the water is discharged into surface waters. Some pathogenic microorganisms can remain longer in an aquatic environment, creating dissemination pathways and environmental reservoirs of antibiotic resistance genes (ARGs)(Perron et al., 2008). Over the past two decades, the presence of antibiotics in water bodies and the subsequent development of microbial resistance have attracted the attention of scientists and the public as a potential concern (Kümmerer, 2009) (Rodríguez-Gil et al., 2010). Hospital wastewater is one of the sources that contribute significantly to the environmental burden of antibiotics and, consequently, to antibiotic resistance (Zhang et al., 2009) (Davies & Davies, 2010) (Harris et al., 2012).

Enterobacterales, particularly *Escherichia coli*, are commensal bacteria present at approximately 10^8 colony-forming units per gram of human stool. In 2024, the WHO list of priority antibiotic-resistant pathogens specifically classifies Enterobacterales resistant to third-generation cephalosporins (3GCEs), including *Escherichia coli* and *Klebsiella pneumoniae*, as “critical priority”. The burden of these bacteria in AMR requires targeted action, especially in low-income countries (Da et al., 2023). In sub-Saharan Africa, several studies indicate *Escherichia coli* as the most common and most resistant Enterobacterales (Conforti et al., 2024). Extended-spectrum beta-lactamase-producing *Escherichia coli* (ESBL *E. coli*) are a real health problem for the populations of these countries because these strains escape medically relevant antibiotics and constitute essential supports in the propagation of resistance genes via mobile genetic elements (Traoré M., 2019). ESBL *E. coli* are commonly found in wastewater worldwide (Bréchet et al., 2015). Recently, authors have established the relationship between wastewater indicators and AMR carriage in the population. This approach shows the scope of wastewater surveillance to inform effective public health interventions (Guessennd et al., 2013). As in many sub-Saharan countries, the problem of ESBL-producing *Escherichia coli* is a concern in Côte d'Ivoire (Lorcy & Dubé, 2018). Indeed, despite the increasing prevalence of ESBL-producing Enterobacterales (BSE), particularly ESBL *E. coli*, their epidemiology is not yet well understood overall and the role of wastewater in their propagation is poorly documented (Laber et al., 1999). According to a study conducted at a university hospital in the country, liquid discharge from hospitals can contain most of the multi-resistant bacteria involved in nosocomial infections (Nuñez & Moreton, 2007). Reducing the spread of the most common multi-resistant bacteria (MRB) (ESBL *E. coli*) and that of the most resistant is one of the challenges facing health authorities in Côte d'Ivoire (Reinthal et al., 2003).

The objective of this work was to estimate the prevalence of resistant Enterobacterales, resistant *Escherichia coli* and BLSE *Escherichia coli* in hospital effluents in the district of Abidjan.

2. MATERIAL AND METHODS

2.1 COLLECTION OF HOSPITAL WASTEWATER

Hospital wastewater was collected from 4 hospital sites between January and December 2020, distributed as follows: 1 university hospital, 2 general hospitals and 1 specialized day hospital for tuberculosis (Figure 1).




FIG. 1. MAPPING OF HOSPITAL WASTEWATER SAMPLING SITES

2.2 SAMPLE SIZES

Twenty-eight campaigns were conducted overall. During each campaign, each of the 4 sites was collected once between 9:00 and 12:00, corresponding to a period of intense activity in

the hospitals. A total of 112 samples were collected during this study. Samples were manually collected in sterile vials from the main manholes (collecting the discharge from all other manholes in the hospital) and then transported within 3 hours, in coolers containing ice packs, to the laboratory for immediate analysis.

2.3 DETERMINATION OF THE PREVALENCE OF RESISTANT ENTEROBACTERIALES

A volume of 100 µl of water samples serially diluted to the 10th degree in saline was plated on McConkey agar without antibiotics to enumerate Enterobacteriales and on McConkey agar supplemented with 4 mg/L ceftazidime to enumerate resistant Enterobacteriales. The value of 4 mg/L was retained because it corresponds to the critical concentration above which resistance to ceftazidime in enterobacteriales is detected according to EUCAST recommendations, particularly during antibiogram in liquid medium .

Subsequently, the plates were incubated for 24 to 48 hours at 37°C. The percentage of resistant Enterobacteria was calculated by dividing the total number of colonies on McConkey medium with antibiotic by the number of colonies on McConkey medium without antibiotic.

2.4 DETERMINATION OF THE PREVALENCE OF RESISTANT ESCHERICHIA COLI

For the detection of antibiotic-resistant *Escherichia coli* strains, presumptive Enterobacteriales colonies on McConkey agar supplemented with 4 mg/l ceftazidime were randomly selected on the basis of their macroscopic differences. Five colonies of each morphology type were identified using MALDI–TOF MS (VITEK® MS, BioMérieux, Marcy-l'Étoile, France). The prevalence of resistant *Escherichia coli* was calculated by dividing the number of *Escherichia coli* identified with MALDI–TOF MS by the number of enterobacteriales identified with MALDI–TOF MS.

2.5 DETERMINATION OF THE PREVALENCE OF BLSE-PRODUCING ESCHERICHIA COLI (E. COLI BLSE):

The production of ESBL among the isolated *Escherichia coli* strains was investigated using synergy tests on Mueller-Hinton agar by agar diffusion technique, as previously described (Nimri & Azaizeh, 2012) . The presence of an ESBL was confirmed between a third-generation cephalosporin (C3G) disc (cefotaxime, ceftazidime, cefepime) and amoxicillin + clavulanic acid separated by 3 cm from center to center. The appearance of a synergy image called "champagne cork" between these discs allowed the confirmation of the production of a broad-spectrum beta-lactamase. Subsequently, the number of ESBL *E. coli* was divided respectively by the number of resistant Enterobacteriaceae and by the number of resistant *E. coli* in order to determine the prevalence of ESBL *E. coli* in relation to these 2 sets

2.5 STATISTICAL DATA ANALYSIS

Statistical analysis of the data was performed using IBM SPSS statistics 27 and GraphPad Prism 9.5.0 (730) software (San Diego, California, USA). Values were given as percentages.

3. RESULTS

The prevalence of resistant Enterobacterales in hospital effluents varied between 4.6% and 2.6% with a mean of 3.9% (Table 1).

Table 1. Prevalence of resistant enterobacteria in hospital effluents

Collection sites	Count (CFU/ml)		Prevalence of resistant enterobacteria (%)
	Total Enterobacteria	Resistant Enterobacteria	
Site 1	3.8 10 ⁶	1,75 10 ⁵	4.6
Site 2	3.8 10 ⁶	1,47 10 ⁵	3.9
Site 3	3.66 10 ⁶	1,61 10 ⁵	4.4
Site 4	2.86 10 ⁶	7,48 10 ⁴	2.6
Average	3,53 10 ⁶	1,38 10 ⁵	3.9

Approximately 69.2% of resistant Enterobacterales in hospital effluents are strains of *Escherichia coli* (Table 2).

Table 2. MALDI-TOF MS identification of *Escherichia coli* strains within the population of resistant enterobacterales

Resistant Enterobacterales	Effective MALDI-TOF MS Identification (N)	Prevalence (%)
<i>Escherichia coli</i> species	296	69.2
Non- <i>Escherichia coli</i> species	132	30.8
Total	428	100

32.7% of resistant enterobacterales in hospital effluents are strains of ESBL *Escherichia coli* (Table 3)

Table 3. Distribution of ESBL *Escherichia coli* strains in the population of resistant Enterobacterales

Resistant Enterobacterales	Staff (N)	Prevalence (%)
<i>Escherichia coli</i> ESBL	140	32.7
Resistant enterobacterales other than <i>Escherichia coli</i> ESBL	288	67.3
Total	428	100

47.3% of resistant *Escherichia coli* strains found in hospital effluents produce ESBL (Table 4)

Table 4. Distribution of ESBL *Escherichia coli* strains in the population of resistant enterobacterales

Resistant <i>Echerichia coli</i>	Staff (N)	Prevalence (%)
<i>Escherichia coli</i> ESBL	140	47.3
Non- ESBL <i>Escherichia coli</i>	156	52.7
Total	296	100

4. DISCUSSION

Hospitals are places where the use of antibiotics and the density of individuals favor the dissemination of antibiotic resistance (Russell, 2001). The operation of these structures is accompanied by the discharge of a high quantity of wastewater containing pathogens mainly from patients (Wellington et al., 2013) (LaPara et al., 2011). These germs can be exposed to a wide variety of antibacterials contained for the development of resistance via cellular mutation or acquisition of genetic elements in the form of plasmids or transposons (Conforti et al., 2024). Studies indicate that even when treated by wastewater treatment plants (WWTP), hospital effluents constitute a source of release of antibiotic-resistant pathogens into the downstream environment (Wellington et al., 2013) (LaPara et al., 2011). Wastewater monitoring is a cost-effective and less burdensome way to track changes in RAM carriage within a community (Conforti et al., 2024). In countries of the South, particularly in sub-Saharan Africa where antimicrobial resistance is less well documented, this approach could be an alternative to this problem (Da et al., 2023).

Prevalence of resistant Enterobacterales in discharged hospital wastewater

In wastewater, little information is available on RAM levels in other bacterial populations because most studies in this area have been carried out using coliforms as bacterial indicators (Guardabassi et al., 2002). The results obtained in this study confirm the presence of bacteria of the Enterobacterales family such as *Escherichia coli*, in hospital wastewater. The results of this study show that approximately $1.47 \cdot 10^5$ CFU/ml or 3.9% of the Enterobacterales contained in the wastewater continuously discharged into the environment by these structures are resistant. This prevalence of resistant Enterobacterales is much lower than that of 77.41% noted by authors in samples from rivers influenced by strong anthropogenic pressure characterized by the nearby presence of a hospital, a significant concentration of population and refugees, associated or not with agricultural activity (Hobeika, 2021). Similarly, other studies have shown that approximately 15% of clinical strains of Enterobacterales in hospitals could be resistant to 3rd generation cephalosporins (3GC) with high prevalences recorded in intensive care units (48%), hematology-oncology departments (27%) and the pediatric department (25%) (Mkaouar et al., 2008). Once in hospital effluents, these Enterobacterales have the power to easily transfer genetic information between themselves or with other bacteria in the environment (Yala et al., 2024). The low prevalence of resistant Enterobacterales observed in this study could be related to the probable presence of residues of specific substances in the effluents as suggested by Nunez and Moretton (Nuñez & Moretton, 2007). According to some studies, the prevalence of resistant bacteria in wastewater can vary considerably depending on several factors including the physicochemical parameters and the analytical methods used (Guardabassi et al., 2002) (Gyorgy & Sandor, 2010). For others, this difference in resistance rates can be explained by the variability of epidemiological factors, antibiotic usage policies and levels of hospital hygiene from one country to another (Mkaouar et al., 2008). The application of infection limitation provisions and the determination of the mechanism underlying resistance to 3GC in Enterobacterales are essential to control their dissemination at the health and community level (Yala et al., 2024).

Prevalence of Escherichia coli in the population of resistant Enterobacterales isolated

Among the bacteria isolated from hospital wastewater, the species *Escherichia coli* occupies an important place with a proportion of 69.2%. This high proportion could be explained by the predominance of *Escherichia coli* commonly isolated among the Enterobacterales. Indeed, studies conducted in Togo found *Escherichia coli* (63.93%) as the most frequently isolated species in the group of Enterobacterales followed by *Klebsiella* (22.86%) (Salah et al., 2021). This trend is confirmed by other authors, notably by Ebongue et al. in Cameroon (48.5%) and Kalambry et al. in Mali (44.7%) (Ebongue et al., 2015) (Kalambry et al., 2019). *Escherichia coli* is an Enterobacterales present in the digestive tract and involved in infections, mainly urinary. As such, the bacterium is frequently the therapeutic target of antibiotics (Sonnet, 2020) (Public Health France a, 2023). Third-generation cephalosporins (3GCs) occupy an essential place in the therapeutic arsenal in human health. Thus, the control of resistance to these molecules in Enterobacterales such as *Escherichia coli* is a concern both in hospital and community settings. If in Europe, the average proportion of resistance to 3GCs in *Escherichia coli* decreased from 15.7% to 14.3% between 2018-2022, this is not yet the case in most developing countries (Public Health France a, 2023) (Tagajdid et al., 2010). The share of resistant *Escherichia coli* in this study shows the interest in determining the impact of hospital effluent discharges on the contamination of aquatic ecosystems downstream, particularly at the fauna level.

Prevalence of ESBL Escherichia coli in the population of isolated resistant Escherichia coli

The ESBL *Escherichia coli* rate obtained in this study is 32.7%. The presence of ESBL *Escherichia coli* in hospital effluents has been reported in several studies conducted worldwide (Guessennd et al., 2013) (Cahill et al., 2019) (Fadare & Okoh, 2021). This rate could be explained, on the one hand, by a higher frequency and density of carriage among hospitalized patients (Mkaouar et al., 2008) (Ruppé et al., 2013). On the other hand, the high consumption of antibiotics in hospitals leads to their elimination in the form of compounds that are still active in wastewater. These compounds exert a selection pressure favoring resistant bacteria including ESBL *Escherichia coli*. The presence of disinfectants, widely used in hospitals, can also contribute to the selection of resistant bacteria (Coutu et al., 2013).

Analysis of Table 4 shows a rate of 47.3% of resistant *Escherichia coli* strains found in hospital effluents producing an ESBL. Indeed, in recent years, several studies have highlighted the presence of ESBL *Escherichia coli* strains in hospital effluents, both in developing and developed countries (Drieux-Rouzet & Jarlier, 2014). In their study, Galvin and colleagues investigated the impact of hospital effluent by quantifying antibiotic-sensitive and antibiotic-resistant *Escherichia coli* strains in effluent samples collected at different levels at the discharge of an Irish hospital, in the municipal network upstream and downstream of the hospital sewer connection. They showed that the proportion of antibiotic-resistant *Escherichia coli* strains found in the municipal network was higher downstream of the hospital connection than upstream of the hospital connection (Galvin et al., 2010). These data suggest the importance of organizing the monitoring of antibiotic consumption and bacterial resistance at the environmental level as well as in human and animal health (Public Health France b, 2023).

5. CONCLUSION

This study shows that the phenomenon of antimicrobial resistance is undoubtedly present in bacteria isolated from hospital effluent wastewater, because the germs isolated from these waters are mainly strains of *Escherichia coli* producing ESBL. These effluents discharged without treatment into the lagoon can constitute a source of dissemination of these potentially pathogenic bacteria. Controlling ESBL-producing *Escherichia coli* strains in water remains a public health issue in Côte d'Ivoire. These bacterial strains are the cause of many therapeutic failures. More rigorous environmental management is more than necessary. On the one hand, improving wastewater treatment processes and on the other hand decontaminating hospital effluents by physicochemical and microbiological processes is therefore essential. Also, quantification and genotypic characterization by searching for resistance genes would make it possible to assess the risks associated with possible transfers of resistance genes and infections

COMPETING INTERESTS

The authors declare that they have no conflict of interest

ETHICAL APPROVAL

Access to the various hospital sites for sampling and analysis of discharged effluents was subject to written authorizations from the managers of these structures. These authorizations are kept with the authors.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

- Bréchet, C., Hocquet, D., & Bertrand, X. (2015). Eaux usées et *Escherichia coli* producteur de β -lactamases à spectre étendu. *Journal des Anti-infectieux*, 17.
<https://doi.org/10.1016/j.antinf.2015.03.001>
- Cahill, N., O'Connor, L., Mahon, B., Varley, Á., McGrath, E., Ryan, P., Cormican, M., Brehony, C., Jolley, K. A., Maiden, M. C., Brisse, S., & Morris, D. (2019). Hospital effluent : A reservoir for carbapenemase-producing Enterobacterales? *The Science*

of the Total Environment, 672, 618-624.

<https://doi.org/10.1016/j.scitotenv.2019.03.428>

- Chagas, T. P. G., Seki, L. M., Cury, J. C., Oliveira, J. a. L., Dávila, A. M. R., Silva, D. M., & Asensi, M. D. (2011). Multiresistance, beta-lactamase-encoding genes and bacterial diversity in hospital wastewater in Rio de Janeiro, Brazil. *Journal of Applied Microbiology*, 111(3), 572-581. <https://doi.org/10.1111/j.1365-2672.2011.05072.x>
- Conforti, S., Holschneider, A., Sylvestre, É., & Julian, T. R. (2024). Monitoring ESBL-*Escherichia coli* in Swiss wastewater between November 2021 and November 2022 : Insights into population carriage. *mSphere*, 9(5), e0076023. <https://doi.org/10.1128/msphere.00760-23>
- Coutu, S., Rossi, L., Barry, D. A., Rudaz, S., & Vernaz, N. (2013). Temporal variability of antibiotics fluxes in wastewater and contribution from hospitals. *PloS One*, 8(1), e53592. <https://doi.org/10.1371/journal.pone.0053592>
- Da, L., Somé, D., Yehouenou, C., Somé, C., Zoungrana, J., Ouédraogo, A. S., Lienhardt, C., & Poda G.E.A. (2023). État des lieux de la résistance aux antibiotiques en Afrique subsaharienne. *Médecine et Maladies Infectieuses Formation*, 2(1), 3-12. <https://doi.org/10.1016/j.mmifmc.2023.01.003>
- Davies, J., & Davies, D. (2010). Origins and evolution of antibiotic resistance. *Microbiology and Molecular Biology Reviews: MMBR*, 74(3), 417-433. <https://doi.org/10.1128/MMBR.00016-10>
- Drieux-Rouzet, L., & Jarlier, V. (2014). Bactéries multirésistantes dans l'eau : Modèles des entérobactéries productrices de bêta-lactamase à spectre étendu. *Revue Francophone des Laboratoires. mars 2014;2014(460):75-9.*
- Ebongue, C. O., Tsiatzok, M. D., Mefo'o, J. P. N., Ngaba, G. P., Beyiha, G., & Adio, D. (2015). Evolution de la résistance aux antibiotiques des entérobactéries isolées à l'Hôpital Général de Douala de 2005 à 2012. *The Pan African Medical Journal*, 20(227), Article 227. <https://doi.org/10.11604/pamj.2015.20.227.4770>

Emmanuel, E., Perrodin, E., Keck, G., Blanchard, J., & Vermande, P. (2005).

Ecotoxicological risk assessment of hospital wastewater : A proposed framework for raw effluents discharging into urban sewer network—PubMed.

<https://pubmed.ncbi.nlm.nih.gov/15621348/>

Fadare, F. T., & Okoh, A. I. (2021). Distribution and molecular characterization of ESBL, pAmpC β -lactamases, and non- β -lactam encoding genes in Enterobacteriaceae isolated from hospital wastewater in Eastern Cape Province, South Africa. *PloS One*, 16(7), e0254753. <https://doi.org/10.1371/journal.pone.0254753>

Galvin, S., Boyle, F., Hickey, P., Vellinga, A., orris, C., & Cormican, M. (2010).

Dénombrement et caractérisation des bactéries Escherichia coli résistantes aux antimicrobiens dans les effluents des stations d'épuration municipales, hospitalières et secondaires—PubMed. <https://pubmed.ncbi.nlm.nih.gov/20525867/>

Guardabassi, L., Lo Fo Wong, D. M. A., & Dalsgaard, A. (2002). The effects of tertiary wastewater treatment on the prevalence of antimicrobial resistant bacteria. *Water Research*, 36(8), 1955-1964. [https://doi.org/10.1016/S0043-1354\(01\)00429-8](https://doi.org/10.1016/S0043-1354(01)00429-8)

Guessennd, N. K., Ouattara, M. B., Ouattara, N. D., Nevry, R. K., Gbanon, V., Tiekoura, K.

B., Dosso, M., & Ger, B. M. R. (2013). Étude des bactéries multirésistantes des effluents hospitaliers d'un centre hospitalier et universitaire (CHU) de la ville d'Abidjan (Côte d'Ivoire). *Journal of Applied Biosciences*, 69, 5456-5464.

<https://doi.org/10.4314/jab.v69i0.95071>

Gyorgy, H., & Sandor, S. (2010). *The importance of pseudomonas aeruginosa on environmental safety.*

Harris, S., Cormican, M., & De cummins, F. (2012). *Résidus d'antimicrobiens et bactéries résistantes aux antimicrobiens : Impact sur l'environnement microbien et risque pour la santé humaine—Une revue | Demander un PDF.*

https://www.researchgate.net/publication/241682632_Antimicrobial_Residues_and_Antimicrobial-

Resistant_Bacteria_Impact_on_the_Microbial_Environment_and_Risk_to_Human_Health_-_A_Review

- Hobeika, W. (2021). *Dissémination sous pression anthropique et de la saisonnalité de bactéries résistantes dans les fleuves du Liban*.
- Kalambry, A., Gaudré, N., Dramé, B., Poudiogo, A., Kassogué, A., Koné, H., & Diarra, A. (2019). *Profil de résistance aux bêta-lactamines des entérobactéries isolées des prélèvements urinaires à l'Hôpital du Mali*. *ome* 14.
- Kümmerer, K. (2009). Antibiotics in the aquatic environment—A review—Part II. *Chemosphere*, 75(4), 435-441. <https://doi.org/10.1016/j.chemosphere.2008.12.006>
- Laber, Haberl, R., & Shrestha, R. (1999). Two-stage constructed wetland for treating hospital wastewater in Nepal | Request PDF. *ResearchGate*. [https://doi.org/10.1016/S0273-1223\(99\)00447-3](https://doi.org/10.1016/S0273-1223(99)00447-3)
- LaPara, T. M., Burch, T. R., McNamara, P. J., Tan, D. T., Yan, M., & Eichmiller, J. J. (2011). Tertiary-treated municipal wastewater is a significant point source of antibiotic resistance genes into Duluth-Superior Harbor. *Environmental Science & Technology*, 45(22), 9543-9549. <https://doi.org/10.1021/es202775r>
- Lorcy, A., & Dubé, E. (2018). Les enjeux des bactéries multi-résistantes à l'hôpital. *Anthropologie & Santé. Revue internationale francophone d'anthropologie de la santé*, 16, Article 16. <https://doi.org/10.4000/anthropologiesante.2825>
- Mkaouar, D., Mahjoubi, F., Mezghani, S., Znazen, A., Ktari, S., & Hammami, A. (2008). Étude de la résistance des entérobactéries aux céphalosporines de troisième génération dans les hôpitaux de Sfax, Tunisie (1999–2005). *Médecine et Maladies Infectieuses*, 38(6), 293-298. <https://doi.org/10.1016/j.medmal.2007.11.017>
- Nimri, L. F., & Azaizeh, B. A. (2012). First Report of Multidrug-Resistant ESBL-Producing Urinary Escherichia coli in Jordan. *Microbiology Research Journal International*, 71-81. <https://doi.org/10.9734/BMRJ/2012/1360>

- Nuñez, L., & Moretton, J. (2007). *SciELO Brazil—Disinfectant-resistant bacteria in Buenos Aires city hospital wastewater Disinfectant-resistant bacteria in Buenos Aires city hospital wastewater*. <https://www.scielo.br/j/bjm/a/PBRsGt9tGgJKL53PD6FLYNc/>
- Perron, G. G., Gonzalez, A., & Buckling, A. (2008). The rate of environmental change drives adaptation to an antibiotic sink. *Journal of Evolutionary Biology*, *21*(6), 1724-1731. <https://doi.org/10.1111/j.1420-9101.2008.01596.x>
- Reinthal, F. F., Posch, J., Feierl, G., Wüst, G., Haas, D., Ruckebauer, G., Mascher, F., & Marth, E. (2003). Antibiotic resistance of *E. coli* in sewage and sludge. *Water Research*, *37*(8), 1685-1690. [https://doi.org/10.1016/S0043-1354\(02\)00569-9](https://doi.org/10.1016/S0043-1354(02)00569-9)
- Rodríguez-Gil, J. L., Catalá, M., Alonso, S. G., Maroto, R. R., Valcárcel, Y., Segura, Y., Molina, R., Melero, J. A., & Martínez, F. (2010). Heterogeneous photo-Fenton treatment for the reduction of pharmaceutical contamination in Madrid rivers and ecotoxicological evaluation by a miniaturized fern spores bioassay. *Chemosphere*, *80*(4), 381-388. <https://doi.org/10.1016/j.chemosphere.2010.04.045>
- Ruppé, E., Lixandru, B., Cojocar, R., Büke, C., Paramythiotou, E., Angebault, C., Visseaux, C., Djuikoue, I., Erdem, E., Burduniuc, O., El Mniai, A., Marcel, C., Perrier, M., Kesteman, T., Clermont, O., Denamur, E., Armand-Lefèvre, L., & Andremont, A. (2013). Relative fecal abundance of extended-spectrum- β -lactamase-producing *Escherichia coli* strains and their occurrence in urinary tract infections in women. *Antimicrobial Agents and Chemotherapy*, *57*(9), 4512-4517. <https://doi.org/10.1128/AAC.00238-13>
- Russell, A. D. (2001). Mechanisms of bacterial insusceptibility to biocides. *American Journal of Infection Control*, *29*(4), 259-261. <https://doi.org/10.1067/mic.2001.115671>
- Salah, F. D., Sadji, A. Y., Akolly, K., Bidjada, B., Awoussi, K. S., Abaya, A. M., Amouzou-Glikpa, A., Kutoati, A., Amegan, L., Palanga, K. K., Yanogo, P. K., Kouawo, M. L., Otshudiandjeka, J., Sawadogo, B., Diallo-Ouedraogo, F., Layibo, Y., Patassi, A., Halatoko, W. A., Simporé, J., & Saka, B. (2021). Augmentation de la résistance aux

antibiotiques des Entérobactéries isolées à l'Institut National d'Hygiène de Lomé de 2010 à 2017. *Journal of Interventional Epidemiology and Public Health*, 4(3), Article 3. <https://doi.org/10.11604/JIEPH.suppl.2021.4.3.1089>

Santé publique France a. (2023, décembre). *Résistance aux antibiotiques*.

<https://www.santepubliquefrance.fr/maladies-et-traumatismes/infections-associees-aux-soins-et-resistance-aux-antibiotiques/resistance-aux-antibiotiques>

Santé publique France b. (2023). *Prévention de la résistance aux antibiotiques : Une démarche « Une seule santé »*. Novembre 2023.

<https://www.santepubliquefrance.fr/maladies-et-traumatismes/infections-associees-aux-soins-et-resistance-aux-antibiotiques/resistance-aux-antibiotiques/documents/rapport-synthese/prevention-de-la-resistance-aux-antibiotiques-une-demarche-une-seule-sante-novembre-2023>

Sonnet, M. (2020). *Les entérobactéries face aux céphalosporines de 3ème génération et carbapénèmes : Évolution, actualité, problématique*.

https://matheo.uliege.be/bitstream/2268.2/9758/6/SONNET_Mirabel_TFE_FMV_juin_2020_provisoire.pdf

Tagajdid, M. R., Boumhil, L., Iken, M., Adnaoui, M., & Benouda, A. (2010). Étude de la résistance des souches d' *Escherichia coli* isolées dans les urines aux fluoroquinolones et aux céphalosporines de. *Médecine et Maladies Infectieuses*, 40(2), 70-73. <https://doi.org/10.1016/j.medmal.2008.10.015>

Traoré, M. (2019). La résistance aux antimicrobiens fait peur à la Côte d'Ivoire. *Afrique Sub-Saharienne*. <https://www.scidev.net/afrique-sub-saharienne/news/resistance-antimicrobiens-inquietante-cote-d-ivoire-09122019/>

Wellington, E. M., Boxall, A. B., Cross, P., Feil, E. J., Gaze, W. H., Hawkey, P. M., Johnson-Rollings, A. S., Jones, D. L., Lee, N. M., Otten, W., Thomas, C. M., & Williams, A. P. (2013). The role of the natural environment in the emergence of antibiotic resistance

in Gram-negative bacteria. *The Lancet Infectious Diseases*, 13(2), 155-165.

[https://doi.org/10.1016/S1473-3099\(12\)70317-1](https://doi.org/10.1016/S1473-3099(12)70317-1)

Yala, J.-F., Mabika, M. R., Boucka, B. D., & Lepengue, A. N. (2024). (PDF) Étude phénotypique de la résistance des bactéries isolées des eaux des lacs et rivières de la ville de Franceville aux céphalosporines de troisième génération. *ResearchGate*.
<https://doi.org/10.4314/jab.v119i1.12>

Zhang, X., Lü, X., & Zong, Z. (2012). Enterobacteriaceae producing the KPC-2 carbapenemase from hospital sewage. *Diagnostic Microbiology and Infectious Disease*, 73(2), 204-206. <https://doi.org/10.1016/j.diagmicrobio.2012.02.007>

Zhang, X.-X., Zhang, T., & Fang, H. H. P. (2009). Antibiotic resistance genes in water environment. *Applied Microbiology and Biotechnology*, 82(3), 397-414.
<https://doi.org/10.1007/s00253-008-1829-z>

UNDER PEER REVIEW