

## **Improving maize yield and soil productivity through N management practices in maize-legume Intercropping**

### **Abstract**

A field experiment was conducted to investigate the effects of maize intercropping and nitrogen management on crop productivity. The study comprised two factors: maize intercropping with five treatments (sole maize, skipped row maize, and maize intercropped with greengram, blackgram, or cluster bean) and nitrogen management with three treatments (100%, 75%, and 50% of the recommended nitrogen dose). The results showed that sole maize at 60 x 20 cm spacing and maize intercropped with cluster bean, blackgram, or greengram significantly outyielded sole maize in skipped rows. Nitrogen application at 100% of the recommended dose resulted in higher maize yields, while intercropping with cluster bean showed the highest maize grain equivalent yield and land equivalent ratio. The study highlights the potential of maize intercropping and optimized nitrogen management to enhance crop productivity, reduce soil nutrient depletion, and promote sustainable agriculture practices.

**Keywords:** maize intercropping, nitrogen management, crop productivity, sustainable agriculture.

### **Introduction**

The sustainable productivity level of crops is the need of hour in the present Indian farming. Hence, we must examine every possibility for crop intensification with sustainable nutrition for achieving the sustainability. Maize (*Zea mays* L.) is the third most important cereal crop after wheat and rice in the world, and India is the seventh largest producer of maize. Due to its multidimensional uses, high production potential and adaptability to wide range of environment, we can expect maize to assume a proportionally large and more important role in the food production in coming years. Under crop intensification intercropping with maize is gaining popularity during recent past in southern agro climatic zone of Andhra Pradesh. Hybrid maize being an exhaustive crop, requires high quantity of nutrients particularly nitrogen. The recent maize hybrids are responding to more than 240 kg N ha<sup>-1</sup>. But, the prohibitive cost of fertilizer nitrogen limits the farmers for its liberal application. In the present day's concern about environmental degradation coupled with high cost of nitrogen, there is a need to find out supplemental alternative sources. Legumes, if associated with maize can reduce nitrogen requirement to some extent, besides maintaining soil health. Though considerable quantum of research has been carried out in maize nutrition, much of them was resource exploitive type, but not conserving type and they were unable to answer, proportionate fertilizer nitrogen saving to hybrid maize when legumes are associated. Traditionally, nitrogen recommendations for maize-based intercropping systems have been determined by the response of individual crops, rather than considering the intercropping system as a whole. This approach can lead to suboptimal nitrogen management, resulting in reduced economic returns. By adopting a rational agronomic approach of uniform plant population of both base and inter crop, the production potential and economic viability of maize + legume inter cropping system depends upon nature and type of associated legume and quantity of fertilizer applied. Hence, the present

study was designed with following objectives to explore the possibilities of intercropping of short duration legumes under variable doses of nitrogen to maize and intercropping sequence under different crop geometry.

**Material and Methods:**

The experiment was conducted at the Agricultural Research Farm, Sri CB Singh Memorial Shikshan Sansthan, Jhinhak, Kanpur Dehat, U.P., India, during the Kharif season of 2022-23. The farm is situated in the Indo-Gangetic Plains, between 26.35° N latitude and 80.09° E longitude, with an elevation of 130.00 m above mean sea level. A randomized block design with a factorial concept was employed, with three replications. The treatments consisted of two factors: maize intercropping systems and nitrogen management. The maize intercropping systems included five treatments: sole maize at 60 x 20 cm spacing, skipped row maize, maize + greengram intercropping, maize + blackgram intercropping, and maize + cluster bean intercropping. The nitrogen management factor consisted of three treatments: recommended dose of nitrogen (240 kg ha-1), 75% recommended dose of nitrogen (180 kg ha-1), and 50% recommended dose of nitrogen (120 kg ha-1). Sole crops of greengram, blackgram, and cluster bean were raised separately at 30 x 10 cm spacing outside the experimental layout. The plot size was 6.0 x 5.4 m (gross) and 4.0 x 3.6 m (net). Intercrops were sown at a spacing of 60 x 5 cm. A uniform dose of 60 kg P2O5 ha-1 and 50 kg K2O ha-1 was applied as basal to maize in all plots. Nitrogen was applied in three equal splits, viz., basal, knee stage, and tasseling stage. For the intercrops, 20, 50, and 40 kg N, P2O5, and K2O ha-1, respectively, were applied as basal application at sowing.

**Results and Discussion**

The results of the study revealed that maize intercropping and nitrogen management significantly influenced the growth parameters, yield attributes, and yield of maize. The growth parameters, such as plant height, leaf area index, and dry matter production, were significantly higher in sole maize at 60 x 20 cm spacing, which was at par with maize + cluster bean, maize + blackgram, and maize + greengram intercropping. The yield attributes, such as cob length and girth, number of grains cob-1, and 100 grain weight, were higher in sole maize at 60 x 20 cm planting, closely followed by maize + cluster bean, maize + blackgram, and maize + greengram intercropping. The grain and stover yield of maize was highest in sole maize at 60 x 20 cm, which was at par with the yields of maize + cluster bean, maize + blackgram, and maize + greengram intercropping. The application of 100% recommended dose of nitrogen to maize resulted in significant higher yield attributes and grain and stover yield, while application of 50% recommended dose of nitrogen resulted in significantly lower yield attributes and grain and stover yield.

**Table 1: Plant height (cm) of maize as influenced by maize + legume intercropping and nitrogen management**

Treatment	25 DAS	50 DAS	75 DAS
<b>Maize intercropping</b>			
T <sub>1</sub> : Maize 60 x 20 cm	52.6	126.9	164.9
T <sub>2</sub> : Maize skipped row	48.3	109.8	142.6
T <sub>3</sub> : T <sub>2</sub> + Greengram	48.2	121.6	158.0
T <sub>4</sub> : T <sub>2</sub> + Blackgram	48.6	124.8	158.4
T <sub>5</sub> : T <sub>2</sub> + Cluster bean	52.1	125.1	161.6

SEm ±	2.36	3.61	4.02
CD (P=0.05)	NS	10.3	11.5
<b>Nitrogen management</b>			
N <sub>1</sub> : 100 % Rec. N to maize	52.4	124.2	164.6
N <sub>2</sub> : 75 % Rec. N to maize	50.3	119.2	153.9
N <sub>3</sub> : 50 % Rec. N to maize	48.0	105.3	141.6
SEm ±	1.83	4.01	4.23
CD (P=0.05)	NS	11.5	12.1

**Table 2 : Leaf area index of maize as influenced by maize + legume intercropping and nitrogen management**

Treatment	25 DAS	50 DAS	75 DAS	Harvest
<b>Maize intercropping</b>				
T <sub>1</sub> : Maize 60 x 20 cm	0.212	2.50	1.49	1.28
T <sub>2</sub> : Maize skipped row	0.181	1.94	1.20	1.20
T <sub>3</sub> : T <sub>2</sub> + Greengram	0.199	2.36	1.36	1.25
T <sub>4</sub> : T <sub>2</sub> + Blackgram	0.206	2.39	1.39	1.26
T <sub>5</sub> : T <sub>2</sub> + Cluster bean	0.212	2.41	1.42	1.28
SEm ±	0.07	0.14	0.048	0.014
CD (P=0.05)	NS	0.40	0.14	0.04
<b>Nitrogen management</b>				
N <sub>1</sub> : 100 % Rec. N to maize	0.217	2.38	1.36	1.11
N <sub>2</sub> : 75 % Rec. N to maize	0.201	2.36	1.29	1.09
N <sub>3</sub> : 50 % Rec. N to maize	0.179	2.01	1.14	0.99
SEm ±	0.006	0.108	0.037	0.010
CD (P=0.05)	0.017	0.31	0.11	0.03

**Table 3 : Dry matter production (kg ha<sup>-1</sup>) of maize as influenced by maize + legume intercropping**

and nitrogen management

Treatment	25 DAS	50 DAS	75 DAS	Harvest
<b>Maize intercropping</b>				
T <sub>1</sub> : Maize 60 x 20 cm	421.7	4311	7257	9174
T <sub>2</sub> : Maize skipped row	408.0	4004	6655	8763
T <sub>3</sub> : T <sub>2</sub> + Greengram	412.8	4296	6987	8917
T <sub>4</sub> : T <sub>2</sub> + Blackgram	417.6	4298	7056	9003
T <sub>5</sub> : T <sub>2</sub> + Cluster bean	419.6	4303	7240	9103
SEm ±	4.8	90.9	112.0	104.2
CD (P=0.05)	NS	262	323	301
<b>Nitrogen management</b>				
N <sub>1</sub> : 100 % Rec. N to maize	420.1	4199	6987	9314
N <sub>2</sub> : 75 % Rec. N to maize	418.8	4109	6763	9074
N <sub>3</sub> : 50 % Rec. N to maize	404.2	3802	6213	8517
SEm ±	3.61	102.5	86.7	90.2
CD (P=0.05)	10.6	297	251	260

Table 4 : Yield attributes of maize as influenced by maize + legume intercropping and nitrogen management

Treatment	Cob length (cm)	Cob girth (cm)	No. of grains per cob	100 grain weight (g)
<b>Maize intercropping</b>				
T <sub>1</sub> : Maize 60 x 20 cm	14.4	14.7	333	28.8
T <sub>2</sub> : Maize skipped row	12.1	12.9	252	23.8

T <sub>3</sub> : T <sub>2</sub> + Greengram	13.7	13.4	297	26.4
T <sub>4</sub> : T <sub>2</sub> + Blackgram	13.9	14.4	299	27.5
T <sub>5</sub> : T <sub>2</sub> + Cluster bean	14.4	14.6	324	27.7
SEm ±	0.74	0.54	26.83	1.15
CD (P=0.05)	2.1	1.6	77	3.3
<b>Nitrogen management</b>				
N <sub>1</sub> : 100 % Rec. N to maize	14.8	14.6	333	28.1
N <sub>2</sub> : 75 % Rec. N to maize	14.5	14.3	324	27.7
N <sub>3</sub> : 50 % Rec. N to maize	12.3	13.0	245	24.7
SEm ±	0.57	0.42	20.78	0.88
CD (P=0.05)	1.6	1.2	60	2.5

**Table 5 : Grain yield, stover yield and harvest index of maize as influenced by maize + legume intercropping and nitrogen management**

Treatment	Grain yield (kg ha <sup>-1</sup> )	Stover yield (kg ha <sup>-1</sup> )	Harvest index
<b>Maize intercropping</b>			
T <sub>1</sub> : Maize 60 x 20 cm	4009	5921	40.4
T <sub>2</sub> : Maize skipped row	3071	4277	41.8
T <sub>3</sub> : T <sub>2</sub> + Greengram	3791	5685	40.0
T <sub>4</sub> : T <sub>2</sub> + Blackgram	3864	5771	40.1
T <sub>5</sub> : T <sub>2</sub> + Cluster bean	3905	5825	40.1
SEm ±	86.5	93.7	1.23
CD (P=0.05)	250	271	NS

<b>Nitrogen management</b>			
N <sub>1</sub> : 100 % Rec. N to maize	3838	5809	39.7
N <sub>2</sub> : 75 % Rec. N to maize	3791	5694	40.0
N <sub>3</sub> : 50 % Rec. N to maize	3055	4984	38.0
SEm ±	79.2	83.4	1.01
CD (P=0.05)	229	241	NS

**Table 6 : Biometric observations of greengram as influenced by maize + legume intercropping and nitrogen management**

<b>Treatment</b>	<b>Plant height (cm)</b>			<b>Leaf area index</b>			<b>Dry matter production (kg ha<sup>-1</sup>)</b>		
	<b>25 DAS</b>	<b>50 DAS</b>	<b>Harvest</b>	<b>25 DAS</b>	<b>50 DAS</b>	<b>Harvest</b>	<b>25 DAS</b>	<b>50 DAS</b>	<b>Harvest</b>
(Maize + greengram) + N <sub>1</sub>	10.3	59.4	59.6	0.32	2.89	2.02	110	2401	2492
(Maize + greengram) + N <sub>2</sub>	10.4	59.1	59.1	0.30	2.80	1.98	96	2398	2413
(Maize + greengram) + N <sub>3</sub>	10.3	59.0	59.0	0.30	2.80	1.96	96	2346	2409
Sole greengram	10.5	60.2	59.6	0.33	3.02	2.63	114	2462	2562

N<sub>1</sub>, N<sub>2</sub> and N<sub>3</sub> are 100, 75 and 50 % recommended doses of nitrogen to maize respectively.

**Table 7 : Yield attributes and yield of greengram as influenced by maize + legume intercropping and nitrogen management**

<b>Treatment</b>	<b>Yield attributes</b>				<b>Yield</b>	
	<b>No. of clusters per plant</b>	<b>No. of pods per cluster</b>	<b>No. of seeds per pod</b>	<b>1000 seed weight (g)</b>	<b>Seed yield (kg ha<sup>-1</sup>)</b>	<b>Haulm yield (kg ha<sup>-1</sup>)</b>
(Maize + greengram) + N <sub>1</sub>	11.0	5.3	5.5	37.5	418	860
(Maize + greengram) + N <sub>2</sub>	10.8	4.9	5.2	37.1	394	854

(Maize + greengram) + N <sub>3</sub>	9.6	5.4	4.9	36.1	384	851
Sole greengram	11.6	5.5	5.4	38.0	425	860

N<sub>1</sub>, N<sub>2</sub> and N<sub>3</sub> are 100, 75 and 50 % recommended doses of nitrogen to maize, respectively

**Table 8 : Biometric observations of blackgram as influenced by maize + legume intercropping and nitrogen management**

Treatment	Plant height (cm)			Leaf area index			Dry matter production (kg ha <sup>-1</sup> )		
	25 DAS	50 DAS	Harvest	25 DAS	50 DAS	Harvest	25 DAS	50 DAS	Harvest
(Maize + blackgram) + N <sub>1</sub>	11.2	45.2	49.8	0.32	3.01	2.19	360	2280	2436
(Maize + blackgram) + N <sub>2</sub>	11.0	45.3	48.1	0.29	2.98	2.78	354	2200	2380
(Maize + blackgram) + N <sub>3</sub>	11.0	45.3	47.1	0.30	2.98	2.11	350	2196	2300
Sole blackgram	12.2	47.4	50.1	0.32	3.21	3.01	375	2460	2502

N<sub>1</sub>, N<sub>2</sub> and N<sub>3</sub> are 100, 75 and 50 % recommended doses of nitrogen to maize, respectively.

**Table 9 : Yield attributes and yield of blackgram as influenced by maize + legume intercropping and nitrogen management**

Treatment	Yield attributes				Yield	
	No. of clusters per plant	No. of pods per cluster	No. of seeds per pod	1000 seed weight (g)	Seed yield (kg ha <sup>-1</sup> )	Haulm yield (kg ha <sup>-1</sup> )
(Maize + blackgram) + N <sub>1</sub>	6.2	12.6	3.6	46.6	378	609
(Maize + blackgram) + N <sub>2</sub>	5.3	12.3	4.3	46.0	362	538
(Maize + blackgram) + N <sub>3</sub>	5.2	12.0	4.1	45.7	358	514
Sole blackgram	6.4	13.3	4.2	48.1	421	713

N<sub>1</sub>, N<sub>2</sub> and N<sub>3</sub> are 100, 75 and 50 % recommended doses of nitrogen to maize, respectively.

**Table 10: Biometric observations of cluster bean as influenced by maize + legume intercropping and nitrogen management**

Treatment	Plant height (cm)			Leaf area index			Dry matter production (kg ha <sup>-1</sup> )		
	25 DAS	50 DAS	Harvest	25 DAS	50 DAS	Harvest	25 DAS	50 DAS	Harvest
(Maize + cluster bean) +N <sub>1</sub>	9.9	54.6	70.3	0.7	2.9	1.7	124.8	1286	4402
(Maize + cluster bean) +N <sub>2</sub>	8.8	51.4	70.2	0.6	2.6	1.6	117.0	1050	4208
(Maize + cluster bean) +N <sub>3</sub>	8.3	43.3	69.1	0.5	2.5	1.4	109.8	1072	3891
Sole cluster bean	10.9	55.6	72.0	0.7	3.1	1.9	128.4	1352	4580

The study revealed that maize + legume intercropping and nitrogen management had a significant impact on yield attributes, yield, and soil fertility. The treatment with 100% recommended dose of nitrogen (N<sub>1</sub>) in maize + cluster bean intercropping system showed improved yield attributes and yield, resulting in 5.0 clusters per plant, 16.8 pods per cluster, 5.9 seeds per pod, 42.9g 1000 seed weight, and 1063 kg/ha seed yield. Additionally, the haulm yield was recorded at 2900 kg/ha. The highest maize grain equivalent yield (5790 kg/ha) and land equivalent ratio (1.938) were recorded in the treatment where maize was intercropped with cluster bean (T<sub>5</sub>). Nitrogen management also had a significant impact on maize grain equivalent yield and land equivalent ratio. The highest nitrogen uptake (105.0 kg/ha) was recorded in the treatment where maize was grown alone (T<sub>1</sub>), while intercropping maize with greengram (T<sub>3</sub>) and blackgram (T<sub>4</sub>) resulted in similar nitrogen uptake (97.1 kg/ha). Nitrogen management significantly influenced the nutrient uptake of greengram, blackgram, and cluster bean, with the highest nitrogen uptake recorded in cluster bean (34.0 kg/ha) when 100% recommended nitrogen (N<sub>1</sub>) was applied. The highest soil available nitrogen (177 kg/ha) was recorded in the treatment where 100% recommended nitrogen (N<sub>1</sub>) was applied to maize, while intercropping maize with greengram (T<sub>3</sub>) also resulted in higher soil available nitrogen (176 kg/ha). The highest computed nitrogen balance (125 kg/ha) was recorded in the treatment where maize was intercropped with greengram (T<sub>3</sub>) and cluster bean (T<sub>5</sub>) with 100% recommended nitrogen (N<sub>1</sub>). The growth parameters, yield attributes, and yield of the intercrops, viz., greengram, blackgram, and cluster bean, were higher in their respective sole crops, followed by application of 100%, 75%, and 50% recommended dose of nitrogen to maize. The maize grain equivalent yield and land equivalent ratio were higher in maize + cluster bean intercropping, followed by maize + greengram and maize + blackgram. The application of 100% recommended dose of nitrogen to maize resulted in significant superiority of maize grain equivalent yield and land equivalent ratio. The nutrient uptake by maize was influenced by maize intercropping and nitrogen management. The highest value of nitrogen and phosphorus uptake by maize was associated with 60 x 20 cm planting, which was comparable with that of maize + greengram and maize + blackgram intercrops. The post-harvest soil status of available nitrogen, phosphorus, and potassium was influenced by maize intercropping and nitrogen management. The post-harvest soil status of available nitrogen was highest in maize + greengram, closely followed by maize + blackgram and maize + cluster bean intercropping.

## Conclusion

The study investigated the effects of maize intercropping and nitrogen management on maize growth, yield, and economic returns. The results showed that maize intercropping and nitrogen management significantly influenced maize growth, yield attributes, and yield. Sole maize at 60 x 20 cm spacing and maize + cluster

bean intercropping resulted in higher growth parameters, yield attributes, and yield. The application of 100% recommended dose of nitrogen to maize resulted in significant higher yield attributes and grain and stover yield. The growth parameters, yield attributes, and yield of the intercrops were higher in their respective sole crops, followed by application of 100%, 75%, and 50% recommended dose of nitrogen to maize. In conclusion, the study suggests that maize intercropping with cluster bean and application of 100% recommended dose of nitrogen to maize can result in higher growth parameters, yield attributes, and yield, as well as higher economic returns. Optimizing nitrogen management in maize-based intercropping systems is crucial to achieve higher yields.

## References

1. Arya., M.P.S and Saini., R.I. 1989. Effect of planting geometry on maize and soybean intercropping system under rainfed condition. *Indian Journal of Agronomy* 34(3):322-324.
2. Asmat Ullah, Ashraf Bhatti M, Zulfiqar Ali Gurmani and Imran., M. 2007. Studies on planting patterns of maize (*Zea mays L.*) facilitating legumes intercropping. *Journal of Agricultural Research* 45(2):113-118.
3. Barik., K.C. 1997. Maize + legume intercropping for north central plateau of Orissa. *Legume Research* 20(3/4):218-220.
4. Bhagat., R.K. 2002. Intercropping of pignonpea and maize under rainfed condition. *Journal of Research (BAU)* 14(2):233-235.
5. Bhatt., B.S. and Damor., U.M. 1985. Relative performance of intercrops at graded levels of fertilizers in maize. *Indian Journal of Agronomy* 30: 514-515.
6. Bhattacharya., A and Goutam., R.C. 1996. Performance of maize legume intercropping at different rates of nitrogen. *Annals of Agricultural Research* 17(25):215-216.
7. Chalka., M.K and Nepalia., V. 2005. Production potential and economics of maize (*Zea mays*) intercropped with legumes as influenced by weed control. *Indian Journal of Agronomy* 50(2):119-122.
8. Chittapur., B.M, Babalad H.B, Hiremath., S.M, Kulkarni., G.A and Sardesh Pande D.R. 1994. Intercropping of fodder legumes in grain maize. *Journal of Maharashtra Agricultural University* 19(3):454-455.
9. Dasaraddi., S.V, Hiremeth., S.M and Patil., R.H. 2002. Performance of legumes as intercrops in maize. *Journal of Research BAU*, 14(2):241-244.
10. Dilip Singh and Nepalia., V. 2009. Influence of integrated nutrient management on quality protein maize (*Zea mays*) productivity and soils of Southern Rajasthan. *Indian Journal of Agricultural Sciences* 79(12):1020-1022.
11. Gangwar., B and Kalra., G.S. 1981. Influence of maize legume associations and nitrogen levels on growth and dry matter accumulation of rainfed maize. *Madras Agricultural Journal* 68(7):450-457.
12. Gaur., B.L, Mansion., P.R and Gupta., D.C. 1992. Effect of nitrogen levels and their split on yield of winter maize (*Zea mays*). *Indian Journal of Agronomy* 37(4):816-

817.

13. Halikatti., S.I and Banarasilal. 1998. Production potentiality of maize as influenced by planting geometry, mulching and grain legume intercropping. Karnataka Journal of Agricultural Sciences 11(4):883-888.
14. Ibrahim Mohammad, Tyagi., R.C. and Rammohan Rao., D.S. 1993. Studies on maize – legumes intercropping systems in relation to nitrogen levels. Farming Systems 9(1&2):56-58.
15. Intodia., S.K. 1999. Economically viable cropping system under rainfed condition in south Rajasthan. Annals of Agriculture Research 20(4):504-505.
16. Kaire Moses., G.B, Ikramullah., M and Shaik Mohammad. 2000. Performance of maize in intercropping with legumes at different levels of fertilizers. Crop Research 20(1):149- 151.
17. Khandkar., U.R and Nigam., K.B. 1996. Effect of maize legume intercropping systems on soil fertility. Indian Journal of Agricultural Sciences 66(1):601-603.
18. Shivay., YS, Singh R.P and Pandey., C.S. 1999. Response of nitrogen in maize (*Zea mays*) based intercropping system. Indian Journal of Agronomy 44(2):261-266.
19. Singh MK., Pasu., S.K, Thakur. R and Herman., U.N. 1995. Effect of fertilizer and plant density management on productivity and economics of maize and blackgram intercropping. Indian Journal of Agricultural Sciences 65(1):800-803.
20. Singh., M.K, Thakur., R, Verma., U.N, Pal., S.K and Pasupalak., S. 1998. Productivity and nutrient balance of maize + blackgram intercropping as affected by fertilizer and plant density management of blackgram. Indian Journal of Agronomy 43:495-500.
21. Singh., R.P and Kaushik., M.K. 1987. Nitrogen economy in maize legume intercropping system. Annals of Agriculture Research 8(1):105-109.

UNDER PEER REVIEW