

## **Improving maize yield and soil productivity through N management practices in maize-legume Intercropping**

### **Abstract**

A field experiment was conducted to investigate the effects of maize intercropping and nitrogen management on crop productivity. The study comprised two factors: maize intercropping with five treatments (sole maize, skipped row maize, and maize intercropped with greengram, blackgram, or cluster bean) and nitrogen management with three treatments (100%, 75%, and 50% of the recommended nitrogen dose). The results showed that sole maize at 60 x 20 cm spacing and maize intercropped with cluster bean, blackgram, or greengram significantly outyielded sole maize in skipped rows. Applying nitrogen at the full recommended dose (100%) significantly enhanced maize yields, while the cluster bean intercropping system excelled in terms of maize grain equivalent yield and land equivalent ratio. The study highlights the potential of maize intercropping and optimized nitrogen management to enhance crop productivity, reduce soil nutrient depletion, and promote sustainable agriculture practices.

**Keywords:** maize intercropping, nitrogen management, crop productivity, sustainable agriculture.

### **Introduction**

“Sustainable crop productivity is crucial in Indian agriculture, necessitating innovative strategies for crop intensification with sustainable nutrition. Maize (*Zea mays* L.) is the world's third most important cereal crop and India's seventh largest producer. Its multidimensional uses, high production potential, and adaptability make it an essential crop for food production. In southern Andhra Pradesh's agro-climatic zone, maize intercropping has gained popularity. However, hybrid maize requires high nutrient levels, particularly nitrogen, with recent hybrids responding to over 240 kg N/ha. The prohibitive cost of fertilizer nitrogen limits its application, and environmental concerns necessitate alternative nitrogen sources. Legumes, when associated with maize, can reduce nitrogen requirements while maintaining soil health. Despite considerable research on maize nutrition, most studies focused on resource exploitative approaches rather than conservation-based methods” [1,2]. “Traditional nitrogen recommendations for maize-based intercropping systems have been determined by individual crop responses, leading to suboptimal nitrogen management and reduced economic returns. A rational agronomic approach, considering uniform plant populations of both base and intercrops, is essential for optimizing maize + legume intercropping systems. The production potential and economic viability of these systems depend on the nature and type of associated legume and fertilizer application. This study aims to explore the possibilities of intercropping short-duration legumes with maize under variable nitrogen doses and intercropping sequences using different crop geometries. By adopting a sustainable and conservation-based approach, this research seeks to optimize maize production, reduce environmental degradation, and promote economic viability for farmers” [3,4].

## Material and Methods:

The experiment was conducted at the Agricultural Research Farm, Sri CB Singh Memorial Shikshan Sansthan, Jhinhak, Kanpur Dehat, U.P., India, during the 2022-23 Kharif season. The research site is located in the Indo-Gangetic Plains, with a geographical position of 26.35° N latitude and 80.09° E longitude, and an elevation of 130.00 m above mean sea level. A randomized block design with a factorial concept was employed, with three replications. The experiment consisted of two factors: maize intercropping systems and nitrogen management. The maize intercropping systems included five treatments: sole maize, skipped row maize, and maize intercropped with greengram, blackgram, or cluster bean. The nitrogen management factor comprised three treatments: 100%, 75%, and 50% of the recommended nitrogen dose (240 kg ha<sup>-1</sup>). Sole crops of greengram, blackgram, and cluster bean were raised separately outside the experimental layout. Each plot measured 6.0 x 5.4 m (gross) and 4.0 x 3.6 m (net). Intercrops were sown at a spacing of 60 x 5 cm. A uniform dose of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 50 kg K<sub>2</sub>O ha<sup>-1</sup> was applied as basal to maize in all plots. Nitrogen was split into three equal applications: basal, knee stage, and tasseling stage. For the intercrops, 20, 50, and 40 kg N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O ha<sup>-1</sup>, respectively, were applied as basal at sowing

## Results and Discussion

The study's results showed that maize intercropping and nitrogen management significantly impacted maize's growth parameters, yield attributes, and yield. Sole maize planted at 60 x 20 cm spacing exhibited superior growth parameters, including plant height, leaf area index, and dry matter production, which were comparable to those of maize intercropped with cluster bean, blackgram, and greengram. In terms of yield attributes, sole maize at 60 x 20 cm spacing displayed higher cob length and girth, number of grains per cob, and 100-grain weight, closely followed by maize intercropped with cluster bean, blackgram, and greengram. The grain and stover yield of maize was highest in sole maize at 60 x 20 cm spacing, which was statistically at par with the yields obtained from maize intercropped with cluster bean, blackgram, and greengram. Nitrogen application at 100% of the recommended dose significantly enhanced yield attributes and grain and stover yield, whereas applying 50% of the recommended nitrogen dose resulted in significantly lower yield attributes and grain and stover yield.

**Table 1: Effect of Maize + Legume Intercropping and Nitrogen Management on Plant Height (cm) of Maize**

| Treatments                                     | 25 DAS | 50 DAS | 75 DAS |
|--|--------|--------|--------|
| <b>Maize intercropping</b>                     |        |        |        |
| T <sub>1</sub> : Maize 60 x 20 cm              | 52.6   | 126.9  | 164.9  |
| T <sub>2</sub> : Maize skipped row             | 48.3   | 109.8  | 142.6  |
| T <sub>3</sub> : T <sub>2</sub> + Greengram    | 48.2   | 121.6  | 158.0  |
| T <sub>4</sub> : T <sub>2</sub> + Blackgram    | 48.6   | 124.8  | 158.4  |
| T <sub>5</sub> : T <sub>2</sub> + Cluster bean | 52.1   | 125.1  | 161.6  |
| SEm ±  | 2.36   | 3.61   | 4.02   |
| CD (P=0.05)                                    | NS     | 10.3   | 11.5   |
| <b>Nitrogen management</b>                     |        |        |        |
| N <sub>1</sub> : 100 % Rec. N to maize         | 52.4   | 124.2  | 164.6  |

|                                       |      |       |       |
|---------------------------------------|------|-------|-------|
| N <sub>2</sub> : 75 % Rec. N to maize | 50.3 | 119.2 | 153.9 |
| N <sub>3</sub> : 50 % Rec. N to maize | 48.0 | 105.3 | 141.6 |
| SEm ±                                 | 1.83 | 4.01  | 4.23  |
| CD (P=0.05)                           | NS   | 11.5  | 12.1  |

**Table 2 : Effect of Maize + Legume Intercropping and Nitrogen Management on Leaf Area Index (LAI) of Maize**

| Treatments                                     | 25 DAS | 50 DAS | 75 DAS | Harvest |
|--|--------|--------|--------|---------|
| <b>Maize intercropping</b>                     |        |        |        |         |
| T <sub>1</sub> : Maize 60 x 20 cm              | 0.212  | 2.50   | 1.49   | 1.28    |
| T <sub>2</sub> : Maize skipped row             | 0.181  | 1.94   | 1.20   | 1.20    |
| T <sub>3</sub> : T <sub>2</sub> + Greengram    | 0.199  | 2.36   | 1.36   | 1.25    |
| T <sub>4</sub> : T <sub>2</sub> + Blackgram    | 0.206  | 2.39   | 1.39   | 1.26    |
| T <sub>5</sub> : T <sub>2</sub> + Cluster bean | 0.212  | 2.41   | 1.42   | 1.28    |
| SEm ±  | 0.07   | 0.14   | 0.048  | 0.014   |
| CD (P=0.05)                                    | NS     | 0.40   | 0.14   | 0.04    |
| <b>Nitrogen management</b>                     |        |        |        |         |
| N <sub>1</sub> : 100 % Rec. N to maize         | 0.217  | 2.38   | 1.36   | 1.11    |
| N <sub>2</sub> : 75 % Rec. N to maize          | 0.201  | 2.36   | 1.29   | 1.09    |
| N <sub>3</sub> : 50 % Rec. N to maize          | 0.179  | 2.01   | 1.14   | 0.99    |
| SEm ±  | 0.006  | 0.108  | 0.037  | 0.010   |
| CD (P=0.05)                                    | 0.017  | 0.31   | 0.11   | 0.03    |

**Table 3 : Effect of Maize + Legume Intercropping and Nitrogen Management on Dry Matter Production (kg ha<sup>-1</sup>) of Maize**

| Treatment | 25 DAS | 50 DAS | 75 DAS | Harvest |
|-----------|--------|--------|--------|---------|
|-----------|--------|--------|--------|---------|

| <b>Maize intercropping</b>                     |       |       |       |       |
|--|-------|-------|-------|-------|
| T <sub>1</sub> : Maize 60 x 20 cm              | 421.7 | 4311  | 7257  | 9174  |
| T <sub>2</sub> : Maize skipped row             | 408.0 | 4004  | 6655  | 8763  |
| T <sub>3</sub> : T <sub>2</sub> + Greengram    | 412.8 | 4296  | 6987  | 8917  |
| T <sub>4</sub> : T <sub>2</sub> + Blackgram    | 417.6 | 4298  | 7056  | 9003  |
| T <sub>5</sub> : T <sub>2</sub> + Cluster bean | 419.6 | 4303  | 7240  | 9103  |
| SEm ±  | 4.8   | 90.9  | 112.0 | 104.2 |
| CD (P=0.05)                                    | NS    | 262   | 323   | 301   |
| <b>Nitrogen management</b>                     |       |       |       |       |
| N <sub>1</sub> : 100 % Rec. N to maize         | 420.1 | 4199  | 6987  | 9314  |
| N <sub>2</sub> : 75 % Rec. N to maize          | 418.8 | 4109  | 6763  | 9074  |
| N <sub>3</sub> : 50 % Rec. N to maize          | 404.2 | 3802  | 6213  | 8517  |
| SEm ±  | 3.61  | 102.5 | 86.7  | 90.2  |
| CD (P=0.05)                                    | 10.6  | 297   | 251   | 260   |

**Table 4 : Yield attributes of maize as influenced by maize + legume intercropping and nitrogen management**

| <b>Treatment</b>                            | <b>Cob length (cm)</b> | <b>Cob girth (cm)</b> | <b>No. of grains per cob</b> | <b>100 grain weight (g)</b> |
|---|------------------------|-----------------------|------------------------------|-----------------------------|
| <b>Maize intercropping</b>                  |                        |                       |                              |                             |
| T <sub>1</sub> : Maize 60 x 20 cm           | 14.4                   | 14.7                  | 333                          | 28.8                        |
| T <sub>2</sub> : Maize skipped row          | 12.1                   | 12.9                  | 252                          | 23.8                        |
| T <sub>3</sub> : T <sub>2</sub> + Greengram | 13.7                   | 13.4                  | 297                          | 26.4                        |
| T <sub>4</sub> : T <sub>2</sub> + Blackgram | 13.9                   | 14.4                  | 299                          | 27.5                        |

|  |      |      |       |      |
|--|------|------|-------|------|
| T <sub>5</sub> : T <sub>2</sub> + Cluster bean | 14.4 | 14.6 | 324   | 27.7 |
| SEm ±  | 0.74 | 0.54 | 26.83 | 1.15 |
| CD (P=0.05)                                    | 2.1  | 1.6  | 77    | 3.3  |
| <b>Nitrogen management</b>                     |      |      |       |      |
| N <sub>1</sub> : 100 % Rec. N to maize         | 14.8 | 14.6 | 333   | 28.1 |
| N <sub>2</sub> : 75 % Rec. N to maize          | 14.5 | 14.3 | 324   | 27.7 |
| N <sub>3</sub> : 50 % Rec. N to maize          | 12.3 | 13.0 | 245   | 24.7 |
| SEm ±  | 0.57 | 0.42 | 20.78 | 0.88 |
| CD (P=0.05)                                    | 1.6  | 1.2  | 60    | 2.5  |

**Table 5 : Effect of Maize + Legume Intercropping and Nitrogen Management on Grain Yield, Stover Yield, and Harvest Index of Maize**

| Treatment                                      | Grain yield (kg ha <sup>-1</sup> ) | Stover yield (kg ha <sup>-1</sup> ) | Harvest index |
|--|------------------------------------|-------------------------------------|---------------|
| <b>Maize intercropping</b>                     |                                    |                                     |               |
| T <sub>1</sub> : Maize 60 x 20 cm              | 4009                               | 5921                                | 40.4          |
| T <sub>2</sub> : Maize skipped row             | 3071                               | 4277                                | 41.8          |
| T <sub>3</sub> : T <sub>2</sub> + Greengram    | 3791                               | 5685                                | 40.0          |
| T <sub>4</sub> : T <sub>2</sub> + Blackgram    | 3864                               | 5771                                | 40.1          |
| T <sub>5</sub> : T <sub>2</sub> + Cluster bean | 3905                               | 5825                                | 40.1          |
| SEm ±  | 86.5                               | 93.7                                | 1.23          |
| CD (P=0.05)                                    | 250                                | 271                                 | NS            |
| <b>Nitrogen management</b>                     |                                    |                                     |               |
| N <sub>1</sub> : 100 % Rec. N to maize         | 3838                               | 5809                                | 39.7          |

|                                       |      |      |      |
|---------------------------------------|------|------|------|
| N <sub>2</sub> : 75 % Rec. N to maize | 3791 | 5694 | 40.0 |
| N <sub>3</sub> : 50 % Rec. N to maize | 3055 | 4984 | 38.0 |
| SEm ±                                 | 79.2 | 83.4 | 1.01 |
| CD (P=0.05)                           | 229  | 241  | NS   |

**Table 6 : Biometric observations of greengram as influenced by maize + legume intercropping and nitrogen management**

| Treatment                            | Plant height (cm) |        |         | Leaf area index |        |         | Dry matter production (kg ha <sup>-1</sup> ) |        |         |
|--------------------------------------|-------------------|--------|---------|-----------------|--------|---------|--|--------|---------|
|                                      | 25 DAS            | 50 DAS | Harvest | 25 DAS          | 50 DAS | Harvest | 25 DAS                                       | 50 DAS | Harvest |
| (Maize + greengram) + N <sub>1</sub> | 10.3              | 59.4   | 59.6    | 0.32            | 2.89   | 2.02    | 110  | 2401   | 2492    |
| (Maize + greengram) + N <sub>2</sub> | 10.4              | 59.1   | 59.1    | 0.30            | 2.80   | 1.98    | 96   | 2398   | 2413    |
| (Maize + greengram) + N <sub>3</sub> | 10.3              | 59.0   | 59.0    | 0.30            | 2.80   | 1.96    | 96   | 2346   | 2409    |
| Sole greengram                       | 10.5              | 60.2   | 59.6    | 0.33            | 3.02   | 2.63    | 114  | 2462   | 2562    |

N<sub>1</sub>, N<sub>2</sub> and N<sub>3</sub> are 100, 75 and 50 % recommended doses of nitrogen to maize respectively.

**Table 7 : Effect of Maize + Legume Intercropping and Nitrogen Management on Yield Attributes and Yield of Greengram**

| Treatment                            | Yield attributes          |                         |                      |                      | Yield                             |                                    |
|--------------------------------------|---------------------------|-------------------------|----------------------|----------------------|-----------------------------------|------------------------------------|
|                                      | No. of clusters per plant | No. of pods per cluster | No. of seeds per pod | 1000 seed weight (g) | Seed yield (kg ha <sup>-1</sup> ) | Haulm yield (kg ha <sup>-1</sup> ) |
| (Maize + greengram) + N <sub>1</sub> | 11.0                      | 5.3                     | 5.5                  | 37.5                 | 418                               | 860                                |
| (Maize + greengram) + N <sub>2</sub> | 10.8                      | 4.9                     | 5.2                  | 37.1                 | 394                               | 854                                |
| (Maize + greengram) + N <sub>3</sub> | 9.6                       | 5.4                     | 4.9                  | 36.1                 | 384                               | 851                                |
| Sole greengram                       | 11.6                      | 5.5                     | 5.4                  | 38.0                 | 425                               | 860                                |

N<sub>1</sub>, N<sub>2</sub> and N<sub>3</sub> are 100, 75 and 50 % recommended doses of nitrogen to maize, respectively

**Table 8 : Effect of Maize + Legume Intercropping and Nitrogen Management on Biometric Observations of Blackgram**

| Treatment                            | Plant height (cm) |        |         | Leaf area index |        |         | Dry matter production (kg ha <sup>-1</sup> ) |        |         |
|--------------------------------------|-------------------|--------|---------|-----------------|--------|---------|--|--------|---------|
|                                      | 25 DAS            | 50 DAS | Harvest | 25 DAS          | 50 DAS | Harvest | 25 DAS                                       | 50 DAS | Harvest |
| (Maize + blackgram) + N <sub>1</sub> | 11.2              | 45.2   | 49.8    | 0.32            | 3.01   | 2.19    | 360  | 2280   | 2436    |
| (Maize + blackgram) + N <sub>2</sub> | 11.0              | 45.3   | 48.1    | 0.29            | 2.98   | 2.78    | 354  | 2200   | 2380    |
| (Maize + blackgram) + N <sub>3</sub> | 11.0              | 45.3   | 47.1    | 0.30            | 2.98   | 2.11    | 350  | 2196   | 2300    |
| Sole blackgram                       | 12.2              | 47.4   | 50.1    | 0.32            | 3.21   | 3.01    | 375  | 2460   | 2502    |

N<sub>1</sub>, N<sub>2</sub> and N<sub>3</sub> are 100, 75 and 50 % recommended doses of nitrogen to maize, respectively.

**Table 9 : Effect of Maize + Legume Intercropping and Nitrogen Management on Yield Attributes and Yield of Blackgram**

| Treatment                            | Yield attributes          |                         |                      |                      | Yield                             |                                    |
|--------------------------------------|---------------------------|-------------------------|----------------------|----------------------|-----------------------------------|------------------------------------|
|                                      | No. of clusters per plant | No. of pods per cluster | No. of seeds per pod | 1000 seed weight (g) | Seed yield (kg ha <sup>-1</sup> ) | Haulm yield (kg ha <sup>-1</sup> ) |
| (Maize + blackgram) + N <sub>1</sub> | 6.2                       | 12.6                    | 3.6                  | 46.6                 | 378                               | 609                                |
| (Maize + blackgram) + N <sub>2</sub> | 5.3                       | 12.3                    | 4.3                  | 46.0                 | 362                               | 538                                |
| (Maize + blackgram) + N <sub>3</sub> | 5.2                       | 12.0                    | 4.1                  | 45.7                 | 358                               | 514                                |
| Sole blackgram                       | 6.4                       | 13.3                    | 4.2                  | 48.1                 | 421                               | 713                                |

N<sub>1</sub>, N<sub>2</sub> and N<sub>3</sub> are 100, 75 and 50 % recommended doses of nitrogen to maize, respectively.

**Table 10: Effect of Maize + Legume Intercropping and Nitrogen Management on Biometric Observations of Cluster Bean**

| Treatment                              | Plant height (cm) |        |         | Leaf area index |        |         | Dry matter production (kg ha <sup>-1</sup> ) |        |         |
|--|-------------------|--------|---------|-----------------|--------|---------|--|--------|---------|
|  | 25 DAS            | 50 DAS | Harvest | 25 DAS          | 50 DAS | Harvest | 25 DAS                                       | 50 DAS | Harvest |
| (Maize + cluster bean) +N <sub>1</sub> | 9.9               | 54.6   | 70.3    | 0.7             | 2.9    | 1.7     | 124.8  | 1286   | 4402    |

|  |      |      |      |     |     |     |       |      |      |
|--|------|------|------|-----|-----|-----|-------|------|------|
| (Maize + cluster bean) +N <sub>2</sub> | 8.8  | 51.4 | 70.2 | 0.6 | 2.6 | 1.6 | 117.0 | 1050 | 4208 |
| (Maize + cluster bean) +N <sub>3</sub> | 8.3  | 43.3 | 69.1 | 0.5 | 2.5 | 1.4 | 109.8 | 1072 | 3891 |
| Sole cluster bean                      | 10.9 | 55.6 | 72.0 | 0.7 | 3.1 | 1.9 | 128.4 | 1352 | 4580 |

The study demonstrated that maize + legume intercropping and nitrogen management had a profound impact on yield attributes, yield, and soil fertility. Notably, the treatment with 100% recommended dose of nitrogen (N1) in maize + cluster bean intercropping system exhibited improved yield attributes and yield, resulting in 5.0 clusters per plant, 16.8 pods per cluster, 5.9 seeds per pod, 42.9g 1000 seed weight, and 1063 kg/ha seed yield. Additionally, the haulm yield was recorded at 2900 kg/ha. The highest maize grain equivalent yield (5790 kg/ha) and land equivalent ratio (1.938) were recorded in the treatment where maize was intercropped with cluster bean. Nitrogen management also played a crucial role in influencing maize grain equivalent yield and land equivalent ratio. The highest nitrogen uptake (105.0 kg/ha) was recorded in the treatment where maize was grown alone, while intercropping maize with greengram and blackgram resulted in similar nitrogen uptake (97.1 kg/ha). Furthermore, nitrogen management significantly influenced the nutrient uptake of greengram, blackgram, and cluster bean, with the highest nitrogen uptake recorded in cluster bean (34.0 kg/ha) when 100% recommended nitrogen (N1) was applied. The study also revealed that the highest soil available nitrogen (177 kg/ha) was recorded in the treatment where 100% recommended nitrogen (N1) was applied to maize, while intercropping maize with greengram also resulted in higher soil available nitrogen (176 kg/ha). The highest computed nitrogen balance (125 kg/ha) was recorded in the treatment where maize was intercropped with greengram and cluster bean with 100% recommended nitrogen (N1). “The growth parameters, yield attributes, and yield of the intercrops, including greengram, blackgram, and cluster bean, were higher in their respective sole crops, followed by application of 100%, 75%, and 50% recommended dose of nitrogen to maize. The maize grain equivalent yield and land equivalent ratio were higher in maize + cluster bean intercropping, followed by maize + greengram and maize + blackgram. The application of 100% recommended dose of nitrogen to maize resulted in significant superiority of maize grain equivalent yield and land equivalent ratio. Moreover, the nutrient uptake by maize was influenced by maize intercropping and nitrogen management. The highest value of nitrogen and phosphorus uptake by maize was associated with 60 x 20 cm planting, which was comparable with that of maize + greengram and maize + blackgram intercrops. The post-harvest soil status of available nitrogen, phosphorus, and potassium was influenced by maize intercropping and nitrogen management. The post-harvest soil status of available nitrogen was highest in maize + greengram, closely followed by maize + blackgram and maize + cluster bean intercropping” [17,18].

## Conclusion

This study examined the effects of maize intercropping and nitrogen management on maize growth, yield, and economic returns. The results revealed that maize intercropping and nitrogen management significantly impacted maize growth, yield attributes, and yield. Sole maize at 60 x 20 cm spacing and maize + cluster bean intercropping exhibited superior growth parameters, yield attributes, and yield. The application of 100% recommended dose of nitrogen to maize resulted in substantially higher yield attributes and grain and stover yield. The growth parameters, yield attributes, and yield of the intercrops were highest in their respective sole crops, followed by the application of 100%, 75%, and 50% recommended dose of nitrogen to maize. In conclusion, this study suggests that intercropping maize with cluster bean and applying 100% recommended dose of nitrogen to maize can lead to improved growth parameters, yield attributes, and

yield, as well as enhanced economic returns. Optimizing nitrogen management in maize-based intercropping systems is crucial for achieving higher yields.

#### Disclaimer (Artificial intelligence)

##### Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

##### Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. There are no use of any ai technologies
- 2.
- 3.

#### References

1. Arya., M.P.S and Saini., R.I. 1989. Effect of planting geometry on maize and soybean intercropping system under rainfed condition. *Indian Journal of Agronomy* 34(3):322-324.
2. Asmat Ullah, Ashraf Bhatti M, Zulfiqar Ali Gurmani and Imran., M. 2007. Studies on planting patterns of maize (*Zea mays L.*) facilitating legumes intercropping. *Journal of Agricultural Research* 45(2):113-118.
3. Barik., K.C. 1997. Maize + legume intercropping for north central plateau of Orissa. *Legume Research* 20(3/4):218-220.
4. Bhagat., R.K. 2002. Intercropping of pignonpea and maize under rainfed condition. *Journal of Research (BAU)* 14(2):233-235.
5. Bhatt., B.S. and Damor., U.M. 1985. Relative performance of intercrops at graded levels of fertilizers in maize. *Indian Journal of Agronomy* 30: 514-515.
6. Bhattacharya., A and Goutam., R.C. 1996. Performance of maize legume intercropping at different rates of nitrogen. *Annals of Agricultural Research* 17(25):215-216.
7. Chalka., M.K and Nepalia., V. 2005. Production potential and economics of maize (*Zea mays*) intercropped with legumes as influenced by weed control. *Indian Journal of Agronomy* 50(2):119-122.
8. Chittapur., B.M, Babalad H.B, Hiremath., S.M, Kulkarni., G.A and Sardesh Pande D.R. 1994. Intercropping of fodder legumes in grain maize. *Journal of Maharashtra Agricultural University* 19(3):454-455.
9. Dasaraddi., S.V, Hiremeth., S.M and Patil., R.H. 2002. Performance of legumes as intercrops in maize. *Journal of Research BAU*, 14(2):241-244.
10. Dilip Singh and Nepalia., V. 2009. Influence of integrated nutrient management

- on quality protein maize (*Zea mays*) productivity and soils of Southern Rajasthan. *Indian Journal of Agricultural Sciences* 79(12):1020-1022.
11. Gangwar., B and Kalra., G.S. 1981. Influence of maize legume associations and nitrogen levels on growth and dry matter accumulation of rainfed maize. *Madras Agricultural Journal* 68(7):450-457.
  12. Gaur., B.L, Mansion., P.R and Gupta., D.C. 1992. Effect of nitrogen levels and their split on yield of winter maize (*Zea mays*). *Indian Journal of Agronomy* 37(4):816-817.
  13. Halikatti., S.I and Banarasilal. 1998. Production potentiality of maize as influenced by planting geometry, mulching and grain legume intercropping. *Karnataka Journal of Agricultural Sciences* 11(4):883-888.
  14. Ibrahim Mohammad, Tyagi., R.C. and Rammohan Rao., D.S. 1993. Studies on maize – legumes intercropping systems in relation to nitrogen levels. *Farming Systems* 9(1&2):56-58.
  15. Intodia., S.K. 1999. Economically viable cropping system under rainfed condition in south Rajasthan. *Annals of Agriculture Research* 20(4):504-505.
  16. Kaire Moses., G.B, Ikramullah., M and Shaik Mohammad. 2000. Performance of maize in intercropping with legumes at different levels of fertilizers. *Crop Research* 20(1):149- 151.
  17. Khandkar., U.R and Nigam., K.B. 1996. Effect of maize legume intercropping systems on soil fertility. *Indian Journal of Agricultural Sciences* 66(1):601-603.
  18. Shivay., YS, Singh R.P and Pandey., C.S. 1999. Response of nitrogen in maize (*Zea mays*) based intercropping system. *Indian Journal of Agronomy* 44(2):261-266.
  19. Singh MK., Pasu., S.K, Thakur. R and Herman., U.N. 1995. Effect of fertilizer and plant density management on productivity and economics of maize and blackgram intercropping. *Indian Journal of Agricultural Sciences* 65(1):800-803.
  20. Singh., M.K, Thakur., R, Verma., U.N, Pal., S.K and Pasupalak., S. 1998. Productivity and nutrient balance of maize + blackgram intercropping as affected by fertilizer and plant density management of blackgram. *Indian Journal of Agronomy* 43:495-500.
  21. Singh., R.P and Kaushik., M.K. 1987. Nitrogen economy in maize legume intercropping system. *Annals of Agricultural Research* 8(1):105-109

UNDER PEER REVIEW