

Linear Estimation in the Type II Generalized Logistic Distribution under Progressive Censoring

ABSTRACT

Generalized distributions have become increasingly popular in applications. They are highly flexible in data analysis, especially with skewed data, which are common in many applications. The Generalized Logistic Distribution (GLD) and its special cases have recently received a lot of interest in the literature. We derived estimators of the unknown parameters of type II Generalized Logistic Distribution (Type II GLD) based on progressively type II censored data. A variety of point estimation methods is employed. We considered the best linear unbiased estimator (BLUE) and the best (affine) linear equivariant estimator (BLEE). In addition, we considered Bayesian estimation. Simulation approaches were used to study the estimators and compare them with the maximum likelihood estimator (MLE) in a range of progressive censoring schemes. The mean squared error (MSE) and bias were employed as comparison criteria. An example based on real data is presented.

Keywords: Point Estimation; Best Linear Unbiased Estimation; Best linear equivariant estimation; Type II Generalized Logistic Distribution, Progressive Censoring

1. INTRODUCTION

Considerable attention has been paid in the literature to inference in parametric distributions based on progressively censored data. Balakrishnan and Sandhu (1995) considered progressive Type II censored sample to find the best linear unbiased estimators to estimate the parameters of the exponential distributions. In addition, they found the maximum likelihood estimators (MLE's) and found that they are equal to the BLUE's of the two-parameter exponential distribution. Also, they drew the attention to the fact that the accuracy of the estimators of the location and scale parameters (BLUE) depends on r , n and m but not the progressive censoring scheme R . The generalized exponential distribution was studied by Kundu and Pradhan (2009). They considered Bayesian inference of the parameters of based on the progressively censored data assuming independent gamma priors for the scale and shape parameters. Bayes estimates are approximated using Lindley's approximation and by the importance sampling and Markov chain Monte Carlo techniques. The authors noted that the Bayes estimates have strong advantages over the MLEs, if suitable prior information is available. The generalized Rayleigh distribution was considered by Kousik Maiti and Kayal (2019) where they considered estimation of parameters and reliability characteristics a under progressive type-II censored sample. The MLEs and Bayes estimates of the parameters were obtained under various loss functions. Salah (2020) considered estimating the unknown parameters of α -power exponential distribution under progressively Type II censored data using the MLEs. He found the approximate best linear unbiased estimators (ABLUE's) as an initial guess of the MLEs. The author discovered that ABLUEs and MLEs are so closely related of the exponential distribution with two parameters. This closeness provides good initial estimates of MLEs. Aly and Bleed (2013) considered Bayesian estimation of the generalized logistic distribution based on progressively censored data under accelerated testing.

In this paper, we shall consider the type II generalized logistic distribution whose probability density function is given by

42

43
$$f(x|\lambda, \mu, \sigma) = \frac{\lambda^\alpha}{\sigma\Gamma(\alpha)} \exp[-\alpha \frac{x-\mu}{\sigma}] \exp[-\lambda \exp \frac{x-\mu}{\sigma}], -\infty < x, \mu < \infty; \sigma, \alpha, \lambda > 0. \quad (1)$$

44 Nassar and Elmasri (2012); Azizpour and Asgharzadeh (2018) and Aljarrah et al. (2020) studied
 45 MLEs for the Generalized Logistic Distribution and other distributions under progressive censoring.
 46 Balakrishnan and Hossain (2007) found that the approximate maximum likelihood estimators
 47 (AMLEs) and the MLEs have similar performance in terms of bias and variance. Moreover, Rimawi
 48 and Baklizi (2021) investigated the type II Generalized Logistic Distribution estimators based on type
 49 II progressive censoring data. They analyzed the MLE and the Lindley's approximation to the Bayes
 50 estimator.

51

52 In this work, we will derive approximate linear estimators of the parameters of the type II generalized
 53 logistic distribution using type II progressively censored data. Progressive censoring is a type of
 54 censoring where we have n units that are placed simultaneously on the life-testing experiment.
 55 Immediately following the first failure, r_1 surviving units are randomly chosen and removed from the
 56 experiment. Immediately after the second failure, r_2 items are withdrawn and so on. The procedure is
 57 continued until all r_m remaining units are removed after the m^{th} failure. Note that the r_i 's are fixed
 58 prior to study. If $r_1 = r_2 = \dots = r_m = 0$, then $n = m$ which corresponds to the complete sample, while
 59 when $r_1 = r_2 = \dots = r_{m-1} = 0$, we have $r_m = n - m$ which corresponds to the conventional Type II
 60 right-censoring scheme.

61

62 **2. APPROXIMATE BEST LINEAR UNBIASED ESTIMATORS**

63 Linear statistics have an easy and accurate structure. Researchers have been interested in using linear
 64 inference for parametric distributions with ordered data in a variety of applications because of their
 65 ease and accuracy. Suppose we have $(X = X_{1:m:n}, \dots, X_{m:m:n})$ be a random vector of progressively
 66 Type-II censored order statistics from a distribution with location parameter μ and scale parameter σ .
 67 Let $Y = (Y_{1:m:n}, \dots, Y_{m:m:n})$ be such that:

68
$$Y_{j:m:n} = \frac{X_{j:m:n} - \mu}{\sigma}, j = 1, \dots, m. \quad (2)$$

69 Let $W = \sigma(Y - E(Y))$, $b = E(Y)$, $\theta = (\mu, \sigma)$ and $B = [\mathbb{1}, b]$. It follows that X can be presented as a
 70 linear equation:

71
$$X = \mu \cdot \mathbb{1} + \sigma \cdot Y = \mu \cdot \mathbb{1} + \sigma \cdot E(Y) + W = [\mathbb{1}, b] \begin{pmatrix} \mu \\ \sigma \end{pmatrix} + W = B \theta + W. \quad (3)$$

72 Let Σ be the covariance matrix $cov(Y)$, assuming Σ is regular, and non-singular covariance matrix,
 73 then

74
$$\Sigma = \Delta \Sigma_{Y^R} \Delta. \quad (4)$$

75

76 The best linear unbiased estimator (BLUE) for the parameters under study depends on the evaluation
 77 of the variance covariance matrix of the order statistics from the progressively censored data. This
 78 matrix is very complicated and can not be obtained in closed form. An approximate best linear
 79 unbiased estimator is available. It is derived in Balakrishnan and Cramer (2014). We will apply this
 80 approximation to the location and scale parameters of our model as follows:

81 Suppose we have $m \geq 2$ and $n = \sum_{j=1}^m r_j + 1$, the BLUE estimators of μ and σ are given by

82
$$\hat{\mu}_{LU} = \frac{1}{\Delta} \cdot ((b' \Sigma^{-1} b)(\mathbb{1}' \Sigma^{-1} X) - (\mathbb{1}' \Sigma^{-1} b)(b' \Sigma^{-1} X)), \quad (5)$$

83
$$\hat{\sigma}_{LU} = \frac{1}{\Delta} \cdot ((\mathbb{1}' \Sigma^{-1} \mathbb{1})(b' \Sigma^{-1} X) - (\mathbb{1}' \Sigma^{-1} b)(\mathbb{1}' \Sigma^{-1} X)), \quad (6)$$

84 where $\Delta = ((\mathbb{1}' \Sigma^{-1} \mathbb{1})(b' \Sigma^{-1} b) - (\mathbb{1}' \Sigma^{-1} b)^2) > 0$.

85 In order to find the approximate covariance matrix, we calculate the following quantities;

86
$$\gamma_j = n - j + 1, j = 1, \dots, n, \quad c_r = \prod_{j=1}^r \gamma_j, r = 1, \dots, m, \quad d_r = \prod_{j=1}^r (\gamma_j + 1), r = 1, \dots, m,$$

87

88
$$e_r = \prod_{j=1}^r (\gamma_j + 2), r = 1, \dots, m, \quad a_r = \frac{d_r}{e_r}, r = 1, \dots, m, \quad b_r = \frac{c_r}{d_r}, r = 1, \dots, m,$$

89
$$EU_r = \Pi_r = 1 - b_r, r = 1, \dots, m, \quad COVU_r U_s = (a_r - b_r) b_s, r = 1, \dots, m, s = 1, \dots, m.$$

90 The last quantity $COVU_r U_s$ gives the approximate covariance matrix Σ_{U^R} . Now Calculate the diagonal
 91 matrix Δ with diagonal elements $\left(\frac{1}{f(F^{-1}(\Pi_1))}, \dots, \frac{1}{f(F^{-1}(\Pi_r))}\right)$ where

92 $f(x) = \frac{e^{-\alpha\left(\frac{x_i-\mu}{\sigma}\right)}}{\left(1+e^{-\left(\frac{x_i-\mu}{\sigma}\right)}\right)^{\alpha+1}}$ and $F(x) = 1 - \left[\frac{e^{-\left(\frac{x_i-\mu}{\sigma}\right)}}{1+e^{-\left(\frac{x_i-\mu}{\sigma}\right)}}\right]^\alpha$. We obtain the required covariance matrix,

93 $\Sigma = \Delta \Sigma_{U^R} \Delta$.

94 The best linear equivariant estimators (BLEE) are approximated in a similar manner. Using the same
 95 notation used for the BLUEs, and let $\Delta_1 = \Delta + (\mathbb{1}' \Sigma^{-1} \mathbb{1})$ we obtain

96
$$\hat{\mu}_{LE} = \frac{1}{\Delta_1} \cdot ((b' \Sigma^{-1} b + 1)(\mathbb{1}' \Sigma^{-1} X) - (\mathbb{1}' \Sigma^{-1} b)(b' \Sigma^{-1} X)), \quad (7)$$

97
$$\hat{\sigma}_{LE} = \frac{1}{\Delta_1} \cdot ((\mathbb{1}' \Sigma^{-1} \mathbb{1})(b' \Sigma^{-1} X) - (\mathbb{1}' \Sigma^{-1} b)(\mathbb{1}' \Sigma^{-1} X)). \quad (8)$$

98

99

100 3. BAYESIAN ESTIMATION OF LOCATION AND SCALE PARAMETERS

101 Bayesian statistical methods begin with established 'prior' beliefs and update them with data to
 102 generate 'posterior' beliefs that can be used to make inferences. Based on this technique, we will
 103 derive Bayes estimators for the parameters of the type II generalized logistic distribution (GLD)
 104 location and scale parameters (μ and σ) with type II progressively censored data.

105 To facilitate comparison with the classical estimators, we will assume non-informative prior
 106 distributions for the location and scale parameters, that is, $\pi(\mu) = 1$ and $\pi(\sigma) = 1/\sigma$. The likelihood
 107 function is given by

108
$$l(data|\mu, \sigma) \propto \frac{1}{\sigma^m} \prod_{i=1}^m f(z_{i:m:n}) [1 - F(z_{i:m:n})]^{r_i}. \quad (9)$$

109 Therefore, the joint posterior density of, μ and σ given the data, is given by

110
$$\pi(\mu, \sigma|data) \propto \frac{1}{\sigma} l(data|\mu, \sigma), -\infty < \mu < \infty, \sigma > 0. \quad (10)$$

111 The Bayes estimator of a function of the parameters, say $t = t(\mu, \sigma)$ under the squared error loss
 112 function is given by its posterior expectation

113
$$\hat{t} = \int_0^\infty \int_{-\infty}^\infty t(\mu, \sigma) \pi(\mu, \sigma|data) d\mu d\sigma. \quad (11)$$

114 This integral is difficult to obtain analytically and therefore we can approximate it using either
 115 importance sampling procedures or the Lindley approximation.

116 Importance Sampling can be explained as a weighted average of random samples taken from another
 117 distribution $h_p(x)$ "importance sampling" density function to estimate an expectation with respect to
 118 the target density function $f_x(x)$. The prior distribution of μ and σ are non-informative priors for the
 119 location and scale parameters (μ and σ)

120
$$\pi_1(\mu) = 1, -\infty < \mu < \infty, \quad (12)$$

121
$$\pi_2(\sigma) = \frac{1}{\sigma}, \sigma > 0. \quad (13)$$

122 The joint prior distribution is

123
$$\pi(\mu, \sigma) = \frac{1}{\sigma}, -\infty < \mu < \infty, \sigma > 0. \quad (14)$$

124

125 It follows that the posterior distribution is given by

126
$$\pi(\mu, \sigma|data) = k \frac{\alpha^m}{\sigma^{m+1}} \prod_{i=1}^m \left\{ \frac{1}{\left(1+e^{-\left(\frac{x_i-\mu}{\sigma}\right)}\right)} \left(\frac{e^{-\left(\frac{x_i-\mu}{\sigma}\right)}}{1+e^{-\left(\frac{x_i-\mu}{\sigma}\right)}} \right)^{\alpha(R_i+1)} \right\}$$

127
$$\propto \left\{ \frac{e^{m/\sigma}}{m^{m-1}} \left(1 + e^{-\left(\frac{\mu-\bar{x}}{\sigma/m}\right)}\right)^2 \prod_{i=1}^m \left\{ \frac{e^{-\alpha(R_i+1)-1}\left(\frac{x_i-\mu}{\sigma}\right)}{\left(1+e^{-\left(\frac{x_i-\mu}{\sigma}\right)}\right)^{\alpha(R_i+1)+1}} \right\} \right\}. \quad (15)$$

128

129 We can rewrite the posterior function as:

130
$$\pi(\mu, \sigma|data) \propto f_1(\mu) f_2(\sigma) h(\mu, \sigma), \quad (16)$$

131 where $f_1(\mu) = \left\{ \frac{m}{\sigma} \frac{e^{-\frac{\mu-\bar{x}}{\sigma/m}}}{\left(1+e^{-\frac{\mu-\bar{x}}{\sigma/m}}\right)^2} \right\}$, this is the logistic distribution with parameters $\bar{x} = \frac{\sum_{i=1}^m x_i}{m}$ and σ/m .

132 $f_2(\sigma) = \left\{ \frac{m^{m-1}}{\Gamma(m-1)} \left(\frac{1}{\sigma}\right)^m e^{-m/\sigma} \right\}$, which is the inverse gamma distribution's pdf with parameters $m - 1$
 133 and m , and

$$134 \quad h(\mu, \sigma) = \left\{ \frac{e^{m/\sigma}}{m^{m-1}} \left(1 + e^{-\frac{\mu-\bar{x}}{\sigma/m}}\right)^2 \prod_{i=1}^m \left\{ \frac{e^{-(\alpha(R_i+1)-1)\left(\frac{x_i-\mu}{\sigma}\right)}}{\left(1+e^{-\frac{x_i-\mu}{\sigma}}\right)^{\alpha(R_i+1)+1}} \right\} \right\}. \quad (17)$$

135 To find the estimate of the GLD parameters we do the following steps:

136 *Algorithm 1:*

137 Step 1: Generate σ from inverse gamma distribution with parameters $m - 1$ and m .

138 Step 2: Generate μ from the logistic distribution with parameters $\bar{x} = \frac{\sum_{i=1}^m x_i}{m}$ and σ/m , where σ is
 139 generated from Step 1.

140 Step 3: Repeat steps 1 and 2 to obtain $((\mu_1, \sigma_1), (\mu_2, \sigma_2), \dots, (\mu_N, \sigma_N))$.

141 Step 4: Calculate the Bayes estimate as $\sum_{i=1}^N t(\mu_i, \sigma_i) h((\mu_i, \sigma_i)) / \sum_{i=1}^N h((\mu_i, \sigma_i))$.

142

143

144

4. SIMULATION STUDY

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

A Monte Carlo simulation study is conducted to investigate and compare the performance of the estimators under various experimental situations. We considered various progressive censoring schemes as explained in tables 1 – 6 below, corresponding to sample sizes of 50, 70 and 90. The location and scale parameters were set to zero and one respectively. The parameter α is taken to be 0.5, 1 and 1.5 to cover the various shapes of the distribution. We used the algorithm proposed by Balakrishnan and Sandhu (1996) to generate progressive Type II censored samples from Type II GLD. The findings are presented in Tables 1 and 6. We used 5000 replications in all our simulation runs.

The results include the biases and mean squared errors for the estimators developed in this paper in addition to the Lindley's approximation of the Bayes estimators and the maximum likelihood estimators developed and studied in Balakrishnan and Hossain (2007) and Rimawi and Baklizi (2021).

Table 1. Results of Simulation for parameter μ with GLD ($\alpha = 1.5, \mu = 0, \sigma = 1$)

N	m	Scheme	MLE	Lindley	I.S	BLUE	BLEE
50	30	(0*29,20)					
		Bias	-0.0316	-0.0411	-1.7436	0.0295	0.0101
		MSE	0.0010	0.0017	3.0400	0.0660	0.0648
	30	(0*10,2*10,0*10)					
		Bias	-0.0293	-0.0466	-1.3551	2.2187	2.1775
		MSE	0.0009	0.0022	1.8362	4.9878	0.0648
30	(20,0*29)						
	Bias	-0.0092	-0.0929	-0.8390	2.6077	2.5681	
	MSE	0.0001	0.0086	0.7040	6.8653	0.0648	
50	40	(0*39,10)					
		Bias	-0.0160	-0.0226	-1.2661	0.0172	0.0094

		MSE	0.0003	0.0005	1.6030	0.0497	0.0493
	40	(0*15,1*10,0*15)					
		Bias	-0.0137	-0.0421	-1.0062	0.9233	0.9108
		MSE	0.0002	0.0018	1.0125	0.9019	0.0493
	40	(10,0*39)					
		Bias	-0.0067	-0.0586	-0.7654	1.1288	1.1166
		MSE	0.0000	0.0034	0.5858	1.3237	0.0493
70	40	(0*39,30)					
		Bias	-0.0246	-0.0294	-1.7559	0.0285	0.0129
		MSE	0.0006	0.0009	3.0832	0.0506	0.0495
	40	(0*10,2*15,0*15)					
		Bias	-0.0246	-0.0366	-1.2942	2.6859	2.6498
		MSE	0.0006	0.0013	1.6750	7.2640	0.0495
70	50	(0*49,20)					
		Bias	-0.0147	-0.0224	-1.4289	0.0164	0.0085
		MSE	0.0002	0.0005	2.0419	0.0389	0.0385
	50	(0*20,2*10,0*20)					
		Bias	-0.0166	-0.0557	-1.0992	1.5217	1.5064
		MSE	0.0003	0.0031	1.2083	2.3542	0.0385
	50	(20,0*49)					
		Bias	-0.0101	-0.0557	-0.7403	1.8189	1.8040
		MSE	0.0001	0.0031	0.5481	3.3470	0.0385
90	50	(0*49,40)					
		Bias	-0.0248	-0.0259	-1.7668	0.0183	0.0053
		MSE	0.0006	0.0007	3.1217	0.0406	0.0401
	50	(0*15,2*20,0*15)					
		Bias	-0.0153	-0.0312	-1.3673	2.8937	2.8620
		MSE	0.0002	0.0010	1.8696	8.4135	0.0401
90	60	(0*59,30)					
		Bias	-0.0076	-0.0180	-1.5100	0.0143	0.0067
		MSE	0.0001	0.0003	2.2800	0.0323	0.0321
	60	(0*20,2*15,0*25)					
		Bias	-0.0067	-0.0252	-1.1241	2.0089	1.9925
		MSE	0.0000	0.0006	1.2636	4.0679	0.0321
	60	(30,0*59)					
		Bias	-0.0029	-0.0420	-0.7201	2.2792	2.2635
		MSE	0.0000	0.0018	0.5185	5.2268	0.0321

162
163
164
165
166

Table 2. Results of Simulation for parameter μ with GLD ($\alpha = 1.0, \mu = 0, \sigma = 1$)

N	m	Scheme	MLE	Lindley	LS	BLUE	BLEE
50	30	(0*29,20)					
		Bias	-0.0145	-0.0260	-1.2894	0.0078	-0.0010
		MSE	0.0002	0.0007	1.6625	0.0649	0.0648
	30	(0*10,2*10,0*10)					
		Bias	-0.0223	-0.0400	-0.8053	1.8900	1.8698
		MSE	0.0005	0.0016	0.6485	3.6369	0.0648
	30	(20,0*29)					
		Bias	-0.0030	-0.0845	-0.2378	2.4078	2.3881
		MSE	0.0000	0.0071	0.0565	5.8622	0.0648
50	40	(0*39,10)					
		Bias	-0.0044	-0.0148	-0.7395	-0.0040	-0.0056
		MSE	0.0000	0.0002	0.5468	0.0584	0.0584

	40	(0*15,1*10,0*15)					
		Bias	-0.0108	-0.0322	-0.4200	0.6519	0.6492
		MSE	0.0001	0.0010	0.1764	0.4834	0.0584
	40	(10,0*39)					
		Bias	0.0044	-0.0779	-0.1488	0.9265	0.9239
		MSE	0.0000	0.0061	0.0221	0.9169	0.0584
70	40	(0*39,30)					
		Bias	-0.0140	-0.0206	-1.3127	0.0046	-0.0028
		MSE	0.0002	0.0004	1.7231	0.0482	0.0482
	40	(0*10,2*15,0*15)					
		Bias	-0.0094	-0.0276	-0.7473	2.3503	2.3314
		MSE	0.0001	0.0008	0.5585	5.5720	0.0482
	40	(30,0*39)					
		Bias	-0.0027	-0.0730	-0.1854	2.8241	2.8059
		MSE	0.0000	0.0053	0.0344	8.0237	0.0482
70	50	(0*49,20)					
		Bias	-0.0020	-0.0148	-0.9359	-0.0020	-0.0045
		MSE	0.0000	0.0002	0.8759	0.0432	0.0432
	50	(0*20,2*10,0*20)					
		Bias	-0.0093	-0.0213	-0.5268	1.1800	1.1749
		MSE	0.0001	0.0005	0.2775	1.4356	0.0432
	50	(20,0*49)					
		Bias	-0.0081	-0.0561	-0.1273	1.5672	1.5622
		MSE	0.0001	0.0032	0.0162	2.4993	0.0432
90	50	(0*49,40)					
		Bias	-0.0120	-0.0179	-1.3227	0.0062	-0.0002
		MSE	0.0001	0.0003	1.7496	0.0385	0.0384
	50	(0*15,2*20,0*15)					
		Bias	-0.0150	-0.0156	-0.8236	2.5062	2.4892
		MSE	0.0002	0.0002	0.6784	6.3193	0.0384
90	60	(0*59,30)					
		Bias	-0.0057	-0.0175	-1.0327	0.0018	-0.0010
		MSE	0.0000	0.0003	1.0664	0.0346	0.0346
	60	(0*20,2*15,0*25)					
		Bias	-0.0045	-0.0221	-0.5478	1.6323	1.6258
		MSE	0.0000	0.0005	0.3001	2.6990	0.0346
	60	(30,0*59)					
		Bias	0.0012	-0.0510	-0.1158	2.0324	2.0260
		MSE	0.0000	0.0026	0.0134	4.1650	0.0346

167
168
169
170
171

Table 3. Results of Simulation for parameter μ with GLD ($\alpha = 0.5, \mu = 0, \sigma = 1$)

N	m	Scheme	MLE	Bayesian	Importance	BLUE	BLEE
50	30	(0*29,20)					
		Bias	0.0155	-0.0507	-0.3528	-0.0283	-0.0219
		MSE	0.0002	0.0026	0.1245	0.0997	0.0989
	30	(0*10,2*10,0*10)					
		Bias	-0.0015	-0.0836	0.3704	0.8626	0.8792
		MSE	0.0000	0.0070	0.1372	0.8430	0.0989
	30	(20,0*29)					
		Bias	0.0007	-0.2832	1.1404	1.6587	1.6758
		MSE	0.0000	0.0802	1.3005	2.8502	0.0989
50	40	(0*39,10)					
		Bias	0.0140	-0.0257	0.3215	-0.0389	-0.0319
		MSE	0.0002	0.0007	0.1033	0.1003	0.0987

40	(0*15,1*10,0*15)						
	Bias	0.0081	-0.1002	0.8464	0.0444	0.0564	
	MSE	0.0001	0.0100	0.7164	0.1007	0.0987	
40	(10,0*39)						
	Bias	0.0062	-0.2277	1.2132	0.4070	0.4193	
	MSE	0.0000	0.0519	1.4719	0.2644	0.0987	
70	40	(0*39,30)					
	Bias	0.0072	-0.0312	-0.4076	-0.0225	-0.0183	
	MSE	0.0001	0.0010	0.1661	0.0720	0.0715	
40	(0*10,2*15,0*15)						
	Bias	-0.0026	-0.0649	0.4517	1.2506	1.2631	
	MSE	0.0000	0.0042	0.2040	1.6354	0.0715	
40	(30,0*39)						
	Bias	0.0013	-0.2201	1.1894	2.0300	2.0426	
	MSE	0.0000	0.0484	1.4147	4.1924	0.0715	
70	50	(0*49,20)					
	Bias	0.0022	-0.0221	0.0621	-0.0313	-0.0263	
	MSE	0.0000	0.0005	0.0039	0.0723	0.0713	
50	(0*20,2*10,0*20)						
	Bias	0.0092	-0.0650	0.7188	0.3066	0.3177	
	MSE	0.0001	0.0042	0.5167	0.1653	0.0713	
50	(20,0*49)						
	Bias	0.0082	-0.1819	1.2491	0.8419	0.8532	
	MSE	0.0001	0.0331	1.5603	0.7801	0.0713	
90	50	(0*49,40)					
	Bias	0.0094	-0.0294	-0.4368	-0.0169	-0.0138	
	MSE	0.0001	0.0009	0.1908	0.0563	0.0560	
50	(0*15,2*20,0*15)						
	Bias	0.0023	-0.0443	0.3366	1.3371	1.3468	
	MSE	0.0000	0.0020	0.1133	1.8439	0.0560	
50	(40,0*49)						
	Bias	0.0066	-0.1864	1.2254	2.2811	2.2910	
	MSE	0.0000	0.0348	1.5017	5.2593	0.0560	
90	60	(0*59,30)					
	Bias	0.0086	-0.0152	-0.0725	-0.0217	-0.0178	
	MSE	0.0001	0.0002	0.0053	0.0563	0.0558	
60	(0*20,2*15,0*25)						
	Bias	0.0041	-0.0531	0.6870	0.5890	0.5989	
	MSE	0.0000	0.0028	0.4719	0.4027	0.0558	
60	(30,0*59)						
	Bias	0.0071	-0.1501	1.2685	1.1942	1.2042	
	MSE	0.0001	0.0225	1.6090	1.4820	0.0558	

172
173

Table 4. Results of Simulation for parameter σ with GLD ($\alpha=1.5, \mu=0, \sigma=1$)

N	m	Scheme	MLE	Bayesian	Importance	BLUE	BLEE
50	30	(0*29,20)					
		Bias	-0.0289	-0.0009	0.3606	0.0558	0.0291
		MSE	0.0008	0.0000	0.1300	0.0290	0.0253
	30	(0*10,2*10,0*10)					
		Bias	-0.0211	-0.0069	0.0971	1.2428	1.1861
		MSE	0.0004	0.0000	0.0094	1.5704	0.0253
	30	(20,0*29)					
		Bias	-0.0154	0.0060	0.0508	1.1522	1.0979
		MSE	0.0002	0.0000	0.0026	1.3535	0.0253
50	40	(0*39,10)					
		Bias	-0.0190	0.0063	0.1550	0.0460	0.0278
		MSE	0.0004	0.0000	0.0240	0.0198	0.0174

40	(0*15,1*10,0*15)						
	Bias	-0.0152	0.0001	0.0689	0.6908	0.6614	
	MSE	0.0002	0.0000	0.0047	0.4949	0.0174	
40	(10,0*39)						
	Bias	-0.0134	0.0010	0.0526	0.6559	0.6272	
	MSE	0.0002	0.0000	0.0028	0.4479	0.0174	
70	40	(0*39,30)					
	Bias	-0.0189	-0.0043	0.3667	0.0448	0.0247	
	MSE	0.0004	0.0000	0.1345	0.0216	0.0192	
40	(0*10,2*15,0*15)						
	Bias	-0.0154	-0.0017	0.0614	1.4195	1.3730	
	MSE	0.0002	0.0000	0.0038	2.0347	0.0192	
70	50	(0*49,20)					
	Bias	-0.0153	0.0000	0.2044	0.0359	0.0210	
	MSE	0.0002	0.0000	0.0418	0.0159	0.0144	
50	(0*20,2*10,0*20)						
	Bias	-0.0126	0.0015	0.0639	0.9904	0.9617	
	MSE	0.0002	0.0000	0.0041	0.9955	0.0144	
50	(20,0*49)						
	Bias	-0.0100	0.0015	0.0413	0.9326	0.9047	
	MSE	0.0001	0.0000	0.0017	0.8843	0.0144	
90	50	(0*49,40)					
	Bias	-0.0178	-0.0025	0.3658	0.0389	0.0228	
	MSE	0.0003	0.0000	0.1338	0.0173	0.0228	
50	(0*15,2*20,0*15)						
	Bias	-0.0108	-0.0062	0.0843	1.5284	1.4892	
	MSE	0.0001	0.0000	0.0071	2.3518	0.0155	
90	60	(0*59,30)					
	Bias	-0.0115	-0.0008	0.2394	0.0315	0.0188	
	MSE	0.0001	0.0000	0.0573	0.0134	0.0123	
60	(0*20,2*15,0*25)						
	Bias	-0.0092	-0.0006	0.0529	1.2133	1.1860	
	MSE	0.0001	0.0000	0.0028	1.4845	0.0123	
60	(30,0*59)						
	Bias	-0.0121	0.0044	0.0405	1.1111	1.0851	
	MSE	0.0001	0.0000	0.0016	1.2469	0.0123	

174
175
176
177
178
179
180
181
182

Table 5. Results of Simulation for parameter σ with GLD ($\alpha=1.0, \mu=0, \sigma=1$)

N	m	Scheme	MLE	Lindley	I.S	BLUE	BLEE
50	30	(0*29,20)					
		Bias	-0.0256	0.0105	0.1913	0.0559	0.0298
		MSE	0.0007	0.0001	0.0366	0.0285	0.0247
	30	(0*10,2*10,0*10)					
		Bias	-0.0189	-0.0015	0.0560	1.4334	1.3733
		MSE	0.0004	0.0000	0.0031	2.0801	0.0247
	30	(20,0*29)					
		Bias	-0.0144	-0.0049	0.0532	1.3737	1.3151
		MSE	0.0002	0.0000	0.0028	1.9125	0.0247
50	40	(0*39,10)					
		Bias	-0.0150	0.0064	0.0746	0.0473	0.0292
		MSE	0.0002	0.0000	0.0056	0.0199	0.0173

40	(0*15,1*10,0*15)						
	Bias	-0.0160	0.0019	0.0416	0.7485	0.7182	
	MSE	0.0003	0.0000	0.0017	0.5779	0.0173	
40	(10,0*39)						
	Bias	-0.0103	-0.0013	0.0398	0.7399	0.7098	
	MSE	0.0001	0.0000	0.0016	0.5651	0.0173	
70	40	(0*39,30)					
	Bias	-0.0173	0.0067	0.1925	0.0424	0.0228	
	MSE	0.0003	0.0000	0.0371	0.0209	0.0188	
40	(0*10,2*15,0*15)						
	Bias	-0.0161	-0.0015	0.0332	1.6443	1.5946	
	MSE	0.0003	0.0000	0.0011	2.7228	0.0188	
40	(30,0*39)						
	Bias	-0.0091	-0.0003	0.0343	1.5484	1.5005	
	MSE	0.0001	0.0000	0.0012	2.4167	0.0188	
70	50	(0*49,20)					
	Bias	-0.0130	0.0095	0.0982	0.0349	0.0202	
	MSE	0.0002	0.0001	0.0096	0.0157	0.0142	
50	(0*20,2*10,0*20)						
	Bias	-0.0115	0.0011	0.0292	1.1164	1.0863	
	MSE	0.0001	0.0000	0.0009	1.2608	0.0142	
50	(20,0*49)						
	Bias	-0.0088	-0.0008	0.0325	1.0805	1.0509	
	MSE	0.0001	0.0000	0.0011	1.1820	0.0142	
90	50	(0*49,40)					
	Bias	-0.0149	0.0006	0.1943	0.0357	0.0200	
	MSE	0.0002	0.0000	0.0378	0.0167	0.0152	
50	(0*15,2*20,0*15)						
	Bias	-0.0129	0.0017	0.0374	1.7541	1.7123	
	MSE	0.0002	0.0000	0.0014	3.0922	0.0152	
90	60	(0*59,30)					
	Bias	-0.0126	0.0030	0.1154	0.0308	0.0183	
	MSE	0.0002	0.0000	0.0133	0.0132	0.0121	
60	(0*20,2*15,0*25)						
	Bias	-0.0100	-0.0007	0.0269	1.3707	1.3420	
	MSE	0.0001	0.0000	0.0007	1.8909	0.0121	
60	(30,0*59)						
	Bias	-0.0081	0.0004	0.0262	1.3090	1.2812	
	MSE	0.0001	0.0000	0.0007	1.7258	0.0121	

183
184
185
186
187

Table 6. Results of Simulation for parameter σ with GLD ($\alpha = 0.5, \mu = 0, \sigma = 1$)

N	M	Scheme	MLE	Bayesian	Importance	BLUE	BLEE
50	30	(0*29,20)					
		Bias	-0.0206	0.0537	0.0684	0.0528	1.0274
		MSE	0.0004	0.0029	0.0047	0.0275	0.0241
	30	(0*10,2*10,0*10)					
		Bias	-0.0170	-0.0005	0.0779	1.7266	1.6609
		MSE	0.0003	0.0000	0.0061	3.0060	0.0241
	30	(20,0*29)					
		Bias	-0.0151	-0.0060	0.1052	1.8265	1.7584
		MSE	0.0002	0.0000	0.0111	3.3607	0.0241
50	40	(0*39,10)					
		Bias	-0.0124	0.0022	0.0422	0.0506	-0.0319
		MSE	0.0002	0.0000	0.0018	0.0208	0.0179

	40	(0*15,1*10,0*15)					
		Bias	-0.0169	0.0018	0.0696	0.8021	0.7697
		MSE	0.0003	0.0000	0.0048	0.6616	0.0179
	40	(10,0*39)					
		Bias	-0.0132	-0.0071	0.0963	0.8504	0.8172
		MSE	0.0002	0.0001	0.0093	0.7414	0.0179
70	40	(0*39,30)					
		Bias	-0.0189	0.0416	0.0590	0.0466	0.0275
		MSE	0.0004	0.0017	0.0035	0.0207	0.0182
	40	(0*10,2*15,0*15)					
		Bias	-0.0140	-0.0017	0.0670	2.0821	2.0260
		MSE	0.0002	0.0000	0.0045	4.3539	0.0182
	40	(30,0*39)					
		Bias	-0.0121	-0.0085	0.0948	2.1116	2.0549
		MSE	0.0001	0.0001	0.0090	4.4772	0.0182
70	50	(0*49,20)					
		Bias	-0.0093	0.0114	0.0332	0.0383	0.0234
		MSE	0.0001	0.0001	0.0011	0.0160	0.0143
	50	(0*20,2*10,0*20)					
		Bias	-0.0113	0.0030	0.0657	1.2792	1.2465
		MSE	0.0001	0.0000	0.0043	1.6509	0.0143
	50	(20,0*49)					
		Bias	-0.0106	-0.0089	0.0832	1.3285	1.2951
		MSE	0.0001	0.0001	0.0069	1.7796	0.0143
90	50	(0*49,40)					
		Bias	-0.0146	0.0334	0.0548	0.0354	0.0202
		MSE	0.0002	0.0011	0.0030	0.0161	0.0147
	50	(0*15,2*20,0*15)					
		Bias	-0.0134	-0.0001	0.0459	2.2139	2.1669
		MSE	0.0002	0.0000	0.0021	4.9164	0.0147
	50	(40,0*49)					
		Bias	-0.0081	-0.0030	0.0860	2.3172	2.2686
		MSE	0.0001	0.0000	0.0074	5.3844	0.0147
90	60	(0*59,30)					
		Bias	-0.0121	0.0179	0.0277	0.0312	0.0188
		MSE	0.0001	0.0003	0.0008	0.0131	0.0120
	60	(0*20,2*15,0*25)					
		Bias	-0.0076	-0.0008	0.0602	1.6402	1.6085
		MSE	0.0001	0.0000	0.0036	2.7023	0.0120
	60	(30,0*59)					
		Bias	-0.0088	-0.0036	0.0773	1.6694	1.6373
		MSE	0.0001	0.0000	0.0060	2.7990	0.0120

188
189
190
191
192
193
194
195
196
197
198
199

5. REAL DATA EXAMPLE: BREAKDOWN OF AN INSULATING FLUID

To evaluate and analyze the quality of transformers and their insulating fluids, a variety of tests have been devised. To explain this, for example, let's consider the Dielectric Breakdown Test, which assesses an insulating liquid's capacity to endure electrical stress up to the point of failure. It displays the voltage at which there will be a breakdown. Moisture, dirt, and conductive particle contamination will induce failure at levels below what is considered tolerable. Nelson (1982) provided a data for the breakdown of an insulating fluid testing experiment. This data collection was examined and evaluated by Balakrishnan and Hossain (2007) examining Type II generalized logistic distribution inference under progressive Type II censoring. Balakrishnan and Hossain (date ?) evaluated and examined the data set that fits the Type II Generalized Logistic Distribution and finding out that MLE and Approximate MLE are very close in the inferencing. In this example $n=19$ and $m=8$ with $\alpha=1$. The data and the results are shown in Tables 7 and 8 below.

200

201 **Table 7. Insulating Fluid Data**

I	1	2	3	4	5	6	7	8
x_i	-1.6608	-0.2485	-0.0409	0.2700	1.0224	1.5789	1.8718	1.9947
r_i	0	0	3	0	3	0	0	5

202

203 **Table 8. Parameter Estimates Based on Insulating Fluid Data**

Estimator	σ	μ
MLE	0.9027	1.8757
Bayesian – Lindley’s Approach	0.9716	1.8511
Bayesian – Importance Sampling	1.4455	-0.2370
BLUE	1.4211	2.5867
BLEE	1.2786	2.4809

204

205 The results show that the MLE and the Bayes estimator based on Lindley’s approximation are close to
206 each other and somewhat smaller than the linear estimators. Based on our simulation study, the former
207 estimators are more precise and reliable.

208

209

210

211 **4. SUMMARY AND CONCLUSION**

212

213 In this study, based on progressively type II censored data, we considered point estimation of location
214 and scale parameters in type II Generalized Logistic Distribution (Type II GLD). We developed three
215 estimators (ABLUE and ABLEE and Importance Sampling Estimator) for the unknown parameters.
216 We also included the maximum likelihood estimators (MLE) and Bayes estimators approximated by
217 the Lindley’s Approach for comparison purposes.

218 The results of the simulation study reveal that MLE and Lindley’s approximation to the Bayes
219 estimator perform better than the other estimators developed in this paper. They have the smallest bias
220 and MSE values as shown during the simulation study. As for the effect of the parameter α value on
221 the location and scale estimator’s bias and MSE values, ee got better results for smaller values of α .

222 The conclusion of this work is that the MLE has the overall best performance for estimating the
223 parameters of the type II generalized logistic distribution. However, for small sample sizes, numerical
224 problems can occur. In such situations, the approximate linear estimators like the ABLUE and
225 ABLEE can provide a viable alternative. The Bayes estimator performs very well too, especially the
226 approximation based on Lindley’s approach.

227

228

229 **ACKNOWLEDGEMENTS**

230

231 The authors would like to thank the editor and the referees for their helpful comments that
232 improved the presentation of the paper. This research was supported by a grant from the Office of
233 Research Support at Qatar University, grant no. QUST-2-CAS-2021-155.

234

235

236

237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Aljarrah, M., Fayome, F. and Lee, C. Generalized logistic distribution and its regression model. *Journal of Statistical Distributions and Applications*. 2020: 7(7), <https://doi.org/10.1186/s40488-020-00107-8>.
2. Azizpour, M., and Asgharzadeh, A. Inference for the Type-II Generalized Logistic Distribution with Progressive Hybrid Censoring. *Journal of Statistical Research of Iran*, 2018: 14 (2):189-217.
3. Balakrishnan N., and Cramer, E. *The art of progressive censoring, Application to Reliability and Quality*. Birkhauser; 2014.
4. Balakrishnan, N. and Hossain, A. (2007), Inference for the Type II generalized logistic distribution under progressive Type II censoring, *Journal of Statistical Computation and Simulation*. 2007: 77(12): 1013 – 1031.
5. Balakrishnan N., and Sandhu, R. A. A Simple Simulational Algorithm for Generating Progressive Type-II Censored Samples, *The American Statistician*. 1995: 49(2): 229-230.
6. Balakrishnan, N., and Sandhu, R. A. Best Linear Unbiased and Maximum Likelihood Estimation for Exponential Distributions under General Progressive Type-II Censored Samples, *The Indian Journal of Statistics, Series B*. 1996: 58(1): 1-9.
7. Kousik, M. and Kayal, S. Estimation of parameters and reliability characteristics for a generalized Rayleigh distribution under progressive type-II censored sample, *Communications in Statistics - Simulation and Computation*. 2019: 50(11): 3669 – 3698.
8. Nassar M.M., and Elmasry, A. A study of generalized logistic distributions, *Journal of the Egyptian Mathematical Society*. 2012: 20(2): 126 – 133.
9. Nelson W. *Applied Life Data Analysis*. NewYork : John Wiley and Sons; 1982.
10. Rimawi, Rana and Baklizi, A. Estimation in the Type II Generalized Logistic Distribution Based on Progressively Type II Censored Data. *Proceedings of the GSRD International Conference*. Dec. 2021: 28 – 35.
11. Salah, M.M. On progressive Type-II censored samples from alpha power exponential distribution. *Journal of Mathematics*. Vol. 2020, Article ID 2584184, <https://doi.org/10.1155/2020/2584184>