



SDI EDITORIAL COMMENTS FORM

| EDITORIAL COMMENT'S on revised paper (if any) | Authors' response to editor's comments |
|--|---|
| <p>There are more than hypotheses. What of them was used as a base to calculate sample size or statistics test hypothesis? The author used mean and S.S.(what is this?) and use non-paramethric method.</p> | <p>Dear Editor,</p> <p>Totally 6 hypotheses, 1 of which was the main hypothesis and 5 of which were the sub-hypotheses- were made in order to demonstrate the views of 569 participants- 385 of whom were Turks and 184 of whom were Arabs- concerning human milk banks.</p> <p>For the research in Turkey, 385 participants were considered adequate to represent the population of up to 100 million participants with 95% reliability taking the sample sizes accepted for certain populations into consideration [25]. 385 data was sufficient to calculate the level of significance. It is calculated with the statistical formula where this reference is made.</p> <p>The averages and standard deviations of the expressions in the scale were calculated and comments were made. Interpreted according to the average of the answers given by the participants.</p> <p>Normality test was performed and non-parametric tests were used. Shapiro Wilk and Kolmogorov Smirnov teats were used to test whether or not the research data had normal distribution. It was found as a result that the data did not have normal distribution ($p < 0.000$). Mann Whitney U test was used for two-class data in comparing the differences between independent groups whereas Kruskal Wallis H test was used for multi-class data and Hochberg GT2 test was used to find the groups with the sources of differences.</p> |