

Prevalence *Staphylococcus aureus* isolated from some street Hot Beverages in Abidjan, Côte d'Ivoire.

ABSTRACT

Aims: *Staphylococcus aureus* is one of the main pathogens found in street food, including hot beverages. However, information about *S. aureus* isolated from street hot beverages from coffee carts is very limited. We aimed to characterize *S. aureus* isolated from street hot beverages sold in Abidjan.

Methodology: The identification was made through conventional microbial and biochemical analysis. Macroscopic identification on the Baird Parker agar with tellurite egg yolk. Microscopic observation through Gram staining as well as biochemical tests such as catalase; DNase and coagulase were performed. to confirm staphylococcal strains, the MALDI-TOF mass spectrometry was used. After identification, the susceptibility of the staphylococcal isolates was evaluated using disc diffusion method.

Results: Result showed that most of *Staphylococcus aureus* (18.4%) were isolated from tea. All the strains of *staphylococcus aureus* isolated from street beverages were sensitive to cefoxitin and vancomycin. All the *S. aureus* isolated from milk were resistant to Erythromycin. Although beverages are consumed hot, the presence of *S. aureus* in ready-to-drink beverage may cause infections to consumers.

Conclusion: This drink need attention for the seller and the user to avoid some infections.

Keywords: Staphylococcus aureus, hot beverages, Susceptibility to antibiotics, food safety, Abidjan.

1. INTRODUCTION

Street food have easy access and generate job opportunities, especially for low- or uneducated people [1]. The hot drink culture is highly developed in industrialized countries and constitutes an important part of street catering. The developing country use the type of drink, but several studies have shown that street food use raw materials and ingredients of poor microbiological quality and non-authorized person are involved in this sector of activity [2-3]. Food safety remains a global concern despite efforts made by health and sanitation organizations [5]. Food contamination remains a real and permanent risk, especially in cities where this risk is closely linked to the consumption of ready to eat street foods [6-7]. Foodborne infections are reported worldwide, and it was estimated to be involved in more than 2 million deaths per year, mainly children [8], due solely to food and water inadequate sanitation and hygiene. Across the world and specifically in Africa, several countries have experienced episodes of food poisoning without scientifically proofed of given food involvement. Several bacteria are involved in food poisoning [7, 9].

Staphylococcus aureus is a ubiquitous commensal bacterium on human skins and anterior nares, but frequently causes severe infections in humans [10]. *S. aureus* has been described as one of the most common causal agents of food poisoning associated with the consumption of raw milk cheese and is also listed as one of the pathogens likely to pose the greatest risk to human health [11-13]. The presence of this pathogen in final products is related to several factors but is usually associated with the use of contaminated raw milk and endogenous starter cultures in manufacturing the cheese, as well as with asymptomatic human carriers of *S. aureus*[14-15]. Another serious public health problem associated with *S. aureus* in food is its resistance to antimicrobials because it triggers serious infections with limited therapeutic options [16-17]. Methicillin-resistant *S. aureus* (MRSA) is a leading example of bacterial resistance in the world [16] and is characterized by acquisition of the *mecA* gene, which is located on the chromosomal cassette *mec* (SCC*mec*) [18]. For many years, MRSA staphylococcal isolates have been recognized as pathogens unique to hospitals. However, studies have documented the spread of MRSA in foods [19]. The aim of this study was to investigate the presence of *S. aureus* in some hot beverages (coffee, tea and cocoa) and the antimicrobial resistance profile of staphylococcal isolates from those street hot beverages collected from Abidjan, Côte d'Ivoire.

2. MATERIAL AND METHODS

2.1. Survey

The survey was carried out in five municipalities (Abobo, Adjamé, Yopougon, Cocody and Port-Bouët) of Abidjan to draw the characteristics of consumers of street Hot Beverages.

2.2. Hot beverages sample's collection

Per municipalities, three kinds of hot beverages (tea, coffee, and/or cocoa) were targeted. A total of 400 samples of hot beverage made of tea (139), coffee (138), or cocoa (123) were collected in sterile tubes (Table 1). The samples were collected hot in stomacher paper and transported immediately or within 2 hours in cooler to the laboratory for microbiological analyzes.

Table1. Distribution of samples collected according to the collected places and the kind of hot beverages

Communes	Tea	Coffee	Cocoa	Total
Abobo	28	26	23	77
Adjamé	28	26	26	80
Yopougon	27	29	27	83
Cocody	27	29	25	81
Port-Bouët	28	29	22	79
Total	138	139	123	400

2.3. Microbial Analysis

For the microbiological analysis, 1 ml of each collected beverage samples was used for a decimal dilution. Each dilution (1 ml) was spread on Baird Parker agar (Biokar Diagnostics, France) with egg yolk [20] before its incubation at 37°C for 48 hours for presumptive identification of *Staphylococcus sp.* For the search *S. aureus* referencing strain of *S. aureus* ATCC 25923 was used as a positive control. Macroscopic identification was made on Baird Parker agar with tellurite egg yolk. Microscopic observation through Gram staining as well as biochemical tests such as catalase, DNase and coagulase and fermentation of mannitol were performed.

2.4. Identification of isolates by MALDI-TOF MS

After isolation, the MALDI-TOF (Bruker Daltonics™) mass spectrometry was used to confirm the *Staphylococcus aureus* identification. Samples were covered with 1.5 µl of the matrix solution (Sigma, Lyon, France). After 10 min of sonication, a centrifugation (13000 g, 5min) was performed before transferred to a clean polypropylene tube. The target plate and matrix were then dried at room temperature before identification [21]. A positive control was used (*S. aureus* ATCC 25923).

2.5. Antibiotic susceptibility testing

Antibiotic susceptibilities were determined by the disc diffusion method on Mueller-Hinton agar (Oxoid, France) according standards and recommendations of the Antibiogram Committee of the French Society of Microbiology [22]. The antimicrobial drugs tested and their concentrations on the discs of antibiotics (Bio-Rad, Marnes-LaCoquette, France) were as following: penicillin (PEN 10µg), amoxicillin (AMX 25 µg), cefoxitin (FOX 30 µg), tobramycin (TM 10 µg), gentamicin (GM 15 µg), erythromycin (ERY 15 µg), clindamycin (CMN 2 µg), levofloxacin (LVX 5 µg); norfloxacin (NOR 5 µg), ofloxacin (OFX 5 µg), vancomycin (VAN 30 µg). The reference strain *Staphylococcus aureus* ATCC 25923 was used as test control processed in the same conditions. Isolates were classified as susceptible or resistant to the drug.

2.6. Data Analysis

The data obtained from the survey and observations checklists were analyzed using the SPSS version 20.0 statistical software package, and then exported to Microsoft Excel for scoring. Descriptive analyses were used to summarize the variables of interest and determine relationships between them. The results were expressed as mean ± standard deviations (±SD), frequencies, and percentages. The Chi-squared frequency test (χ^2 test) was used to test the relationships between the variables. Statistically significant differences were based on 95% confidence limits.

3. RESULTS AND DISCUSSION

3.1. Results

3.1.1. Probable origin of contaminants

The probable origin of contaminants is related to hot beverage process are compile in table 2. Thus, during hot beverage preparation, some contaminants occurred related to the process. Also, the highest factor risks were recorded through the treatment of products, materials or ingredients. The cooking and conservation process are also one of the determining factors linked to the beverage poisoning risks. The quality of water used to prepare and rinse recyclable cup could be a supplement contamination factor. Hot beverage contamination varied according to several factors.

Table 2: Probable risk related to beverage consumption sold by street vendors

Category	Probable contaminants level	
Ingredients	Coffee, Cocoa, Tea	+
	Sugar, Lemon, Mint	+
Materials	Thermos for hot water, Spoon, Knife, Scissors	+
	Lemon squeezer	++
Others	Water used to prepare and rinse recyclable cups	++
	Disposable cups	+
	Recyclable cups	++

Handling of products, materials or ingredients	+++
Beverage cooking process	++
Hygiene of vendors	++

+ Low risk, ++ Moderate risk, +++ High risk

3.1.2. Phenotypic Characterization and Species Identification

All staphylococcal isolates were confirmed as gram-positive cocci, catalase positive, and coagulase and DNase positive. Presumptive staphylococcal isolates were confirmed at the species level by proteomic analysis using MALDI TOF MS, validating identification of the species as *S. aureus*. The global contamination rate of the sampled hot beverage was 8.5% and vary according to the kind of beverage (Table 3) and locality of collections (Table 4). According to the kind of beverage, the highest *S. aureus* contamination level was recorded with tea samples (18.4%), followed by cocoa's (8.7%) and coffee's (1.4%).

Table 3. distribution of *S. aureus* strains according to the sampled hot beverages

Beverages	Proportion of <i>S. aureus</i> % (n/N)
Tea	18.4% (26/138)
Coffee	1.4% (2/139)
Cocoa	8.7% (6/123)
Total	8.5% (34/400)

The table 4 indicated that the most contaminated samples by *Staphylococcus aureus* were collected from Yopougon(15.7%) followed by Adjamé(13.7%). The lest contaminated commune was Cocody (1.2%).

Table 4. Distribution of *Staphylococcus aureus* in communes

Commune	<i>S. aureus</i> n (%)
Abobo	2 (2.6)
Adjamé	11 (13.7)
Yopougon	13 (15.7)
Cocody	1 (1.2)
Port-Bouët	6 (7.8)
Total	34 (8.5)

3.1.3. Antibiotic susceptibilities of *Staphylococcus aureus* isolated from street beverages

Globally, there is a variability of resistance profile of the isolated *S. aureus* strains (Figure 1). All the isolated strains from street beverages were sensitive to cefoxitin and vancomycin (Figure 1). In addition, most of *S. aureus* were sensitive to the aminosides and quinolones tested.




Figure 1. Global antibiotic susceptibility of *Staphylococcus aureus* isolated from street beverages

3.1.4. Distribution of *S. aureus* resistance according to the beverages

According to the kind of beverage, it is observed a variability of resistance profile (Figure 2).




Figure 2. Distribution of isolated *S. aureus* resistance according to their origin

3.2. Discussion

For Lieberman *et al.* [23], foods and beverages which are prepared and sold on places like streets, market, festival areas and consumed on the run are known as streetfood. Street foods especially show the eating habits of people living in big cities such as Abidjan. Survey results showed that, majority of hot street beverage vendors were foreigners (88.0%) and illiterate (71.3%). Ma *et al.* [24] reported that preparing and selling food on the streets provide a constant income for millions of uneducated people. Regarding the employment, street food provides a good job opportunity and income for sellers with small capitals and especially for women [25-26]. In addition, Atobla *et al.* [27] showed that vendors' knowledge is poor and had lower education levels, which is also reflected in their largely inadequate facilities and unhygienic behavior while selling food.

The strains of *S. aureus* were isolated from samples of hot drinks, on all the samples analyzed, with high frequencies in tea (67.3%), cocoa (44.6%) and coffee (34.6%). Hot drinks exposed to contamination by *S. aureus* represent a significant risk for human health, in fact this pathogen is supposed to have been eliminated by heat during treatment therefore its presence in hot drinks is generally indicative of bad hygienic conditions as highlighted [28]. These results are evidence of manual contamination. The spread of *S. aureus* by hot drink handlers is a concern and should be given careful attention in the manufacturing and sales chain.

The variation in the resistance to antibiotics of strains of *Staphylococcus aureus* isolated from hot drinks based on coffee, tea and cocoa is important because strains resistant to several antibiotics have been obtained. Resistance varies depending on the antibiotic used.

Note that all strains are sensitive to levofloxacin, norfloxacin, ofloxacin, ceftriaxone and vancomycin. These results have also been reported by Attien et al. [17]. In addition, certain antibiotics which were initially sensitive have increasingly become resistant. Variable resistance was observed depending on the origin of the strains to antibiotics such as penicillin and erythromycin, amoxicillin and gentamicin. This trend was obtained by Resch et al. [29], who showed that strains of coagulase staphylococci isolated from food had resistance to oxacillin. This is reported by Magiorakos et al. [30] who states that today almost all strains of *S. aureus* have acquired resistance to penicillin. These results could be an indicator of a prophylactic route for the treatment of *S. aureus* infections. Penicillin was the first antibiotic discovered in 1929 by Alexander Fleming, *S. aureus* were sensitive at this time. Resistance to oxacillin indicates the presence of methicillin-resistant strains among the strains isolated. This would increase the risk of infection in the event of contamination by *S. aureus* methi-resistant which is very difficult to manage. However, additional tests will clarify the presence of this type of strain.

4. CONCLUSION

To our knowledge, this is the first hot beverage study to assess the characteristic of vendors with hand push mobile carts in Abidjan (Côte d'Ivoire). Street vendors preferred drinking coffee to boost energy to sell. According to them, tea was the most beverage consumed by Ivoirians. The results from this street vendor showed that most street vendors are foreigners and illiterate. These street vendors therefore need training in basic hygiene practice to avoid risks such as food contamination or throwing plastic cups on the street.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

- [1]. Khairuzzaman M, Chowdhury FM, Zaman S, Al Mamun A, Bari ML. Food Safety Challenges towards Safe, Healthy, and Nutritious Street Foods in Bangladesh. *Int. J. Food Sci.* 2014, ID. 483519.
- [2]. Bereda TW, Emerie YM, Reta MA, Asfaw HS. Microbiological Safety Of Street Vended Foods In Jigjiga City, Eastern Ethiopia. *Ethiop. J. HealthSci.* 2016; 26(2): 161–170.
- [3]. Idowu O, Rowland S. Oral Fecal Parasites and Personal Hygiene of Food Handlers in Abeokuta, Nigeria. *Afr. Health Sci.* 2006; 6(3): 160-164.
- [4]. Rane S. Street Vended Food in Developing World: Hazard Analyses. *Ind. J. Microbiol.* 2011; 51(1), 100–106.

- [5]. Ackah M, Gyamfi ET, Anim AK, Osei J, Hansen JK, Agyemang O. Socio-Economic Profile, Knowledge of Hygiene and Food Safety Practices Among Street-Food Vendors In Some Parts Of Accra-Ghana. *J Food Safety*.2011; 13: 191-197.
- [6]. Tambekar D, Jaiswal V, Dhanorkar D, Gulhane P, Dudhane M. Identification of Microbiological Hazards and Safety of Ready-To-Eat Food Vended Streets of Amravati City, India. *J. Appl. Biosci*. 2008; 7: 195-201.
- [7]. Sina H, Attien P, Wélé M, Socohou A, Boukary-Chabi A, Dougnon VT, Baba-MoussaF, AdjanohounA, Baba-Moussa L. Sanitary risk factors and microbial contamination of grilled meats sold in cotonou, Benin. *J Food Security*. 2019; 7(5): 175-182.
- [8]. World Health Organization. Initiative to Estimate the Global Burden of Foodborne Diseases.2015 Available from: http://www.who.int/foodsafety/foodborne_disease/ferg/en/index.html
- [9]. Hernández-Cortez C, Palma-Martínez I, Gonzalez-Avila LU, Guerrero-MandujanoA, Solís RC, Castro-Escarpulli G. Food poisoning caused by bacteria (food toxins). *Poisoning: From Specific Toxic Agents to Novel Rapid and Simplified Techniques for Analysis*,IntechOpen, 2017; 33DOI: 10.5772/intechopen.69953. Available from: <https://www.intechopen.com/chapters/56521>
- [10]. Kluytmans J, van Belkum A, Verbrugh H: Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. *Clin Microbiol Rev* 1997, 10(3):505-520.
- [11]. Spanu V, Spanu C, ViridisS, Cossu F, Scarano C, De Santis EPL. Virulence factors and genetic variability of *Staphylococcus aureus* strains isolated from raw sheep's milk cheese. *Int J Food Microbiol*. 2012, 153(1-2): 53-57.
- [12]. Sugrue I, Tobin C, Ross RP, Stanton C, Hill C. Foodborne pathogens and zoonotic diseases. *InRaw milk*, 2019; 259-272.
- [13]. Castro RD, Pedroso SHSP, Sandes SHC, Silva GO, Luiz KCM, Dias RS, Filho RAT, Figueiredo HCP, Santos SG, Nunes AC, Souza MR. Virulence factors and antimicrobial resistance of *Staphylococcus aureus* isolated from the production process of Minas artisanal cheese from the region of Campo das Vertentes, Brazil *J Dairy Sci*. 2020;103(3):2098-2110. doi: 10.3168/jds.2019-17138.
- [14]. Castro RD, Oliveira LG, Sant'Anna FM, Luiz LMP, Sandes SHC, Silva CIF, SilvaAM, NunesAC, PennaCFAM, Souza MR. Lactic acid microbiota identification in water, raw milk, endogenous starter culture, and fresh Minas artisanal cheese from the Campo das Vertentes region of Brazil during the dry and rainy seasons. *J Dairy Sci*.2016; 99(8): 6086-6096.
- [15]. Costanzo N, Ceniti C, Santoro A, Clausi MT, Casalnuovo F. Foodborne Pathogen Assessment in Raw Milk Cheeses. *Int J Food Sci*. 2020 :3616713. doi: 10.1155/2020/3616713.
- [16]. Ibed AN, Hamim SS. Molecular detection of methicillin resistant *staphylococcus aureus* isolated from burns infection in Al-nasiriyah city. *World J Pharmaceutical Sci*, 2014: 950-954.
- [17]. Attien P, Sina H, Moussaoui W, Dadié T, Chabi-Sika K, Djeni T, Bankole HS, Kotchoni SO, Edoh V, Prevost G, Dje M, Baba-Moussa L. Prevalence and antibiotic resistance of *Staphylococcus* strains isolated from meat products sold in Abidjan streets (Ivory Coast). *Afr J Microbiol Res*. 2013; 7 (26): 3285-3293

- [18]. Turlej AGATA, Hryniewicz WALERIA, Empel J. Staphylococcal cassette chromosome mec (Sccmec) classification and typing methods: an overview. *Polish J Microbiol*, 2011; 60(2): 95.
- [19]. Crago B, Ferrato C, Drews SJ, Svenson LW, Tyrrell G, Louie M. Prevalence of *Staphylococcus aureus* and methicillin-resistant *S. aureus* (MRSA) in food samples associated with foodborne illness in Alberta, Canada from 2007 to 2010. *Food Microbiol*. 2012; 32(1), 202-205.
- [20]. Baird-Parker AC. Foodborne salmonellosis. *Lancet*, 1990; 336: 1231-1235.
- [21]. Pfeleiderer A, Lagier JC, Armougom F, Robert C, Vialettes B, Raoult D. Culturomics identified 11 new bacterial species from a single anorexia nervosa stool sample. *Eur J Clin Microbiol Infect Dis*. 2013, 32(11): 1471-1481.
- [22]. CASFM "Comité de l'antibiogramme de la Société Française de Microbiologie : Recommandations 2020 V.1.1," 181 p, 2020. Available on https://www.sfm-microbiologie.org/wp-content/uploads/2020/04/CASFM2020_Avril2020_V1.1.pdf
- [23]. Lieberman HR, Stavinoha T, McGraw S, White A, Hadden L, Marriott BP. Caffeine use among active-duty U.S. Army soldiers, *J Acad Nutr Diet*, 2012; 112 (6): 902–912, e901–e904, <https://doi.org/10.1016/j.jand.2012.02.001>.
- [24]. Ma L, Chen H, Yan H, Wu L, Zhang W. Food safety knowledge, attitudes, and behavior of street food vendors and consumers in Handan, a third-tier city in China, *BMC Public Health*. 2019; 19, 1128, <https://doi.org/10.1186/s12889-019-7475-9>.
- [25]. Marangoni F, Pellegrino L, Verduci E, Ghiselli A., Bernabei R, Calvani R, Cetin I, Giampietro M, Perticone F, Piretta L, Giacco R, Vecchia CL, Brandi ML, Ballardini D, Banderali G, Bellentani S, Canzone G, Cricelli C, Faggiano P, Ferrara N, Flachi E, Gonnelli S, Macca C, Magni P, Marelli G, Marrocco W, Miniello VL, Origo C, Pietrantonio F, Silvestri P, Stella R, Strazzullo P, Troiano E, Poli A. Cow's Milk Consumption and Health: A Health Professional's Guide, *J Amer Coll Nutr*, 2018; 38, 197-208, <https://doi.org/10.1080/07315724.2018.1491016>.
- [26]. Medeiros LC, Hillers VN, Chen G, Bergmann P, Kendall V, Schoreder M. Design and development of food safety knowledge and attitude scales for consumer food safety education, *J Amer Diet Assoc*, 2004; 104 (11), 1671-1677, <https://doi.org/10.1016/j.jada.2004.08.030>.
- [27]. Atobla K, Akoa E, Bonny AC, Dadié A, Karou TG, Niamke S. Pork supply chain, consumption, and risk factors for infections of consumers in the Abidjan district (Côte d'Ivoire), *Food Environ Safety*, 2018; 16 (17), 20-31.
- [28]. Giaccone V, Catellani P, Alberghini L. Food as cause of human salmonellosis. *Salmonella a dangerous foodborne pathogen*, Ed. Barakat SM Mahmoud, 2012 ; 47-72.
- [29]. Resch M, Nagel V, Hertel C. Antibiotic resistance of coagulase-negative staphylococci associated with food and used in starter cultures. *Int J Food Microbiol*. 2008; 127(1-2), 99-104.

- [30]. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S, Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J, Struelens MJ, Vatopoulos A, Weber JT, Monnet DL. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect.* 2012 18(3):268-81. doi: 10.1111/j.1469-0691.2011.03570.x.

UNDER PEER REVIEW