

IMPACT ON KNOWLEDGE OF BLACKGRAM GROWERS IN PERIYAR VAIGAI COMMAND AREA OF MADURAI DISTRICT

ABSTRACT

The World Bank Supported TN IAM (Irrigated Agriculture Modernization) Project is a follow up of IAMWARM presently it was called as (Irrigated Agriculture Modernization and Water-Bodies Restoration and Management) Project which has made significant development and impacts in the Tamil Nadu state by modernizing irrigation infrastructure, improving water use efficiency, enhancing yield and productivity of agriculture in a climate resilient production systems, diversification towards high value crops, strengthening the institutional reforms through Participatory Irrigation Management (PIM) and Water Users Association (WUA). Madurai District of Tamil Nadu was purposively selected for this study because Tamil Nadu Irrigated Agriculture Modernization Project was conducted under Tamil Nadu Agricultural University. The foremost objective of the study is to assess the knowledge level of respondents in study area. According to crop production technology aspects revealed that (79.00%) of the beneficiaries possessed knowledge in using VBN 6 variety for cultivation. Majority (80.00%) of the beneficiaries were possess knowledge on (cultural control) fixation of light traps, crop rotation (77.50%) and sowing carry out in proper season.

Key words: TN IAM project, Water use efficiency, Crop production, beneficiaries.

1. Associate Professor (Agrl.Extn), DAE&RS, AC&RI, TNAU, Madurai

Introduction

Agriculture is the single largest consumer of water in the state, using 75% of the state's water. Out of 40000 tanks only 14098 Tanks are under the maintenance of the Water Resources Department with the rest under the Panchayat unions. Irrigation through a combination of canals, wells and tanks increases the reliability and availability

of water for farming and is essential for cultivating crops in much of state. About 3 million ha of land (54 percent of the total crop land) is under irrigation. An Expert Committee on "Development and Management of Water Resources" observed that bringing additional area into cultivation is remote but the challenge is how best to bridge the gap in cultivation by reducing demands, by effective water management and by adoption of modern agricultural techniques (Micro Irrigation etc.) The Committee recommended Integrated Water Resources Management & convergence of various Departments for development and management of water resources in Tamil Nadu. This study will be much useful to researcher, extension workers and policy makers as to what extent the technology were adopted by the beneficiaries besides reflecting on the impact of the project on farm and home of TN-IAMP beneficiaries.

Materials and methods

Tamil Nadu state was purposively selected for the study as the researcher belongs to the same state. Madurai District of Tamil Nadu was purposively selected for this study because Tamil Nadu Irrigated Agriculture Modernization Project was conducted under Tamil Nadu Agricultural University. Sirumalaiyar and sathaiyar sub-basins were covered by the Madurai district which is having more ayacut area. Madurai district consists of 7 taluks. Among the seven taluks, Madurai north and Vadipatti taluk were selected. The sample size of 120 beneficiaries were selected by using purposive sampling method. The data collection was done with the use of a well-structured and pre-tested interview schedule. The reliability of the data could be more in the interview method. The collected data were analyzed by descriptive statistics and Karl Pearson's coefficient of correlation using SPSS software.

Results and Discussion

Practice- wise level of knowledge

The practice wise knowledge level of TN-IAMP beneficiaries was studied and the results are present in Table 1.

Table 1 Distribution of respondents according to their practice wise extent of knowledge (n=120)

S. No	Practices	Knowledge level	
		Number	Per cent
I	Crop production technologies		
1.	Land should be free from volunteer plants like dry root rot	76	63.33
2.	land should be with proper drainage facility	91	75.83
3.	Land should be red soil or alluvial soil	108	90.00
4.	Make the land to fine tilth condition	95	79.17
5.	Application of 12.5 t/ha of FYM/coir pith during land preparation	84	70.00
6.	Mid-July to mid-august and mid-January to mid-February is suitable season	88	73.33
7	VBN 6 is the variety used for black gram cultivation	94	79.00
8	Duration of VBN 6 is 65-70 days	120	100.00
9.	Carbendazim and thiram are mostly used chemicals for seed treatment	85	70.83
10	Treat the seeds with carbendazim or thiram @ 2g/kg of seeds 24 hrs before sowing	83	69.17
11	Treat the seeds with <i>pseudomonas fluorescens</i> @ 10 g/kg or <i>trichoderma viride</i> @ 4 g/kg of seeds 24 hrs before sowing	87	72.50

12	Rhizobium is mostly used bio fertilizer for treating seeds	78	65.00
13.	Treat the seeds with rhizobium @ 600 g/ha of seeds required along with 500 ml of rice kanji	79	65.83
14.	Rhizobium and phosphobacteria are bio fertilizer used for basal application	82	68.33
15	In case of seed treatment is not done, 10 packets of rhizobium (2kg/ha) and 10 packets of phosphobacteria (2kg/ha) along with 25 kg of FYM and 25 kg of sand applied before sowing	92	76.67
16.	20 kg/ha seed required	104	86.67
17.	Seed dibbling is method of sowing	67	55.83
18.	45X10 cm is spacing	71	59.17
19.	Application of 5 t/ha of FYM during last ploughing of land preparation	74	61.67
20.	25 kg/ha of nitrogen fertilizer applied as basal	67	55.83
21.	50 kg/ha of phosphorous fertilizer applied as basal	56	46.60
22.	25 kg/ha of potassium fertilizer applied as basal	66	55.00
23.	Pendimethalin and fluchloralin are mostly used pre-emergence herbicides	81	67.50

24.	Spraying of pendimethalin @ 2.5 l/ha or fluchoralin 30% EC @1.5 l/ha at 3 rd or 5 th day after sowing with hand sprayer	81	67.50
25.	Soil digging should be done at 30 th day after sowing to prevent weeds	81	67.50
II	Crop protection technologies		
26.	Sowing should be done in proper season	92	76.60
27.	Crop rotation should be done	93	77.50
28.	Fixation of light traps to attract gram pod borer adult stage	96	80.00
29.	Collect all stages of pest and destroy it	56	46.67
30.	Removal of yellow mosaic virus affected plants from the field to control viral diseases	67	55.83
31.	Fix the pheromone trap @ 12 no/ha to control borer	62	51.67
32.	To control yellow mosaic virus, seed treatment of seeds with imidacloprid @ 5 ml/kg of seeds 24 hrs before sowing	92	76.67
33.	Seed treatment with <i>trichoderma viride</i> @ 4g/kg or <i>pseudomonas fluorescens</i> @ 10 g/kg of seeds to control dry root rot	70	58.33
34.	Spraying of 0.1% carbendazim on plant parts to control dry root rot	67	55.83
35.	Spraying of carbendazim @ 2.5 kg/ha on plant parts to control powdery mildew disease	79	65.83

	Harvest and storage		
36.	Pods turns brown or black with hard seeds inside pods are harvesting indices	94	78.33
37.	Plants are uprooted by hand or cut with sickle at the bottom of the plants	95	79.17
38.	Beating with flexible bamboo stick or by machinery to separation of seeds from harvested plants	92	76.67
39.	Drying of seeds by open sun dry up to 8-9 per cent of moisture content in seeds	86	71.67
40.	Agro moisture meter is instrument used to check moisture content in seeds	59	49.17
41.	Discoloured and broken seeds should be eliminated before grading	77	64.17
42.	Grading is done by sieving the seeds with BSS 7X7 wire mesh sieve	78	65.00
43.	Store the seeds in gunny bags or cloth bags for short term storage (8-9 months) with seed moisture of 8-9 per cent	70	58.33
44	Store the seeds in polyline gunny bags for medium term storage (12-15 months) with seed moisture of 8-9 per cent	78	65.00
45	Store the seeds in polythene bags for long term storage (more than 15 months) with seed moisture of 8 per cent	93	77.50

Knowledge level on crop production technologies

It could be observed from above Table 1 according to crop production technology aspects reveals that (79.00%) of the beneficiaries possessed knowledge in using VBN 6 variety for cultivation and duration of crop be 65-70 days and maximum number of beneficiaries (90.00%) had knowledge on using alluvial / red soil land and using recommended seed rate of cultivation.

With regard to equal level of knowledge accompanied by beneficiaries on several practices possess (70.83%) on recommended chemicals and averagely (70.00%) of beneficiaries possess knowledge on land preparation, maintaining recommended seasons (73.33%), bio-fertilizer usage on seed treatment and land preparation (60 – 70 per cent) and recommended spacing (59.17%) on crop production aspects.

Knowledge level on crop protection technologies

It could be observed from above Table 1 regarding crop protection measures beneficiaries are good in knowledge level, whereas majority (80.00%) of the beneficiaries were possess knowledge on (cultural control) fixation of light traps, crop rotation (77.50%) and sowing carry out in proper season.

With regard to equal distribution (55-76 per cent) of beneficiaries possess knowledge on biological and chemical control of pest & diseases.

Knowledge level on harvest and storage

It could be observed from the Table 1 majority of the beneficiaries (78.33%) were possess knowledge on harvest indices (colour turns brown to black), harvesting methods (79.17%) (Hand removal / by sickle) and processing (76.67%) (Separation of seeds and drying).

Conclusion

The success and experiences of the project need to be regularly and continuously shared among the black gram growers through publications, websites, newsletter and mass media. It has been observed from the study on the part of farmers, scientists and extension functionaries have been enhancing the adoption levels to create impact on the TN-IAMP beneficiaries. These linkage need to continue and strengthened all the stakeholders for perusing positive impact. Mostly 80.00 per cent of the TN-IAMP black gram growers were medium level in information seeking behavior. Major sources of information being trainings, demonstrations and publications. Hence, interventions on these aspects need to be continued and strengthened to earn more benefit to all black gram growers of project.

References

1. *Palanisamy, A. (2011). Impact Of TN-IAMWARM Project on the Farm and Home of Precision Farming Beneficiaries- An Analysis, M.Sc(Agri) Thesis, Department of Agricultural Extension and Rural Sociology, TNAU, Coimbatore.*

2. Ravichandran, V. K & Prakash, K. C. (2015). *Socio-Economic Impact of SRI and Traditional Rice Cultivation in Villupuram District of Tamil Nadu: Experiences from TN-IAMWARM Project*, *International Journal of Agricultural Sciences*, Vol.11 No1, PP. 166-171.
3. Sindhuja, P., & Shanthasheela, M. (2017). *Profile of Precision Farming Vegetable Beneficiaries in Ongur Sub-Basin Under TN-IAMWARM project*, *International Journal of Research in Agriculture and Forestry*, vol 4(2), P: 10-15.
4. Sangeetha, S. 2009. *Study on Factors Influencing the Adoption of Precision Farming Technologies in Tomato Cultivation*. Unpub. M.Sc. (Ag.) Thesis, AC&RI, TNAU, Coimbatore.
5. Sudha, T. 2008. *Prospects of precision farming in Dharmapuri district – A multidimensional analysis*, Unpub. M.Sc.(Ag.) Thesis, AC&RI, TNAU, Madurai.

6.