

# **Crop residue based complete feed for enhancing livestock performance- A review**

---

## **Abstract**

The estimated projected data regarding demand and supply of feed and fodder shows the 13.20% and 18.43% deficit between demand and supply of dry and green fodder to livestock, respectively. Shortage of land for cultivation of fodders and increased human as well as livestock population has led to heavy competition for feed grains that necessitates using the crop residues for the feeding of livestock. These crop residues are rich in fibre and low in other nutrients; they also have low palatability and digestibility. The best way to efficiently utilize the crop residues is to prepare complete feeds from pretreated crop residues + concentrates mixture. Pretreatment of crop residues increases digestibility by increasing rate of delignification, increasing enzymatic hydrolysis of cellulose, hemicelluloses and thus increasing the glucose yield. The complete feed is a quantitative mixture of all dietary ingredients to provide the specific nutrient requirement for various physiological functions of livestock. Crop residue-based complete feed could be prepared in mash, block and pellet (Expander and extruder) form. In this system, all feed ingredients including roughages, are proportioned, processed and mixed into a uniform blend. Different researchers undertook feeding trials of complete animal feed and conventional animal feed and reported better performance in terms of weight gain, milk yield and reproductive performance on feeding complete feed pellets or complete feed blocks to ruminants as compared to conventional ration. In conclusion, crop residue based complete feed could improve ruminant performance and reduces cost of feeding per kg gain or yield and thereby improving livestock based rural economy in developing countries like India.

**Keywords:** *Complete feed, Crop residue, Livestock, Performance*

## **INTRODUCTION**

Total livestock population as per 20<sup>th</sup> livestock census in India is 535.78 million, showing rise of 4.6% over the previous census. Therefore increased livestock population requires have feed and fodder requirement. Feed cost accounts about 60 to70% of total livestock production cost. Increased competition for grains between human being and livestock necessitates consideration of economical feed resources, treatment methods besides technologies for production of high valued livestock feed using them.

In India, different crop residues are produced every year. Some of them are utilized as fodder for the animals, burned as such to remove waste in land, or used as fuel and for composting. Day by day the feed and fodder requirement is increasing due to increased livestock population, farm mechanization etc. but, feed and fodder supply is deficit due to less availability of land for fodder cultivation, heavy competition for grains between human beings and animals. About 3, 62,793 million tonnes of crop residues are generated each year in India and large quantities of crop residues produced are burnt, contributing grossly to environmental pollution. Hence, it needs proper utilization and disposal.

To overcome the fodder shortage due to increase in livestock population, use of crop residues in the animal ration is must. The best way to efficiently utilize the crop residues is to prepare complete feeds from crop residues + concentrates mixture.

**Table 1: Demand and supply estimates of dry and green forages (million tonnes) [1]**

Year	Demand		Supply		Deficient		Deficit as %	
	Dry	Green	Dry	Green	Dry	Green	Dry	Green
2010	508.9	816.8	453.2	525.5	55.72	291.3	10.95	35.66
2020	530.5	851.3	467.6	590.4	62.85	260.9	11.85	30.65
2030	568.1	911.6	500.0	687.4	68.07	224.2	11.98	24.59
2040	594.9	954.8	524.4	761.7	70.57	193.0	11.86	20.22
2050	631.0	1012.7	547.7	826.0	83.27	186.6	13.20	18.43

Major crop residues produced from different crops are Maize stover, maize husk, Sorghum starw, groundnut straw, wheat straw, rice straw, arhar straw, gram straw, sugarcane bagasse, sugarcane tops, molasses, millet straw, soybean straw etc. The inclusion level of these crop residues in different complete livestock feed varies from 17.5 to 75 percent as per the physiological status of the animal and nutritive value of different crop residues.

### **Constraints in sole feeding of crop residues [2]**

- Crop residues are rich in fibre and low in nitrogen, minerals and vitamins.
- Their palatability, digestibility is low and therefore crop residues cannot form a sole ration for livestock.
- Crop residues are seasonally produced & have year-to-year variation in their production, hence need to be conserved.
- The crop residues are bulky in nature. Hence, they are expensive to transport, even for short distances.
- Some of the residues contains anti-nutritional factors e.g., sorghum and maize Stover contains tannin and paddy straw contains oxalate
- It is possible to bridge the gap between availability and requirement of roughages by processing of crop residues by using suitable method to improve nutrient availability of crop residues.

### **Concept of complete feed from crop residue**

- The complete feed is a quantitative mixture of all dietary ingredients to provide the specific nutrients requirement [3].
- In this system, all feed ingredients including roughages are proportioned, processed and mixed into a uniform blend.
- The product is used for feeding of livestock as a sole source of nutrients.
- The complete feeds can provide an adequate supply of balanced nutrients to the animal by controlling the concentrate to roughage ratio.
- It helps in improving utilization of low-quality fibrous crop residues.
- The system avoids feed wastage by promoting feed intake and by preventing refusal of the unpalatable feedstuff portions with reduction in feeding cost.
- Eating and rumination time can also reduced due to feeding crop residue based complete feeds to animals.
- In complete rations the proportion of other ingredients varies as per physiological status of animal and nutritive value of ingredients.
- These complete diets can be processed into mash or pellets or feed block.
- Mash feed can be pelleted by adopting either traditional steam pelleting or recently developed expander-extruder technology.

### **Pretreatment on different crop residues include following methods [4]**

Following are the pre-treatment methods mainly used for crop residues,

1. Physical method- It includes mechanical, thermal, ultrasound, electrokinetic and microwave method of pretreatment.
2. Chemical treatment includes treatment of acid, alkali, oxidative and organosolv treatment on crop residues.
3. Biological pretreatment method includes fungal, enzymatic and bacterial treatment on different crop residues.
4. Combined pretreatment includes steam explosion, extrusion and thermochemical methods of treatment.

Commonly used pretreatment methods are alkali treatment, urea treatment, steam explosion, ozonolysis and enzyme treatment.

### **Advantages of pretreatment on crop residues**

1. It improves palatability of crop residues and thereby intake.
2. It breaks lingo-cellulosic bond improves digestibility of crop residues.
3. It enhances enzymatic hydrolysis and thereby conversion of cellulose into fermentable sugars.
4. It enhances delignification of crop residues and thereby improves quality of crop residues. Melted lignin acts as a binding material in complete feed pellet or block making.
5. It increases glucose yield leads to improvement in animal production.

Crop residue-based complete feed could be prepared in mash, pellet, block or expander extruder pellet form.

### **Complete feed improves animal performance by following ways**

1. Stabilize rumen environment & improves nutrient utilization
2. Less fluctuation in release of ammonia and efficient utilization of ruminal NPN
3. Improves fibre digestion hence acetate production
4. Due to maintained Roughage: Concentrate ratio it maintains milk quality
5. Increase in energy density in feed
6. Increased digestibility of protein and improves nitrogen balance

- Improves total volatile fatty acid production (Propionate and acetate)

### Proportion of straws in complete feeds

- 30 to 70 percent roughages can be included in complete rations depending upon the physiological status of animals to be fed.
- The roughage content of complete diets can be as follows,

For high-yielding animals -30–40 percent

For growing animals -40–50 percent and

For dry animals -60–70 percent,

Whereas, Protein content in complete ration varies from 11 to 13 percent.

### The proportion of straws in complete feeds for milking animals

Table 2: Complete feed block is composed of forage, concentrate and other supplementary nutrients in desired proportions capable of fulfilling an animal's nutrient requirement[5].

Sr. No.	Milk yield of animal	Proportion of straw
1	5-10 liters/day	60%
2	10-15 liters/day	50%
3	15-20 liters/day	40%

### Ingredients of the concentrate mixture [5]

Ingredients includes-

- Oil cakes/ meals as a protein source
- Molasses, grains, grain byproducts as an energy source
- Bypass fat and protein for the preparation of complete diet for high yielders
- Micro minerals, vitamins
- Feed additives- probiotics, bentonite, enzymes, antioxidants, herbal extracts, etc.

### Advantages of complete feed [5]

- Complete feed has potential to fulfill the nutrient requirement of animals.
- It is possible to exploit locally available animal feed resources in complete feed to overcome the deficit of feed/fodder.
- There is scope for incorporation of tree leaves/shrubs in routine diets.
- Due to preparation of complete feeds quality feed is available round the year and is cost-effective.
- Complete feed prepared from bulky roughage sources is cheaper in storage and transportation.
- Complete ration feeding saves time and labour.
- Complete feed can correct multi-nutrient deficiencies in animals.
- Enhances utilization & palatability of poor quality/unconventional feeds through complete feeds.
- Reduces feed wastage

- It prevents selective feeding by animal.
- Complete feed facilitates farm management by allowing complete mechanization and more flexibility to include a wide range of alternative feeds.
- Low methane emission is possible by feeding complete feeds from animals.
- Feeding of complete feeds leads to improvement in productive & reproductive performance of animals.

### **Comparative study of the effect of feeding the complete feed in various forms**

Comparative growth performance study in goats and lambs on feeding crop residue-based complete mash Vs complete pellet feed was studied by [6]. Final body weight (Kg), ADG(g/d) and daily feed intake (Kg DM) was significantly higher and feed to gain ratio was better in goats and lambs fed pelleted crop residue based complete feed than complete mash feed.

[7] They reported significant ( $p < 0.05$ ) improvement in feed conversion efficiency with marked reduction in cost of production by feeding complete feed block than complete mash feed to Corriedale lambs, while DM and digestibility of gross nutrients, nutritive value and retention of major minerals did not change between the groups.

The higher average milk yield in complete feed block fed than in mash fed lactating buffaloes during an on-farm trial. Farmers were of opinion that complete feed blocks enhanced milk yield and were also easy at feeding and storage [8].

Effect of feeding processed sweet sorghum crushed residue-based complete rations in Murrah buffalo calves was studied by [9]. They found higher final body weight (224.70 kg), ADG (585.33 g/d), Feed intake (4.56 kg), better FCR (7.80) and low cost/Kg gain (Rs. 52.44) in Murrah buffalo calves fed Sweet sorghum crushed residue based complete expander extruder pellets than in Murrah buffalo calves fed either Sorghum straw based complete mash feed or Sweet sorghum crushed residue based complete mash feed

### **Milk production studies based on crop residue-based complete feed**

Dry matter intake was 28.2, 26.7 and 15.0 percent less on complete diets containing cotton stalks, maize cobs and sunflower heads, respectively, compared with the conventional diet. Average milk yield increased by 11.3, 11.5, 23.5 percent and cost of feed/kg milk yield reduced by 21.5, 23.4 and 25.2 percent on feeding complete diets containing cotton stalks, maize cobs and sunflower heads, respectively, compared with the conventional diets. The diets maintained 6–8 litres/day of milk yield in lactating Murrah buffalo. The results indicated that cotton stalks, maize cobs and sunflower heads could be incorporated in complete diets as sole roughage sources without any adverse effect on milk production in lactating Murrah buffalo [10]. The milk yield was 16.7 and 11.4 litres/day on feed blocks produced from premium and lower-quality sorghum Stover, respectively[11], indicating the feasibility of medium levels of milk production on crop-residue-based complete feeds.

[12] They reported increase in milk yield by 15.2% and FCM by 17.5% after feeding complete feed based on sunflower heads. They also reported higher fat and SNF by 19.7 % and 15.9%, respectively.

[13] An increase in milk yield was reported [by 16.23 % in buffaloes fed a complete diet containing concentrate+ sorghum fodder+ wheat straw in 50:35:15 proportions.

[14] It was observed that the milk fat, lactose, SNF and total solids were highest in cow fed with TMR than the conventional feed.

### **Meat production studies based on crop residue-based complete feed**

[10] They reported that the complete diets supported 87–108 g average daily gains (ADG) in Nellore ram lambs and 71 to 81 g ADG in local male kids in the intensive-feeding system. The ADG was significantly ( $P < 0.05$ ) higher on complete diets compared with the respective conventional diet in both species. The dry matter intake/kg gain decreased by about 20 percent in sheep and 15 percent in goats, while the cost of feed/kg gain decreased by 13 to 32 percent in lambs and 13–22 percent in kids compared with the conventional diet. Dressing percentage and meat: bone ratio was optimum and almost similar for complete and conventional diets in sheep and goats.

[15] Fed complete rations containing wheat straw and black gram straw at 60% level in crossbred sheep and reported the ADG (g/d) was 60 and 62.5, respectively. Whereas [16] found significantly higher ADG in Bharat Merino lambs fed complete diets containing 50% roughage and 50% concentrate than Bharat merino lambs fed conventional diet.

[17] Formulated Arhar based pelleted complete feed with roughage: Concentrate ratio, 60:40 with two different protein levels and fed to 12 non-descript goats (divided into two groups) for 140 days. The ADG found in the two groups were 75.7 and 72.9 g/d, respectively.

### **Effect of crop residue based complete feed on animal reproduction**

[18] He reported early maturity and age at first calving by 4-6 months in heifers. However, reduction in problems like anoestrus and repeat breeding with overall improvement in reproductive performance was also reported on feeding crop residue based complete feed.

### **CONCLUSION**

Higher demand of milk and meat in the country requires newer feed resources with lower input cost for economical livestock rearing. Higher production of food grains i.e. cereals, pulses, oilseeds and agro-industrial products generates larger source of crop residues. Demand and supply gap of roughages and concentrates can be bridged through effective utilization of crop residues. Feed processing technology provides better opportunity for converting poor quality crop residues with proper supplementation into high valued feed. Crop residue based complete feed increases digestibility by increasing rate of delignification, increasing enzymatic hydrolysis of cellulose, hemicelluloses and thus increasing the livestock performance. Strategic use of crop residues for complete feed production can support sustainable livestock production system and can further prevent environmental pollution.

### **REFERENCES**

1. Dikshit AK and Birthal PS. India's livestock feed demand: Estimates and projections, *Agri Eco Res Rev.* 2010;23(1): 15-28.
2. Owen E and Jayasuriya MCN. Use of crop residues as animal feeds in developing countries. *Res Develop Agri.* 1989; 6 (3):129-138.
3. Lailer PC, Dahiya SS, Lal D and Chauhan TR. Complete feed for livestock concept, present status and future trend: A review. *The Ind J Anim Sci.* 2005; 75(1):84-91.
4. Kamusoko R, Jingura R, Parawira W and Sanyika W. Comparison of pretreatment methods that enhance biomethane production from crop residues - a systematic review. *Biofuel R J.* 2019; 6(4): 1080-1089. doi: 10.18331/BRJ2019.6.4.4.
5. Singh M, Bhanotra A, Kujur AST, Singh AK, Wani SA. Complete feed block technology- a fruitful innovation. 2016b [Internet]. [cited 14th February 2020]. Available from: <https://www.biotecharticles.com/Biology-Article/Complete-Feed-Block-Technology-A-Fruitful-Innovation-3604.html>.
6. Li B, Xuezhao S, Qin H, Guiguo Z, Tingting W, Peihua Y, Yuhua H, Wannian T, Rongquan L, Changsheng L, Jianping L, Chunqing W and Baijun S. Pelleting of a Total Mixed Ration Affects Growth Performance of Fattening Lambs. *Frontiers in Veterinary Science*, 2021; volume 8, Article no. 629016. <https://www.frontiersin.org/article/10.3389/fvets.2021.629016>. DOI=10.3389/fvets.2021.629016

7. Afzal Y, Ganai AM, Mattoo FA and Shad FI. Comparative dietary evaluation of oats straw and tree leaf based complete feed in block and mash form in sheep. *Ind J Small Rumin*, 2009;15:212–216.
8. Das MM, Singh KK, Samanta AK, Kundu SS. and Sharma SD. Effect of complete feed blocks on milk yield and its composition in buffalo. In: *Proc. XI Anim Nutr Con*, Jabalpur, 5-7<sup>th</sup> January. 2004;78.
9. Sessaiah CV, Rao SJ, Reddy YR, Nagalakshmi D, Mahender M. and Harikrishna C. Effect of feeding processed sweet sorghum (*Sorghum bicolor* (L) moench) crushed residue based complete ration on growth performance and feeding behavior of murrah buffalo calves. *Vet World*, 2013; 6(3):151-155. doi:10.5455/vetworld.2013.151-155.
10. Reddy R. and Reddy G. Processed crop residues to improve ruminant performance, 2011; <https://www.thebeefsite.com/articles/2821/processed-crop-residues-to-improve-ruminant-performance/>
11. Anandan S, Khan AA, Devulapalli R, Jeethander R. and Blummel M. Comparison of sorghum stover based complete feed blocks with a conventional feeding practice in a peri urban dairy. *Anim Nutr feed technol*, 2010; 10:12-17.
12. Rao KS, Reddy MR. and Reddy GVN. Utilization of sunflower crop residues in the rations of dairy cattle. *Ind J Anim Nutr*, 1999;16 (2): 155-159.
13. Thakur SS. and Tomar SK. On-farm evaluation of a nutritionally balanced total mixed ration on the lactation performance of buffaloes. *Ind J Dairy Sci*, 2007; 60(3): 414-416.
14. Gaafer MA, Hamed MA, Halim A, Mohi ED, Kotob FA, Reidy EL and Bassiouni IM. Productive and reproductive performances of lactating cows and buffaloes fed Total mixed ration. *Philipp. J. Vet. Anim. Sci*, 2011; 37: 131-142.
15. Jadhav SE. and Deshmukh SV. Evaluation of complete rations containing black gram straw and wheat straw in sheep. *Ind J Anim Nutr*, 2001; 18(2): 190-193.
16. Karim A, Shaikh, Mehta BS, Sureshkumar S. and Verma DL. Growth performance and carcass traits of Bharat Merino lambs maintained under intensive feeding and grazing with supplementation. *Ind J Anim Sci*, 2004; 74: 977-979.
17. Bhadane KP, Rekhate DH. and Dhok AP. Nutrient utilization in goats fed Arhar straw based pelleted complete feed. *Ind J Anim Nutr*, 2004; 21: 127–129.
18. Walli TK. Straw based densified complete feed block technology. In [H.P.S. Makkar], ed. *Proceedings of FAO E-Conference on successes and failures with animal nutrition practices and technologies in developing countries*, pp. 41–44, 1– 30 Sept. 2010, FAO, Rome, Italy.