

Short Research Article

Effect on Productivity And Profitability Of Toria through Cluster Frontline Demonstrations (CFLD) In Morigaon District

ABSTRACT

In Morigaon district of Assam, 716 demonstrations on High Yielding Variety (HYV) of toria TS-38 were conducted by Krishi Vigyan Kendra, Morigaon during five consecutive years from 2015-16 to 2019-20. The study revealed that the yield of toria with HYV and improved production technology has been increased significantly as compared to local varieties and farmer's practice. The demo yield was obtained in range from 9.5 to 12 q/ha whereas in farmers practice it was recorded from 5.5 to 8.4 q/ha. The percentage increase over farmer's practice was obtained from 26.67 to 72.73%. the average technology gap, extension gap and technology index were found to be -0.9 q/ha, 3.62 q/ha and -9.00 q/ha respectively. Introducing high yielding varieties with improved production technology, the yield of toria can be enhanced to a great extent with economic conditions of farmers of the community.

Key words: Demonstrations, HYV, Farmer's practice, Improved practice, Toria.

INTRODUCTION

Toria is one of the most important rabi oilseed crop grown in Assam. Being flood affected, huge areas of Morigaon district remains fallow during the Sali season and the farmers mostly prefer toria as a short duration crop before going for cultivation of boro paddy.

The production and productivity of local variety is comparatively low due to lack of irrigation facility, but this variety is still existing due to its distinguish character of growing in rainfed condition. TS -38 is newly developed variety of toria with high yield potential and high oil content as compared to other local varieties of toria available in the district. Realizing the importance of high yielding varieties, cluster frontline demonstrations were conducted in different locations of Morigaon district starting from 2015-16 with improved production technology. The front-line demonstration (FLD) is an important method of transferring the latest package of practices in totality to farmers conducting the FLDs in Morigaon district of

Assam. Further, these demonstrations are designed carefully where provisions are made for speedy dissemination of demonstrated technology among farming community through

organization of other supportive extension activities, such as field days, trainings and group discussion.

MATERIALS AND METHODS

Krishi Vigyan Kendra Morigaon conducted cluster frontline demonstrations of toria in different operational areas of the district, to disseminate the improved production technology and high yielding varieties of toria. From 2015-16 to 2019-20, a total of 716 demonstrations were conducted involving 716 farmers and covering 280 ha area in rabi season. Each demonstration was conducted on an area of 0.4 ha, and an area adjacent to the demonstration plot was kept as farmers' practices. The package of improved technologies like required seed rate, nutrient management, seed treatment and whole packages were used in the demonstrations. Before conducting the demonstrations, training and method demonstrations of seed treatment with PSB and Azotobacter were imparted to the selected farmers. The sowing was done from mid-October to mid-November in broadcasting manner. Data were collected from both the improved and farmers practices and analyzed for cost of cultivation, net income, benefit cost ratio etc. the technology gap, extension gap and technology index were calculated by using the formula as given below

Technology gap = Potential yield (kg/ha) – Demonstration Yield (kg/ha)

Extension gap = Demonstration Yield (kg/ha) – Farmer's yield (kg/ha)

Technology index = {(Potential yield – Demonstration yield)/ Potential yield} x 100

Table 1: Cultivation practices followed for improved practices and farmers practices under

CFLD

Particulars	Improved practices (IP)	Farmers Practice (FP)
Variety	TS-38	Local
Seed Rate	8-10 kg/ha	15-20 kg/ha
Seed treatment	PSB @ 50 gm/kg seed and Azotobacter @ 50 gm/kg seed	Nil
Land preparation	4-5 ploughings (Depending on soil type) to obtain a good tilth	2-3 ploughings

Time of sowing	Mid October to Mid November	October to December
Fertilizer application	Organic cultivation with vermicompost	Imbalanced use of chemical fertilizers
Micronutrient	Borax @ 7.5 kg/ha	Nil
Plant protection	Need based application	Nil

RESULTS AND DISCUSSION

Yield: The average yield of toria variety TS-38 were found 9.5, 12, 11.5, 12 and 9.5 q/ha during the year 2015-16 to 2019-20 respectively, however under farmer's practice the average yield was 7.5, 8.4, 7.5, 7.5 and 5.5 q/ha during respective years. The percentage increase in yield over control was found 26.67, 42.86, 53.33, 60.00 and 72.73 during the year 2015-16 to 2019-20 respectively. The result is in conformity with the findings of Tiwari *et al* (2001), Raj *et al* (2013) and Lydia *et al* (2020).

The Cluster Frontline Demonstrations (CFLD) have given a good impact over the farming community of Morigaon district of Assam. The high yielding variety of toria, TS-38 has enhanced the productivity of toria as compared to other local varieties as well as has increased the income of the farmers. In subsequent years it was observed that farmers are using their own seeds for the next year also, farmer to farmer seed exchange was observed within the village and outside through nearby farmers, relatives and friends. This finding was also in corroboration with the findings of Poonia *et al* (2010) and Lydia Z *et al* (2020).

Table 2. Productivity, technology gap, extension gap and technology index of toria variety TS-38 under CFLD.

Year	Area (ha)	No. of farmers	Seed yield (q/ha)			% increase over control	Tech. Gap (q/ha)	Extension gap (q/ha)	Tech. Index (%)
			Potential	Demo	Control				
2015-16	20	50	10	9.5	7.5	26.67	0.5	2	5
2016-17	30	75	10	12	8.4	42.86	-2	3.6	-20
2017-18	30	75	10	11.5	7.5	53.33	-1.5	4	-15
2018-19	50	125	10	12	7.5	60.00	-2	4.5	-20

2019-20	150	391	10	9.5	5.5	72.73	0.5	4	5
---------	-----	-----	----	-----	-----	-------	-----	---	---

Yield gap: The yield gap is the difference between the farmers potential yield and its current yield. In the present study the yield gap was categorized into technological and extension gap. From the data (Table 2) it was observed that the technology gap ranges from -2 to 0.5 q/ha. The negative technology gap implies that the farmers obtained more yield than its potential yield and it may be attributing due to soil fertility status, timely sowing, weather condition and improved management of the crop. The result indicates the positive impact of introducing high yielding varieties in the locality.

Extension Gap : The extension gap ranges between 2 to 4.5 q/ha during the period of study indicates the lack of knowledge in appropriate time of sowing, soil fertility management, and improved production technology on cultivation of toria. Similar findings were observed by Karate *et al* (2011) in oilseeds and Saikia *et al* (2018) and Jha *et al* (2020) in black gram and Bezbaruah *et al* (2020) in green gram. Need based extension literatures, trainings, workshop and advisories from extension specialists may be required to mitigate the extension gap of the farming community.





Fig 1: Field experiment

Technology index: From the data (Table 2), the technology index ranges from -20 to 5%. The lower value of technology index means the more feasibility of the technology (Jeengar *et al* (2006). From the present study it was revealed that introduction of high yielding toria variety TS-38 has a great impact over the farming community in terms of enhancement in productivity and economic conditions of the farmers in the district.

Economic return: The input and outputs prices of commodities prevailed during the study of demonstrations were taken for calculating the gross return, cost of cultivation, net return and benefit: cost ratio (Table 2). The cost of cultivation, gross return, net return and benefit: cost ratio was calculated considering the input and output costs of the commodities in their respective years. Cultivation of toria under improved technologies gave higher returns of Rs. 28500.00, 45425.00, 45800.00, 43500.00 and 36100.00 per ha during 2015-16 to 2019-20 respectively. The benefit cost ratio was 1.96, 2.67, 1.72, 2.09 and 1.29 from 2015-16 to 2019-20 respectively. This may be due to higher yield obtained under improved technologies compared to farmers practice. The superiority of recommended package of practices under frontline demonstration over farmers' practice was also reported by Mitra *et al*(2010), Balai *et al*(2012), Raj *et al* (2013), Jyothi *et al* (2016) and Chaudhary *et al* (2018).

Table 3. Gross return (Rs/ha), Gross cost (Rs/ha), Net return (RS/ha) and B:C Ratio of HYV toria TS-38 under CFLD.

Year	Area	No of Demo/Farmer	Results		GC	GR	NR	B:C
			Demo (q/ha)	Check (q/ha)				
2015-16	20	50	9.5	7.5	14500	28500	14000	1.96
2016-17	30	75	12	8.4	17013	45425	28412	2.67
2017-18	30	75	11.5	7.5	26600	45800	19200	1.72
2018-19	50	125	12	7.5	20750	43500	22750	2.09
2019-20	150	391	9.5	5.5	27835	36100	8265	1.29

CONCLUSION

From the findings it can be concluded that introduction of high yielding variety and improved production technology can reduce the technology gap resulting in increased productivity of toria as well as higher income of the farmers in the district. Awareness and knowledge on high yielding varieties among farmers can reduce the technology and extension gap of the farmers. Therefore, there is a need to provide proper technical support and guidance from appropriate source for better production and upliftment of the farming community of the district.

The frontline demonstrations on toria conducted at Morigaon district of Assam during 2015-16 to 2019-20 at the farmers' field revealed that the adoption of improved technologies significantly increased the yield of the crop and also the net returns to the farmers. There was 51.11 per cent increase in yield observed in demonstrated plot over farmers plot (Table 2). Lydia Z *et al* (2020) also reported 42.58 cent increased yield in demonstration plot over farmer's practice. TS -38 released by Assam Agricultural University has been recommended in all India level during the year November 2021. There is a need to further disseminate these improved technologies among the farmers with effective extension methods like training, farmers conventions, field days and demonstrations. The farmers should be encouraged to adopt the recommended package of practices for realizing higher returns.

REFERENCES

1. Balai C M, Meena R P, Meena B L and Bairwa R K (2012). Impact of frontline on rapeseed and mustard yield improvement. *Indian Res J Extn Edu* 12 (2): 113-116.
2. Bezbaruah R, Deka R S (2020). Impact of Cluster Frontline Demonstration on Productivity and Profitability of Greengram in Morigaon District of Assam *J Krishi Vigyan*, 9 (1) : 164-169
3. Chaudhary R P, Choudhary G K, Prasad R, Singh R and Chaturvedi A K (2018). Impact Assessment of Front Line Demonstration on Mustard Crop. *Int J Curr Microbiol App Sci Special Issue 7*: 4737-4742
4. Jyothi S V, Mounica D, Pavani U and Dhanu S (2016). Popularization of maize production technology through front line demonstration in tribal areas of East Godavari. *J Krishi Vigyan* 4 (2): 80-8
5. Jeengar K L, Panwar P and Pareek O P (2006). Front line demonstration on maize in Bhilwara district of Rajasthan, *Current Agri* 30 (1/2):115-116

6. Jha AK, Chatterjee K, Mehta BK, Kumari, M Effect of technological interventions of cluster frontline demonstration (CFLDs) on productivity and profitability of black gram (*Vigna Mungo* L.) in Sahibganj district of Jharkhand(2020) . *Int. J of Chemical Studies* 8(5): 2124-2127.
7. Katara S, Pandey SK, Mustaaafa M. Yield gap analysis of rapeseed-mustard through frontline demonstrations. *Agric. Update*. 2011; 6:5-7
8. Lydia Z , Salam G S1 , Nanita D , Singh I M and Prakash N (2020) Impact Assessment of Front Line Demonstration on Toria. *J Krishi Vigyan* 9 (1) : 160-163
9. Mitra B and Samajdar T (2010). Yield gap analysis of rapeseed and mustard through frontline demonstrations. *Agril Ext Rev* 22 (2):16-17.
10. Poonia T C and Pithia M S (2011). Impact of front line demonstrations of chickpea in Gujarat. *Legume Res* 34(4): 304-307.
11. Raj A D, Yadav V and Rathod J H (2013). Demonstrations (FLD) on the Yield of Pulses. *Int J Sci and Res Pub* 3 (9): 1- 4
12. Saikia N, Deb Nath K, Chowdhury P (2018). Impact of cluster frontline demonstrations on popularization of blackgram var. PU 31 in Cachar district of Barak Valley region of Assam. *Journal of Pharmacognosy and Phytochemistry*. 7(4):940-942.
13. Tiwari K B and Saxena A (2001). Economic analysis of FLD of oilseed in Chhindwara. *Bhartya Krishi Anusandhan Patrika* 16 (3&4): 185-189