

Repair of Cerebrospinal Fluid Rhinorrhea in a patient presenting empty sella syndrome: Case Report and Review of Literature

Abstract:

INTRODUCTION: Empty Sella syndrome occurs when intracranial contents herniate through the sellar diaphragm filling the sella turcica with CSF and giving the radiographic appearance of an absent pituitary gland, the association between spontaneous CSF leaks and empty sella is not clear.

CASE PRESENTATION: We report a rare case of a 67 years-old woman who was admitted to our department with a complaint of dripping clear fluid from the left side of her nose. Clinical examination and biological, radiological investigations found a cerebrospinal fluid (CSF) leakage through a minute bony defect in the floor of the sella associated with empty Sella syndrome. Treatment consisted of the reparation of the leak using a fascia lata graft, and the sphenoidal sinus was obliterated with fat.

CONCLUSION: the otolaryngologist should consider empty sella syndrome as a diagnostic possibility when evaluating patients with non-traumatic CSF rhinorrhea

Key Words

Empty Sella syndrome, cerebrospinal fluid leakage, Endonasal endoscopic approach

Introduction

Primary empty sella is characterized by the spontaneous herniation of the subarachnoid space into the pituitary fossa. It is generally asymptomatic and rarely requires any management, however, if symptomatic, the most commonly reported signs and symptoms include headache, cerebrospinal fluid (CSF) leakage, visual deficits, and endocrine dysfunction (1).

But the CSF leakage and visual deficits are the only manifestations for which surgical management should be indicated (1).

The endonasal endoscopic approach is a safe and effective method, offering a better reparation of sinonasal cerebrospinal fluid leak and lowering morbidity (2). This technique provides a success rate greater than 90% (3).

We report the rare case of a patient who presents cerebrospinal fluid rhinorrhea revealing empty sella turcica syndrome.

Case report

A 67-year-old obese woman had a five-month history of dripping clear fluid from the left side of her nose. The fluid was said to be sweet-tasting. There was no associated nasal congestion, and the drainage was worse after bending forward.

The medical history found arterial hypertension, multiparity, history of a headache without head trauma, or meningitis. No endocrine disorder, and no visual disturbances such as diplopia, blind spots, or blurring.

The examination indicated a clear fluid that could be seen flowing from the left side of the nose. The site of origin was high in the nasal cavity. There were no abnormal masses or polyps in the nose or nasopharynx.

The beta-2 transferrin level of the fluid that was dripping from the nose was 18,6 mg/l.

Computerized axial tomography (CT) indicated a decreased density in the sella, the density being similar to the density of CSF rather than that of pituitary tissue. This suggested the possibility of an empty sella Fig 1.




Figure 1: Cerebral CT axial (a) and sagittal view (b): decreased density in the sella, the density is similar to the density of CSF rather than that of pituitary tissue. This suggested the possibility of an empty sella and the presence of a nibbled appearance at the level of the floor of the seller turcique (arrow).

Magnetic resonance imaging (MRI) showed the typical herniation of the subarachnoid space into the sella Fig 2.




Figure 2: Cerebral MRI sagittal (a) and coronal view (b): the presence of intra-sellar fluid formation in continuity with the chiasmatic cistern and identical signal to CSF in all sequences corresponding to an empty sella turcica without mass effect on the pituitary stalk.

The patient underwent a full functional endoscopic sinus surgery exploration and with the use of fluorescein-soaked cottonoids, we find a minute bony defect in the floor of the sella with CSF leaking through this defect. The leak was repaired using fascia lata graft Fig 3, and the sphenoidal sinus was obliterated with fat.




Figure 3: Intraoperative view showing the extraction of a fascia lata graft from the right thigh (a) and its placement in the sphenoid sinus (b).

The CSF leaking stopped after the surgery, but the patient developed a persistent headache.

The patient was treated with oral acetazolamide and a cough suppressant and a laxative, and her symptoms improved without a recurrence of CSF leaking. No recurrence was present at 6 months of follow-up.

Discussion

Empty Sella is defined as a hernia of subarachnoid space in the sella turcica also called arachnoidocele. It is a term used in radiology to describe the discovery of "empty sella space" on magnetic resonance imaging and computed tomography with a flattened pituitary gland and an elongated stalk. It can be complete if the cerebrospinal fluid fills more than 50% of space in the sella and the thickness of the gland is less than 2 mm or partial if less than 50% of the sellar space is filled with CSF (4).

Regarding its etiology, it can be primary if there is no pathological mechanism in the sellar region preceding the pituitary lesion or secondary if it is consecutive to a particular pathological mechanism (4).

Primary Empty Sella is the less common of the two entities. Currently, its cause is not fully understood, but researchers have proposed several mechanisms including chronic intracranial hypertension, incompetence or complete absence of the diaphragm sellae, and temporary expansion followed by regression of the pituitary gland (5).

The primary syndrome is commonly seen in middle-aged, obese, and hypertensive women. The disorder may be a sign of idiopathic intracranial hypertension. Among the risk factors, there is obesity in women and multiparity (6). This may be the case with our patient

The clinical presentation of primary empty sella is silent and asymptomatic, not requiring any management. In very rare cases, patients develop cerebrospinal fluid leaks or visual deficits that require surgical treatment (1).

The etiology of CSF leaks is diverse: post-surgical iatrogenic (endoscopic sinus surgery is a common cause of iatrogenic CSF leak), trauma is also a common cause, followed by congenital meningoencephalocele, meningioma/pituitary tumor (7).

While the etiology of traumatic CSF leaks is well known, spontaneous leaks are not and they are also important causes of CSF fistulas, associated with high intracranial pressure, the causes of which are partially empty sella turcica, empty sella turcica, base cranial erosions, hydrocephalus, and intracranial arteriovenous malformation (2).

Our patient had signs of an empty sella on MRI, responsible for raised ICP. Full functional endoscopic sinus surgery exploration uncovered a tiny bone defect in the floor of the sella with CSF leaking through this defect

The association between the radiographic finding of an empty sella and the clinical presentation of spontaneous CSF leak is much more common than would be expected (8).

The study of Schlosser. and al (8) and the study of Shetty and al (9) have noted the association between spontaneous CSF leaks and an empty sella. Shetty et al¹⁶ reviewed 11 patients with spontaneous sphenoid CSF leaks and noted that 63% had completely empty sellae and an additional 27% had partially empty sellae, similar to Schlosser's study results of 67% and 33%, respectively (8,9).

The indications for endoscopic closure are the accessibility of the leak endoscopically such as localization at (Sphenoid, Cribriform plate, Frontal sinus excepted for posterior wall, Fovea ethmoidal/Lateral lamina, Septal olfactory dural extension) after confirming it by using immunofixation of beta 2–transferring (2).

The steps of the endoscopic surgical technique require intraoperative localization of the fluorescein leak and preoperatively by magnetic resonance imaging and computed tomography (CT) (2).

Various grafts including muscle, fascias (lata, temporalis, and rectus abdominis), bone, mucoperiosteum, mucochondrium, dural flaps, fat, and various synthetic dural substitutes may be positioned with different techniques (onlay and/or underlay, dumbbell, bath plug, and fat obliteration) to provide a stable scaffold at the site of the defect, with almost identical postoperative success rates. Free grafts (fascia lata) are advantageous by providing an easily manageable mass allowing a satisfactory field of intraoperative vision with decreased tissue tension (10). In our patient, we have used a fascia lata graft.

Various materials are used to support the secure position of a graft. Fat is a useful supplement and can act as glue to hold the fascia in place and generally render fibrin glue unnecessary. 7 In addition, free fat is the preferred material for obliterating the sphenoid sinus and mounting the graft on the sinus wall (10), like our patient. Fibrin glue can be used as an adjuvant to improve graft adhesion and secure the graft in the correct position (10).

Among causes of recurrent sinonasal cerebrospinal fluid (CSF) leaks after repair, raised intracranial pressure is the most common reason (2).

Conclusion

These cases of non-traumatic CSF rhinorrhea present a diagnostic challenge.

This case demonstrates that the otolaryngologist should consider empty sella syndrome as a diagnostic possibility when evaluating patients with non-traumatic CSF rhinorrhea.

Endoscopic closure is now the treatment of choice in the majority of patients with sinonasal cerebrospinal fluid (CSF) leaks.

UNDER PEER REVIEW

Ethical approval

I certify that this kind of manuscript does not require ethical approval by the Ethical Committee of our institution.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

Registration of research studies

This is a case report that does not require a research registry.

REFERENCES

1. Guinto G, del Valle R, Nishimura E, Mercado M, Nettel B, Salazar F. Primary empty sella syndrome: the role of visual system herniation. *Surg Neurol.* 1 juill 2002;58(1):42-7.
2. Mirza S, Thaper A, McClelland L, Jones NS. Sinonasal cerebrospinal fluid leaks: management of 97 patients over 10 years. *The Laryngoscope.* 2005;115(10):1774-7.
3. Chang C. Septal cartilage plug technique in spontaneous cerebrospinal fluid rhinorrhea postoperatively diagnosed with partial empty sella syndrome. *J Craniofac Surg.* 2014;25(4):1408-9.
4. Miljic D, Pekic S, Popovic V. Empty Sella. In: Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K, et al., éditeurs. *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000 [cité 4 janv 2022]. Disponible sur: <http://www.ncbi.nlm.nih.gov/books/NBK532084/>

5. Ucciferro P, Anastasopoulou C. Empty Sella. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 [cité 4 janv 2022]. Disponible sur: <http://www.ncbi.nlm.nih.gov/books/NBK541002/>
6. Izizag BB, Ngandu A, Mbiso DL. Syndrome de la selle turcique vide: à propos d'un cas. *Pan Afr Med J.* 21 août 2019;33:317.
7. Hegazy HM, Carrau RL, Snyderman CH, Kassam A, Zweig J. Transnasal endoscopic repair of cerebrospinal fluid rhinorrhea: a meta-analysis. *The Laryngoscope.* 2000;110(7):1166-72.
8. Schlosser RJ, Bolger WE. Significance of Empty Sella in Cerebrospinal Fluid Leaks. *Otolaryngol Neck Surg.* 1 janv 2003;128(1):32-8.
9. Shetty PG, Shroff MM, Fatterpekar GM, Sahani DV, Kirtane MV. A retrospective analysis of spontaneous sphenoid sinus fistula: MR and CT findings. *Am J Neuroradiol.* 2000;21(2):337-42.
10. Ozturk O, Polat S, Uneri C. Endoscopic endonasal management of cerebrospinal fluid rhinorrhea. *J Craniofac Surg.* juill 2012;23(4):1087-92.