

Case report: Recurrent Calf Muscle Myositis in Pediatric Patient

Abstract

Focal myositis is a rare entity characterized by acute onset and localized skeletal muscle inflammation.

A four-year-old boy presented with recurrent muscle myositis. He had a high level of muscle enzymes.

Discharge after receiving fluid and analgesia.

Keywords: Recurrent, Muscle, Myositis.

Introduction

Heffner (1) described focal myositis as a rare entity characterized by acute onset and localized skeletal muscle inflammation of unknown origin. It's restricted to one muscle with a benign course without sequelae or recurrence (2).

This case report aims to describe a child affected with recurrent focal myositis. Furthermore, we reviewed the literature to summarize clinical, laboratory, management, and outcome data on recurrent focal myositis in childhood.

Case report:

First visit to ED:

A four-year-old boy presented to the emergency department (ED) with fever, cough, and runny nose for four days, one-day history of calf muscle pain, and an inability to walk. No headache, no vomiting, and no joint pain or swelling. He is medically free, but all families had upper respiratory tract infection symptoms.

On examination: normal vital signs, calf muscles were tender, hot, no calf muscle swelling, normal joint and skin examination, normal cardiovascular, chest, abdomen, and neurological examination.

Other systemic exams were normal.

Investigations: positive COVID-19 PCR Test, CK-Total was 11135 U/L[39-308 U/L], LDH 686 U/L[120-300 U/L], Aldolase 65.3 U/L[<7.7 U/L], normal CRP and ESR, normal urine for myoglobinuria, Normal Complete blood count, normal serum electrolytes and renal function, normal urine analysis and culture, PCR for MRSA not detected, and normal ECG.

Treatment: fluid for hydration, analgesia, and bed rest. After four days, he was discharged in good condition with normal CK-Total.

Second visit to ED:

After two months after the first attack patient presented to ED with fever, symptoms of the upper respiratory tract infection for two days, and an inability to walk with calf muscle pain for one day. No headache, vomiting, diarrhea, or joint pain.

Examination: normal vital signs, tender calf muscles, normal joint and skin. Normal other physical examination.

Investigations: negative COVID-19 PCR Test, CK-Total was 1197 U/L [39-308 U/L], normal complete blood counts, normal serum electrolytes and renal function, normal urine analysis and culture, no myoglobinuria, normal CRP and ESR.

Treatment: good hydration with fluid and analgesia with bed rest.

Discussion

Muscle myositis is an episode of a muscle mass enlargement characterized by recurrent focal myositis over days to months(1.,3,4). Most published reports are single cases or case series involving adults in their studies (5). Then, relapsing episodes of focal myositis account for up to 18% of focal myositis cases. (2)

Focal myositis is a focal inflammatory pseudo tumor usually restricted to one skeletal muscle with a benign course, prognosis, and self-remission. The nature of recurrence seems to be related to the etiology of the illness (2, 6, 7, 8, 9). Viruses and infection were the leading causes of focal myositis, but no specific organism was identified. Toti(10) tried PCR looking for EBV, herpes simplex virus types 1 and 2, and cytomegalovirus but had negative results.

Laboratory findings in patients with focal myositis showed both normal or slightly elevated inflammation indices (white blood count, CRP, ESR) and muscle enzymes (CPK, aldolase, and LDH)(11).

Lorenzo Schantz et al. reported 13 year old with recurrent high muscle enzymes with pain diagnosed with recurrent myositis treated with conservative management (12).

Conclusion

Although focal myositis is considered a self-limited disease, we believe a long-term clinical and biochemical follow-up is needed.

References

- 1-Heffner RR., Armbrustmacher VW., Earle KM. Focal myositis. *Cancer* 1977;40(1):301–6.
- 2-Yanmaz Alnigenis MN., Kolasinski SL., Kalovidouris AE. Focal myositis: a review of 100 previously published cases and a report of 2 new cases. *Clin Exp Rheumatol* 17(5):631.
- 3-Manganelli S., De Stefano R., Malandrini A., Selvi E., Frati E., Gambelli S., et al. Bilateral recurrent focal myositis of gastrocnemius muscles after BCG vaccination. *Rheumatology (Oxford)* 2002;41(9):1074–6.
- 4-Auerbach A., Fanburg-Smith JC., Wang G., Rushing EJ. Focal myositis: a clinicopathologic study of 115 cases of an intramuscular mass-like reactive process. *Am J Surg Pathol* 2009;33(7):1016–24
- 5- Teixeira F., Peixoto D., Costa JA., Bogas M., Taipa R., Melo Pires M., et al. Recurrent Focal myositis: a rare inflammatory myopathy. *Acta Reum Port* 39(2):172–5.
- 6- Smith AG, Urbanits S, Blaivas M, Grisold W, Russel JW. Clinical and pathologic features of focal myositis. *Muscle Nerve* 2000;23:1569-75.
- 7- Naughton M, Jessop JD, Williams BD. Idiopathic recurrent non-suppurative focal myositis: a report of two cases. *Br J Rheumatol* 1994;32:1101-1104.
- 8- Garcia-Consuegra J, Morales C, Gonzales J, Merino R. Relapsing focal myositis: a case report. *Clin Exp Rheumatol* 1995;13:395-397.
- 9- Sieb JP, Ries F, Träber F, Kaminski M. Recurrent focal myositis. *Muscle Nerve* 1997;20:1205-1206. ses of an intramuscular mass-like reactive process. *Am J Surg Pathol* 2009;33:1016-24.
- 10- Toti P, Romano L, Villanova M, Zazzi M, Luzi P. Focal myositis: a polymerase chain reaction analysis for a viral etiology. *Hum Pathol.* 1997;28(1):111–113. [[PubMed](#)] [[Google Scholar](#)]

11- Galloway HR, Dahlstrom JE, Bennett GM. Focal myositis. *Australas Radiol.* 2001;45(3):347–349.

12-Corradin M, Alaggio R. Muscles **Focal Myositis in paediatric age.** *Ligaments Tendons J*, 5(1):45-50, 01 Jan 2015

UNDER PEER REVIEW