

Elemental Concentration and Potential Ecological Risk Assessment of Mangrove Surface Sediments in Conakry, Guinea

ABSTRACT

Surface sediments of Conakry mangrove, Guinea were sampled and analyzed to assess their concentration and potential ecological risks. Seventeen trace elements and five major elements were analyzed using XRF. Mean, STDEV, correlation matrix, CF, C_d , EF, PI, MPI, Er^1 , RI and PCI were used for data analysis. The results indicated contrasting element concentrations with location. Cl; Fe_2O_3 and Sc; MnO had the highest and lowest average concentrations among the trace and major elements used for the study. Relative to the UCC values, As, Pb, Zn, Ni, Cr, V, Cl, Fe_2O_3 , MnO and P_2O_5 had higher values while Cu, Sr, Y, Nb, Zr, Th, Sc, F, TiO_2 and CaO had lower values. Fe_2O_3 majorly influenced the concentration of most of the elements. As enrichment had a potentially moderate to considerable ecological risk while Cr, Cu, Pb, Zn and Ni were found to have potentially low ecological risk.

Keywords: Mangrove sediment. Trace element. Major element. Sediment pollution. Element enrichment. Ecological risk.

1. INTRODUCTION

Mangroves are specialized plants that grow in intertidal zones. They are halophytic and have tangled root system. The intricate nature of mangrove root system increases the deposition rate of marine sediment load carried by either tides or waves as well as the upland sediment load carried by runoff. Thus, the mangrove environment is associated with high rate of deposition. Also, sediments deposited in mangrove environment are fluvial in nature and gradually pile up over time. They are often laden with contaminants from diverse sources [1]. Some of these non point sources of contaminants may be industrial, agricultural or even geogenic. Notwithstanding these non point sources of contaminants, the magnitude of heavy metal contamination depends not only on their sources but also on the solid and aqueous phases of physicochemical properties, microbial activities as well as redox conditions [2]. However, among the organic and inorganic contaminants within the mangrove environment, heavy metals account for the considerable sources of poor ecological quality.

Therefore, assessment of elemental concentration in mangrove surface sediments can potentially provide in-depth information on the sediment quality as well as the ecological status of the mangrove environment. Given that mangroves are described as sink for heavy metals [3], it holds vital history of heavy metal concentration which might be either geogenic and/or anthropogenic [1]. There are limited studies on heavy metals in

mangrove surface sediments in Guinea [4] and these studies analyzed only few metals.

It is against this backdrop that this study was undertaken to assess elaborate metal concentrations in Conakry mangrove sediments as well as their probable ecological risk. As such, this study seeks to deepen insight into (a) trace and major element concentration in mangrove surface sediments in Conakry, (b) evaluation of sediment contamination and (c) determination of the ecological risks of metal concentration in mangrove sediment.

2. MATERIALS AND METHODS

2.1. Study Area

Guinea has a coastal stretch of 1,614.5 km and a mangrove area of 2,039 km² [5]. Its mangroves are described as the best developed in West Africa with stands exceeding 25 meters in height and extends up to 160 km inland [6]. The span of mangroves is influenced remarkably by tides. Conakry is situated on an isthmus. It extends as far as the Peninsula of Kaloum and bordered by mangrove swamps. It lies between 9°30' to 9°40' to the north and 13°30' to 13°42' to the west. Conakry receives about six months of rainfall and up to 4,000 mm per annum [7]. Temperature ranges annually are between 23°C and 29°C [8]. Geologically, Conakry is made up of Jurassic mafic to ultramafic rocks of Kakoulima laccolith [9]. However, according to [6, 10] other areas beyond Conakry consist of granitic rocks of precambrian age.




Figure 1. Map of the study area showing mangrove distribution.

2.2. Mangrove Surface Sediment Sampling and Preparation

The surface sediment samples from Conakry mangroves were obtained from seven locations using the bucket auger. Sampled sediments weighing up to 200g were packaged in plastic bags in the field and stored at 4°C in a cooler. To reduce weight,

the homogenized samples were air dried for 48 hours before repackaging them in zip-lock bags and placed in plastic boxes. The samples were exported and analyzed in Geoscience Laboratory, Shimane University, Japan. Sediment samples weighing about 50 g each were oven dried at 160°C for 2 days. Using the automatic agate mortar and pestle grinder, the oven dried samples were made into powder after 20 minutes of grinding. After which about 5g of each of the powdered sediment samples were compressed into briquettes using a force of 200 kN for 60s.

2.3. Laboratory Analysis

Seventeen trace elements; As, Pb, Zn, Cu, Ni, Cr, V, Sr, Y, Nb, Zr, Th, Sc, TS, F, Br and Cl as well as five major elements; TiO₂, Fe₂O₃, MnO, CaO and P₂O₅ were analyzed using X-ray fluorescence (XRF) RIX-200 spectrometer. In accordance with [11], using pressed powder briquettes with average errors being <10% XRF analysis were done.

2.4. Statistical Analysis

The mean and standard deviations of the trace and major element concentrations in the mangrove surface sediment samples and correlation matrix to show the inter-element relationships in the study area were computed using Microsoft Excel 2013. As-Fe₂O₃, Pb-Fe₂O₃, Zn-Fe₂O₃ and Cu-Fe₂O₃ correlation graphs were as well as normalized concentration graph using upper continental crust (UCC) were done using Kaleida Graph 4.0.

2.5. Sediment Pollution Assessment (SPA)

2.5.1. Contamination Factor (CF)

To establish the extent of heavy metal contamination in the mangrove surface sediments of Conakry, the contamination factor was employed. Contamination factor is expressed in line with [12] as:

$$CF = C_{\text{metal}} / C_{\text{background}}$$

Where:

C_{metal} refers to the current metal concentration within the sediments.

$C_{\text{background}}$ refers to the background metal concentration of sediments.

The upper continental crust proposed by [13] was used as the background metal concentration. The CF is interpreted thus: CF < 1: signifies low contamination; 1 ≤ CF < 3: signifies moderate contamination; 3 = CF ≤ 6: signifies considerable contamination and CF ≥ 6: signifies very high contamination.

2.5.2. Degree of Contamination (C_d)

To further enhance pollution determination in sediments, [14] suggested a sedimentological diagnostic tool known as degree of contamination which is defined thus:

$$C_d = \sum CF$$

It is used to calculate the degree of contamination of an element by summing up the CF of a given element in the samples. C_d is interpreted as: $C_d < 6$ shows low C_d ; $6 < C_d < 12$ means moderate C_d ; $12 < C_d < 24$ is considerable C_d while $C_d > 24$ is high C_d and shows strong anthropogenic pollution.

2.5.3. Enrichment Factor (EF)

The determination of the anthropogenic impact of elemental enrichment in the surface sediments of Conakry mangrove was done using the enrichment factor. The EF uses a normalized approach where normalizing elements could be any of Al, Fe and Si [15] as well as Li, Sc, Ti and Zr [16]. This study used Fe as the normalizing element for the computation of the enrichment factor. The anthropogenic sources of Fe are insignificant, thus it was chosen as the element for normalization [17]. Thus, Fe is a conservative tracer and could be used to differentiate between natural and anthropogenic components in sediments [18]. EF is given as:

$$EF = (X_s/Fe_s)_{\text{sediment}} / (X_b/Fe_b)_{\text{background}}$$

Where: X_s refers to the element being considered from the sediment sample and Fe_s refers to the normalizer from an equivalent sample.

X_b is the element being considered from the background while Fe_b is the normalizer from the background.

The upper continental crust values proposed by [13] were used as the background concentration. The interpreted of EF given thus: EF < 2 is depletion to minimal enrichment and pollution; EF 2 - 5 is moderate enrichment and pollution; EF 5 - 20 is significant enrichment and pollution; EF 20

- 40 is very high enrichment and pollution while EF > 40 is extreme enrichment and pollution.

2.5.4. Pollution Index (PI)

The pollution index is a sediment pollution assessment tool developed by [19]. It makes use of the average and maximum contamination factor values of a metal in its computation. It is interpreted as; PI < 0.7 (unpolluted), 0.7 < PI < 1 (slightly polluted), 1 < PI < 2 (moderately polluted), 2 < PI < 3 (severely polluted) and PI > 3 (heavily polluted). The PI is expressed as:

$$PI = \sqrt{\frac{(Cf_{\text{average}})^2 + (Cf_{\text{max}})^2}{2}}$$

Where: the average contamination factor value of a metal is its Cf_{average} while the maximum contamination value of same metal is its Cf_{max} .

2.5.5. Modified Pollution Index (MPI)

The modified pollution index is a sediment pollution assessment tool developed by [20]. Instead of using contamination factor values of metal in its computation, the modified the pollution index uses the average and maximum enrichment factors. The interpretation is as follows; MPI < 1 (unpolluted), 1 < MPI < 2 (slightly polluted), 2 < MPI < 3 (moderately polluted), 3 < MPI < 5 (moderately-heavily polluted), 5 < MPI < 10 (severely polluted) and MPI > 10 (heavily polluted). The PI is expressed as:

$$MPI = \sqrt{\frac{(Ef_{average})^2 + (Ef_{max})^2}{2}}$$

Where: $Ef_{average}$ is average enrichment factor value of a metal while Ef_{max} is the maximum enrichment factor value of same metal.

2.6. Potential Ecological Risk Assessment (PERA)

2.6.1. Potential Contamination Index (PCI)

The potential contamination index was proposed by [21] and it is given as:

$$PCI = (\text{Metal}_{\text{sample maximum}}) / (\text{Metal}_{\text{background}})$$

Where: $\text{metal}_{\text{sample maximum}}$ refers to the maximum concentration of a metal in sediment while $\text{metal}_{\text{background}}$ refers to the background value of the same metal. The upper continental values of [13] were used as background values. It is interpreted thus: $PCI < 1$ is low contamination, $1 < PCI < 3$ is moderate contamination and $PCI > 3$ is severe or very severe contamination.

2.6.2. Potential Ecological Risk Factor (Er^i)

To determine the potential ecological risk of an element in the sediments, the potential ecological risk factor put forth by [14] was used. It is calculated as follows:

$$Er^i = Tr^i \times C_f^i$$

Where: Er^i is the potential ecological risk factor of element (i) while Tr^i is the toxic response factor of same element (i). C_f^i refers to the contamination factor of element (i). The toxic response factors of As, Cr, Cu, Pb, Zn and Ni were given as 10, 2, 5, 5, 1 and 5 respectively [14]. The Er^i values are interpreted as follows: $Er^i < 40$ is low potential ecological risk, $40 \leq Er^i < 80$ is moderate potential ecological risk, $80 \leq Er^i < 160$ is considerable potential ecological risk, $160 \leq Er^i < 320$ is high potential ecological risk and $Er^i \geq 320$ is very high potential ecological risk.

2.6.3. Potential Ecological Risk Index (RI)

The ecological index is a tool used to estimate the ecological risk posed by an element in the sediment. According to [14], it is calculated by summing up the Er^i and is interpreted as follows: $RI < 150$ is low ecological risk, $150 \leq RI < 300$ is moderate ecological risk, $300 \leq RI < 600$ is considerable ecological risk and $RI \geq 600$ is very high ecological risk. The formula for RI is given thus:

$$RI = \sum Er^i$$

3. RESULTS AND DISCUSSION

3.1. Element Concentration in Mangrove Surface Sediments of Conakry

Trace and major element concentrations in the mangrove surface sediments of Conakry are presented on Tables 1 and 2 respectively. Also, the tables presented the comparison of the trace and major element concentrations with the UCC by [13].

Table 1. Trace element concentrations (ppm) in Conakry mangrove surface sediment

| | As | Pb | Zn | Cu | Ni | Cr | V | Sr | Y | Nb | Zr | Th | Sc | TS | F | Br | Cl |
|---------|-------|-------|--------|------|-------|--------|--------|-------|------|------|--------|------|------|---------|--------|------|---------|
| CKRY1 | 26 | 31 | 152 | 7 | 63 | 736 | 190 | 28 | 8 | 7 | 120 | 6 | 4 | 1172 | 47 | 9 | 6570 |
| CKRY2 | 24 | 30 | 148 | 3 | 62 | 698 | 171 | 30 | 8 | 7 | 110 | 5 | 4 | 1038 | 60 | 8 | 5004 |
| CKRY3 | 24 | 31 | 154 | 5 | 74 | 750 | 176 | 41 | 8 | 7 | 94 | 5 | 5 | 1175 | 230 | 10 | 6393 |
| CKRY4 | 23 | 30 | 142 | 5 | 66 | 704 | 154 | 33 | 8 | 7 | 91 | 5 | 4 | 1295 | - | 12 | 9232 |
| CKRY5 | 25 | 29 | 139 | 5 | 58 | 736 | 169 | 31 | 8 | 7 | 166 | 5 | 4 | 1150 | 74 | 9 | 7221 |
| CKRY6 | 54 | 84 | 162 | 9 | 83 | 1122 | 319 | 27 | 9 | 7 | 139 | 5 | 5 | 1128 | 129 | 6 | 8615 |
| CKRY7 | 46 | 72 | 163 | 7 | 84 | 899 | 279 | 34 | 9 | 7 | 128 | 7 | 6 | 986 | 126 | 6 | 5657 |
| Average | 31.71 | 43.86 | 151.43 | 5.86 | 70.00 | 806.43 | 208.29 | 32.00 | 8.29 | 7.00 | 121.14 | 5.43 | 4.57 | 1134.86 | 111.00 | 8.57 | 6956.00 |
| STDEV | 12.74 | 23.59 | 9.20 | 1.95 | 10.44 | 154.65 | 63.92 | 4.69 | 0.49 | 0.00 | 26.30 | 0.79 | 0.79 | 100.38 | 67.49 | 2.15 | 1525.69 |
| UCC | 4.4 | 18 | 70 | 32 | 38 | 80 | 98 | 266 | 17.4 | 12 | 188 | 8.95 | 15 | - | 561 | - | 142 |

Table 2. Major element concentration (ppm) in Conakry Mangrove Surface Sediment

| | TiO ₂ | Fe ₂ O ₃ | MnO | CaO | P ₂ O ₅ |
|---------|------------------|--------------------------------|------|------|-------------------------------|
| CKRY1 | 0.42 | 17.67 | 0.17 | 0.66 | 0.20 |
| CKRY2 | 0.38 | 16.22 | 0.15 | 0.68 | 0.19 |
| CKRY3 | 0.39 | 16.38 | 0.15 | 0.86 | 0.21 |
| CKRY4 | 0.36 | 15.54 | 0.15 | 0.69 | 0.19 |
| CKRY5 | 0.43 | 16.15 | 0.16 | 0.73 | 0.20 |
| CKRY6 | 0.52 | 24.2 | 0.17 | 0.68 | 0.37 |
| CKRY7 | 0.49 | 22.53 | 0.18 | 0.84 | 0.32 |
| Average | 0.43 | 18.38 | 0.16 | 0.73 | 0.24 |
| STDEV | 0.06 | 3.50 | 0.01 | 0.08 | 0.07 |
| UCC | 0.67 | 5.33 | 0.10 | 3.44 | 0.16 |

The trace and major element concentrations obtained from the mangrove surface sediments of Conakry after XRF analysis showed wide variations among elements and at different locations. The elements are concentrated thus; Cl>TS>Cr>V>Zn>Zr>F>Ni>Pb>Sr>As>Br>Y >Nb>Cu>Th>Sc for the trace elements and

Fe₂O₃>CaO>TiO₂>P₂O₅>MnO for the major elements. In comparison to the UCC values by [13], it was found that As, Pb, Zn, Ni, Cr, V and Cl had higher average concentration values while Cu, Sr, Y, Nb, Zr, Th, Sc and F had lower average values. Among the major elements, Fe₂O₃, MnO and P₂O₅ had higher

concentrations while TiO_2 and CaO relative to the UCC values.




Figure 2. Comparison of trace and major element concentration in surface sediments of Conakry mangrove. All values were normalized to the UCC values of [13].

3.2. Comparison of Metal Concentrations in Guinea and Nigeria Mangrove Surface Sediments.

Guinea and Nigeria are both West African countries and thus have lots of climatic similarities. Guinea has the best developed mangroves in West Africa with stands exceeding 25 meters [6] while mangrove concentration in Nigeria is the largest not only in West Africa but also in Africa with stands of up to 40 meters [5]. Both Guinea and Nigeria mangroves are largely influenced by tides and are in the Atlantic East Pacific (AEP) mangrove group.

The graphical comparison of element concentrations in Guinea and Nigeria mangrove surface sediments was based on the data generated by this study and Niger delta mangrove surface sediment data from [22] is shown in Figure 3. Twenty elements were compared and it was found that the Guinea mangrove surface sediments had higher average concentrations in As, Pb, Zn, Ni, Cr, V, F, Cl, Fe_2O_3 , MnO, CaO and P_2O_5 . However, Nigeria mangrove surface sediments had higher average element concentrations in Cu, Sr, Y, Nb, Zr, Th, Sc and TiO_2 . Again, both mangrove areas have similar trend in element concentrations.




Figure 3. Comparison of trace and major element concentration in surface sediments of Guinea and Nigeria mangrove. All values were normalized to the UCC values of [13].

3.3. Inter-element Relationship

The correlation matrix of element concentration in the mangrove surface sediments of Conakry was presented in Table 3. Sources of elements are defined using Fe_2O_3 and TiO_2 as proxies [23, 24]. According to [25], elements that show strong correlations with TiO_2 reflect only natural detrital inputs. This is because in sediments, lithogenic elements usually have a linear relationship with TiO_2 . As such, where no correlation exists between TiO_2 and any given element, it then means that either anthropogenic or additional natural processes may have contributed to elemental enrichment [26].

Based on the analysis of mangrove surface sediment samples obtained from Conakry, there is a strong TiO_2 -As-Pb-Cu-Cr-V, moderate TiO_2 -Zn-Ni-Zr-Sc, weak TiO_2 -Th-F-Cl and negative TiO_2 -Sr-TS-Br associations. This implies that As, Pb, Cu, Cr and V concentration in the sediments are significantly influenced by TiO_2 . TiO_2 has mild influence on the concentration of Zn, Ni, Zr

and Sc. However, TiO_2 has no influence on the concentration of Sr, TS and Br in the sediments. Fe_2O_3 associated strongly with As, Pb, Zn, Cu, Ni, Cr, V and TiO_2 . The Fe_2O_3 -Sc is moderate while Fe_2O_3 -Zr-Th-F-Cl is weak. However, Fe_2O_3 is negatively associated with Sr, TS and Br. This indicates that the concentration of As, Pb, Zn, Cu, Ni, Cr, V, Y and TiO_2 are strongly influenced by Fe_2O_3 . Similarly, Fe_2O_3 has no influence on the concentration of Sr, TS and Br. In general, it could be concluded that Fe_2O_3 is the major geogenic determiner of element concentration in the mangrove surface sediments of Conakry. Also, the concentration of As, Pb, Zn, Cu, Ni, Cr, V and TiO_2 are largely lithogenic while the concentration of Sr, TS and Br are not.

Trace elements that are strongly and positively correlated have a common source and similar enrichment process [27]. Thus, the resultant suites of As-Pb-Zn-Cu-Ni-Cr-V in the surface sediments implies that these elements have a common source and enrichment process. Pb, Zn, Cu and P_2O_5 concentration in sediments are often influenced by anthropogenic factors [28].

This is because they are mostly used in both technological and agricultural activities [29]. As such, the strong positive P_2O_5 -As-Pb-Zn-Cu-Ni-Cr-V relationship implies likely enrichment from a common anthropogenic source such as fertilizer application in surrounding upland areas. Also in sediments, Cu:Zn ratio can be employed in the description of redox conditions [30]. According to [31], an increase in the Cu:Zn ratio indicates a shift towards anaerobic conditions in sediments as a result of increased organic production and marks the

beginning of eutrophication. This study found an increase in the Cu:Zn ratios in all the sampling locations. Specifically, the sampling locations CKRY1, CKRY2, CKRY3, CKRY4, CKRY5, CKRY6 and CKRY7 had Cu:Zn ratios of 1:21.71, 1:49.33, 1:30.80, 1:28.40, 1:27.80, 1:18.00 and 1:23.29 respectively. This shows that the sediments were at varying anoxic conditions as at the time of sampling. The As- Fe_2O_3 , Pb- Fe_2O_3 , Zn- Fe_2O_3 and Cu- Fe_2O_3 diagrams in Fig. 4 express the element correlations with Fe_2O_3 in Conakry mangrove surface sediments.

Table 3. Correlation matrix of trace and major elements in surface sediments of the Conakry mangrove.

| | As | Pb | Zn | Cu | Ni | Cr | V | Sr | Zr | Th | Sc | TS | F | Br | Cl | TiO ₂ | Fe ₂ O ₃ | MnO | CaO | P ₂ O ₅ | |
|--------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------|-------------|-------|-------|-------|-------|-------|------------------|--------------------------------|------|------|-------------------------------|--|
| As | 1 | | | | | | | | | | | | | | | | | | | | |
| Pb | 1.00 | 1 | | | | | | | | | | | | | | | | | | | |
| Zn | 0.81 | 0.82 | 1 | | | | | | | | | | | | | | | | | | |
| Cu | 0.82 | 0.79 | 0.68 | 1 | | | | | | | | | | | | | | | | | |
| Ni | 0.85 | 0.88 | 0.90 | 0.67 | 1 | | | | | | | | | | | | | | | | |
| Cr | 0.96 | 0.95 | 0.75 | 0.84 | 0.80 | 1 | | | | | | | | | | | | | | | |
| V | 0.99 | 0.99 | 0.87 | 0.81 | 0.87 | 0.94 | 1 | | | | | | | | | | | | | | |
| Sr | -0.32 | -0.27 | 0.00 | -0.36 | 0.17 | -0.33 | -0.28 | 1 | | | | | | | | | | | | | |
| Zr | 0.38 | 0.32 | -0.03 | 0.34 | -0.06 | 0.37 | 0.35 | -0.50 | 1 | | | | | | | | | | | | |
| Th | 0.38 | 0.39 | 0.55 | 0.37 | 0.43 | 0.16 | 0.48 | 0.00 | 0.10 | 1 | | | | | | | | | | | |
| Sc | 0.72 | 0.75 | 0.84 | 0.50 | 0.91 | 0.61 | 0.77 | 0.36 | 0.04 | 0.62 | 1 | | | | | | | | | | |
| TS | -0.46 | -0.46 | -0.55 | -0.05 | -0.37 | -0.30 | -0.53 | 0.10 | -0.34 | -0.55 | -0.56 | 1 | | | | | | | | | |
| F | 0.16 | 0.20 | 0.42 | 0.11 | 0.58 | 0.24 | 0.17 | 0.81 | -0.46 | -0.12 | 0.61 | 0.17 | 1 | | | | | | | | |
| Br | -0.83 | -0.81 | -0.75 | -0.53 | -0.60 | -0.74 | -0.86 | 0.38 | -0.53 | -0.46 | -0.62 | 0.84 | 0.13 | 1 | | | | | | | |
| Cl | 0.16 | 0.16 | -0.20 | 0.40 | 0.08 | 0.30 | 0.06 | -0.20 | 0.01 | -0.41 | -0.21 | 0.76 | 0.14 | 0.36 | 1 | | | | | | |
| TiO ₂ | 0.94 | 0.91 | 0.73 | 0.84 | 0.71 | 0.91 | 0.94 | -0.37 | 0.62 | 0.43 | 0.65 | -0.51 | 0.05 | -0.87 | 0.07 | 1 | | | | | |
| Fe ₂ O ₃ | 0.99 | 0.99 | 0.86 | 0.84 | 0.85 | 0.95 | 1.00 | -0.33 | 0.36 | 0.46 | 0.73 | -0.50 | 0.14 | -0.85 | 0.09 | 0.95 | 1 | | | | |
| MnO | 0.75 | 0.72 | 0.66 | 0.78 | 0.55 | 0.62 | 0.79 | -0.38 | 0.52 | 0.80 | 0.60 | -0.51 | -0.21 | -0.74 | -0.09 | 0.85 | 0.80 | 1 | | | |
| CaO | 0.09 | 0.13 | 0.35 | -0.05 | 0.48 | 0.02 | 0.16 | 0.86 | -0.14 | 0.35 | 0.73 | -0.30 | 0.79 | -0.10 | -0.35 | 0.10 | 0.10 | 0.09 | 1 | | |
| P ₂ O ₅ | 1.00 | 1.00 | 0.82 | 0.81 | 0.88 | 0.97 | 0.99 | -0.25 | 0.35 | 0.35 | 0.75 | -0.44 | 0.25 | -0.80 | 0.18 | 0.93 | 0.98 | 0.71 | 0.16 | 1 | |




Figure. 4 a-d Correlations between Fe_2O_3 and As, Pb, Zn and Cu.

3.4. Sediment Pollution Assessment (SPA)

3.4.1. Contamination Factor (CF) and Degree of Contamination (C_d)

The values of contamination factors (CF) and degree of contamination (C_d) of some biogenic elements in the surface sediments of Conakry mangrove are presented in Table 4. The interpretation of the values obtained were based on [12, 14] respectively. The CF of As and Cr in the surface sediment samples ranged from 5.23 to 12.27 and 8.73 to 14.03 with an average of 7.21 and 10.08 as well as C_d of 50.44 and 70.58 accordingly. This indicates that As and Cr contamination are very high with strong anthropogenic input. Cu has a CF range of 0.16 to 0.28 with an

average of 0.18 and C_d of 1.29. Thus, Cu has low contamination and low C_d . The CF values of Pb, Zn and Ni ranged from 1.61 to 4.76; 1.99 to 2.33 and 1.63 to 2.21 with an average CF of 2.45, 2.16 and 1.84 respectively. Also, their C_d values are 17.15, 15.14 and 12.90 correspondingly. Hence, Pb, Zn and Ni are moderately contaminated with a considerable degree of contamination.

Table 4. Contamination factors and degrees of contamination of Conakry mangrove surface sediments.

| | As | Cr | Cu | Pb | Zn | Ni |
|----------------|-------|-------|------|-------|-------|-------|
| CKRY1 | 5.91 | 9.20 | 0.22 | 1.72 | 2.17 | 1.66 |
| CKRY2 | 5.45 | 8.73 | 0.09 | 1.67 | 2.11 | 1.63 |
| CKRY3 | 5.45 | 9.38 | 0.16 | 1.72 | 2.20 | 1.95 |
| CKRY4 | 5.23 | 8.80 | 0.16 | 1.67 | 2.03 | 1.74 |
| CKRY5 | 5.68 | 9.20 | 0.16 | 1.61 | 1.99 | 1.53 |
| CKRY6 | 12.27 | 14.03 | 0.28 | 4.76 | 2.31 | 2.18 |
| CKRY7 | 10.45 | 11.24 | 0.22 | 4.00 | 2.33 | 2.21 |
| Average | 7.21 | 10.08 | 0.18 | 2.45 | 2.16 | 1.84 |
| C _d | 50.44 | 70.58 | 1.29 | 17.15 | 15.14 | 12.90 |

3.4.2. Enrichment Factor (EF)

The enrichment factor values of some elements in the mangrove surface sediments of Conakry are presented in Table 5. The EF values of Pb, Zn, Cu, Ni, V, Sr, Y, Nb, Zr, Th, Sc and F are less than 2. However, Cu, Sr, Y, Nb, Zr, Th, Sc and F could be described as depleted while Pb, Zn, Ni and V are minimally enriched and polluted. As and Cr

had EF values in the range of 2 to 5 and implies they are moderately enriched and polluted. The EF of Cl is less than 20 and implies significant enrichment and pollution. Further more, [32] posits that EF values between 0.5 and 1.5 indicates geogenically enriched elements while EF values greater than 1.5 are indicative of anthropogenically enriched elements. Thus, Pb, Zn, Ni and V enrichment are geogenic while the enrichment of As, Cr and Cl are anthropogenic.

Table 5. Enrichment factors of Conakry mangrove surface sediments.

| | As | Pb | Zn | Cu | Ni | Cr | V | Sr | Y | Nb | Zr | Th | Sc | F | Cl |
|---------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|-------|
| CKRY1 | 1.78 | 0.52 | 0.65 | 0.07 | 0.50 | 2.78 | 0.58 | 0.03 | 0.14 | 0.18 | 0.19 | 0.20 | 0.08 | 0.03 | 13.96 |
| CKRY2 | 1.79 | 0.55 | 0.69 | 0.03 | 0.54 | 2.87 | 0.57 | 0.04 | 0.15 | 0.19 | 0.19 | 0.18 | 0.09 | 0.04 | 11.58 |
| CKRY3 | 1.77 | 0.56 | 0.72 | 0.05 | 0.63 | 3.05 | 0.58 | 0.05 | 0.15 | 0.19 | 0.16 | 0.18 | 0.11 | 0.13 | 14.65 |
| CKRY4 | 1.79 | 0.57 | 0.70 | 0.05 | 0.60 | 3.02 | 0.54 | 0.04 | 0.16 | 0.20 | 0.17 | 0.19 | 0.09 | - | 22.30 |
| CKRY5 | 1.88 | 0.53 | 0.66 | 0.05 | 0.50 | 3.04 | 0.57 | 0.04 | 0.15 | 0.19 | 0.29 | 0.18 | 0.09 | 0.04 | 16.78 |
| CKRY6 | 2.70 | 1.03 | 0.51 | 0.06 | 0.48 | 3.09 | 0.72 | 0.02 | 0.11 | 0.13 | 0.16 | 0.12 | 0.07 | 0.05 | 13.36 |
| CKRY7 | 2.47 | 0.95 | 0.55 | 0.05 | 0.52 | 2.66 | 0.67 | 0.03 | 0.12 | 0.14 | 0.16 | 0.19 | 0.09 | 0.05 | 9.42 |
| Average | 2.03 | 0.67 | 0.64 | 0.05 | 0.54 | 2.93 | 0.61 | 0.04 | 0.14 | 0.17 | 0.19 | 0.18 | 0.09 | 0.06 | 14.58 |

3.4.3. Pollution Index (PI) and Modified Pollution Index (MPI)

The PI and MPI values obtained from the Conakry mangrove surface sediments are shown in Table 6. The values were interpreted based on [19, 20] for PI and MPI respectively. The PI value of Cu is less than 0.7 and implies that Cu is unpolluted. Zn and Ni had PI values less than 3. As such, they are severely polluted. The PI values of Pb,

As and Cr exceeded 3 and therefore are heavily polluted. However,, Cu, Pb, Zn and Ni had MPI values of less than 1. Thus, they are unpolluted. MPI value of As is 2.26 while that of Cr is 3.01. As such, As is moderately polluted while Cr is moderately-heavily polluted.

Table 6. Pollution Index and Modified Pollution Index of Conakry mangrove surface sediments.

| | As | Cr | Cu | Pb | Zn | Ni |
|-----|-------|-------|------|------|------|------|
| PI | 10.06 | 12.22 | 0.24 | 3.79 | 2.25 | 2.03 |
| MPI | 2.26 | 3.01 | 0.06 | 0.87 | 0.68 | 0.59 |

3.5. Potential Ecological Risk Assessment (PERA)

3.5.1. Potential Contamination Index (PCI)

The potential contamination index values obtained varied among elements as

presented in Table 7. It was found that As, Cr and Pb had PCI values above 3. This suggests that these elements have severe or very severe contamination potentials in the Conakry mangrove. Zn and Ni had PCI values of 2.33 and 2.21 respectively and thus, are of moderate contamination potential. However, Cu had a PCI value below 1 and as such, is of potentially low contamination within the Conakry mangrove.

Table 7. Potential contamination index of Conakry mangrove surface sediments.

| | As | Cr | Cu | Pb | Zn | Ni |
|-----|-------|-------|------|------|------|------|
| PCI | 12.27 | 14.03 | 0.28 | 4.67 | 2.33 | 2.21 |

3.5.2. Potential Ecological Risk Factor (Er^i) and Potential Ecological Risk Index (RI)

There are variations in the potential ecological risk factors obtained for same element at different locations. However, the

assessment of the potential ecological risk in the mangroves of Conakry was based on the average ecological risk factor for each of the elements examined. As had an average ecological risk factor of 72.06 and signifies

that As in the Conakry mangrove posed a moderate potential ecological risk. Cr, Cu, Pb, Zn and Ni all had average risk values below 40. This implies that these elements have low potential ecological risk. Similarly, it was found that the RI of As is 504.4 and

indicates that As is of considerable ecological risk. However, Cr, Cu, Pb, Zn and Ni were found to have RI values below 150. Thus, these elements are of low ecological risk. The Er^i and RI values are shown in Table 8.

Table 8. Potential Ecological Risk Factor and Potential Ecological Risk Index of Conakry mangrove surface sediments.

| | As | Cr | Cu | Pb | Zn | Ni |
|---------|-------|--------|------|-------|-------|-------|
| CKRY1 | 59.1 | 18.4 | 1.1 | 8.6 | 2.17 | 8.3 |
| CKRY2 | 54.5 | 17.46 | 0.45 | 8.35 | 2.11 | 8.15 |
| CKRY3 | 54.5 | 18.76 | 0.8 | 8.6 | 2.2 | 9.75 |
| CKRY4 | 52.3 | 17.6 | 0.8 | 8.35 | 2.03 | 8.7 |
| CKRY5 | 56.8 | 18.4 | 0.8 | 8.05 | 1.99 | 7.65 |
| CKRY6 | 122.7 | 28.06 | 1.4 | 23.8 | 2.31 | 10.9 |
| CKRY7 | 104.5 | 22.48 | 1.1 | 20 | 2.33 | 11.05 |
| Average | 72.06 | 20.17 | 0.92 | 12.25 | 2.16 | 9.21 |
| RI | 504.4 | 141.16 | 6.45 | 85.75 | 15.14 | 64.5 |

4. CONCLUSION

The surface sediment samples of Conakry mangrove, Guinea were found to have variations in element concentration with location. These variations may not be entirely source bound but also due to some other factors such as redox conditions and microbial activities. Based on average concentrations, Cl; Fe_2O_3 and Sc; MnO had

the highest and lowest concentrations among the analyzed trace and major elements respectively. TiO_2 has a strong As-Pb-Cu-Cr-V; weak Th-F-Cl and negative Sr-TS-Br associations. Fe_2O_3 was found to have strong association with As,Pb, Zn,Cu, Ni, Cr, V and TiO_2 ; weakly associated with Zr, Th, F and Cl but had negative association with Sr, TS and Br. Thus both TiO_2 and Fe_2O_3 geogenically influenced the concentration of most of the elements analyzed. However, Fe_2O_3 had a greater influence. The observed

high Cu:Zn ratio in all the samples is a strong indication that the sediments were in an anoxic conditions.

As and Cr had very high contamination factors with strong anthropogenic pollution. Pb, Zn and Ni had moderate contamination and a considerable degree of contamination. The enrichment assessment showed that Cl was significantly enriched while As and Cr were moderately enriched. However, Pb, Zn, Ni and V were minimally enriched while Cu, Sr, Y, Nb, Zr, Th, Sc and F were depleted. Generally, the enrichment of As, Cr and Cl were anthropogenic but the enrichment of Pb, Zn, Ni and V were geogenic. The potential ecological risk assessment showed that As had a potentially moderate to considerable ecological risk but Cr, Cu, Pb, Zn and Ni had potentially low ecological risk.

APPENDIX A. SUPPLEMENTARY DATA

Supplementary data to this article can be sourced online from <http://ir.lib.shimane-u.ac.jp/46655>.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly used products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing

company rather it was funded by personal efforts of the authors.

REFERENCES

1. Nwawuiké, N., Ishiga, H., 2018. Elemental Composition of Core Sediments in Niger Delta Mangrove, Nigeria. *Journal of Geography, Environment and Earth Science International*, 16(3), 1-18.
2. Dousova, B., Buzek, F., Rothwell, J., Krejcová, S., Lhotka, M., 2012. Adsorption Behaviour of Arsenic Relating to Different Natural Solids: Soils, Stream Sediments and Peats. *Science of the Total Environment*.
3. Arnason, J. G., Fletcher, B. A., 2003. A 40+ Year Record of Cd, Hg, Pb and U Deposition in Sediments of Patroon Reservoir, Albany County, NY, USA. *Environmental Pollution*, 123, 383 - 391.
4. Onivogui, G., Saidouba Balde, S., Bangoura, K., Barry, M. K., 2013. Évaluation des risques de pollution en métaux lourds (Hg, Cd, Pb, Co, Ni, Zn) des eaux et des sédiments de l'estuaire du fleuve Konkouré (Rep. de Guinée). *Afrique Science*, 09(3), 36 - 44.
5. United Nations Environment Programme., 2007. *Mangroves of Western and Central Africa*. UNEP-Regional Seas Programme/UNEP-WCMC.
6. Wright, H. E., McCullough, J., 2006. An Ecological, Socio-economic and Conservation Overview of Northwestern Guinea. *Rapid Assessment Programme*, 102 - 106. doi: 10.1896/054.041.0106.

- 7.GEF., 2004. GEF Project Brief: Guinea Coastal Marine and Biodiversity Management. African Regional Office.
- 8.Nations Encyclopedia. Retrieved from www.nationsencyclopedia.com on 23rd March, 2020.
- 9.Deckart, K., Feraud, G., Bertrand, H., 1997. Age of Jurassic Continental Tholeiites of French Guyana, Surinam and Guinea: Implications for the Initial Opening of the Central Atlantic Ocean. *Earth and Planetary Science Letters*, 150, 205 - 220.
- 10.Ishiga, H., Diallo, I. M., Malick, B. M. L., Ngulimi, M.F., Magai, P. J., Shati, S.S., 2016. Geochemical Approach to Evaluate Deforest of Mangroves. *Geoscience Dept. Shimane University*, 34, 95 -106.
- 11.Ogasawara, M., 1987. Trace Element Analysis of Rock Samples by X-ray Fluorescence Spectrometer Using Rh Anode Tube. *Bulletin of the Geological Survey Japan*, 38, 57 - 68.
- 12.Tomlinson, D. L., Wilson, J. G., Harris, C. R., Jeffrey, D. W., 1980. Problems in the Assessment of Heavy-metal Levels in Estuaries and the Formation of a Pollution Index. *Helgolander Meeresuntersuchungen*, 33(1-4), 566 - 575.
- 13.Rudnick, R. L., Gao, S., 2003. Composition of the Continental Crust. In *Treatise on Geochemistry* (Holland, H. D., & Turekian, K. K. eds). Elsevier, 3, 1 - 64.
- 14.Hakanson, L., 1980. An Ecological Risk Index for Aquatic Pollution Control: A Sedimentological Approach. *Water Research*, 14, 975 - 1001.
- 15.Seshan, B. R. R., Natesan, U., Deepthi, K., 2010. Geochemical and Statistical Approach for Evaluation of Heavy Metal Pollution in Core Sediments in Southeast Coast of India. *International Journal of Environmental Science and Technology* 7 (2), 291-306.
- 16.Reimann, C., de Caritat, P., 2000. intrinsic Flaws of Element Enrichment Factors (EFs) in Environmental Geochemistry. *Environmental Science and Technology* 34 (24), 5084-5091.
- 17.Helz, G. R., 1976. Trace element inventory for the northern chesapeake bay with emphasis on the influence of man. *Geochemica et Cosmochimica Acta*, 40(6), 573 - 580.
- 18.Goher, M. E., Farhat, H. I., Abdo, M. H., Salem, S.G., 2014. Metal pollution assessment in the surface sediment of Lake Nasser, Egypt. *Egyptian Journal of Aquatic Research*, 40, 213 - 224.
- 19.Nemerow, N. I., 1991. *Stream, Lake, Estuary and Ocean Pollution*. New York: Van Nostrand Reinhold, p.472.
- 20.Brady, J. P., Ayoko, G. A., Martens, W. N., Goonetilleke, A., 2018. Development of a Hybrid Pollution Index for Heavy Metals in Marine and Estuarine Sediments. *Environmental Monitoring Assessment*.
- 21.Davault, V., Rognerud, S., 2001. Heavy Metal Pollution in Sediments of the Pasvik River Drainage. *Chemosphere*, 42, 9 - 18.

22. Nwawuiké, N., 2019. Environmental Risk Assessment of Geochemical Composition and Spatial Dynamics in Sediments of Niger Delta Mangrove, Nigeria. Doctoral Thesis, Dept. of Geoscience Shimane University. <http://ir.lib.shimane-u.ac.jp/46655>.
23. Ishiga, H., Dozen, K., Sampei, Y., 1999. Geochemical Constraints on Marine Invasion and Provenance Change Related to the Opening of the Japan Sea: An Example from the Lower Miocene Shale in Hoda Section, Shimane Peninsula, SW Japan. *Journal of Asian Earth Sciences*, 17, 443 - 457.
24. Roser, B. P., 2000. Whole-rock Geochemical Studies of Clastic Sedimentary Suits. *Memoirs of the Geological Society of Japan*, 57, 73 - 89.
25. Skilbeck, C.G., Cawood, P. A., 1994. Provenance History of a Carboniferous Gondwana Margin Forearc Basin, New England Fold Belt, Eastern Australia: Modal and geochemical constraints. *Sedimentary Geology*, 93, 107 - 133.
26. Dalai, B., Ishiga H., 2013. Identification of Ancient Human Activity Using Multi-element Analysis of Soils at a Medieval Harbor Site in Masuda City, Shimane Prefecture, Japan. *Earth Science (Chikyu Kagaku)*, 67, 75 - 86.
27. Diallo, I. M., Ishiga H., 2016. Evaluation of Trace Metal Contamination in Ise Bay, Mie Prefecture, Central Japan, Based on Geochemical Analysis of Tidal Flat Sediments. *Environment and Pollution*, 5(1), 92 - 109.
28. Dinelli, E., Cortecchi, G., Lucchini, F., Zantedeschi, E., 2005. Sources of Major and Trace Elements in Stream Sediments of the Arno River Catchment (Northern Tuscany, Italy). *Geochemical Journal*, 39, 531 - 545.
29. Cooper, D. C., Thornton, I., 1994. *Drainage Geochemistry in Contaminated Terrains Handbook of Exploratory Geochemistry* (Hale, M., Plant, J. A., eds). Elsevier, 6, 447 - 497.
30. Tolonen, K., Merilainen, J., 1983. Sedimentary Chemistry of a Small Polluted Lake, Galltrask, S. Finland. *Hydrobiologia*, 103, 309 - 318.
31. Kauppila, P., Weckström, K., Vaalgamaa, S., Korhola, A., Pitkanen, H., Reuss, N., Drew S., 2005. Tracing Pollution and Recovery using Sediments in an Urban Estuary, Northern Baltic Sea: Are we far from Ecological Reference Condition. *Marine Ecology Progress Series*, 290, 35 - 53.
32. Zhang, J., Liu, C. L., 2002. Riverine Composition and Estuarine Geochemistry of Particulate Metals in China-Weathering Features, Anthropogenic Impact and Chemical Fluxes. *Estuarine, Coastal and Shelf Science*, 54(6), 1051 - 1070.