

Original Research Article

Effect of Packaging Type on Quality of Red Tilapia Shredded Added by Cayenne Pineapple (*Ananas comosus (L) Merr*) Fruit Solution

ABSTRACT

The purpose of this study was to determine the right type of packaging for organoleptic quality and analysis of the peroxide value of red tilapia. This research uses a method consisting of 3 treatments, namely A (Aluminum Foil), B (plastic *High Density Polyethylene*), C (plastic *Low Density Polyethylene*) using 20 panels as replicates. Parameters observed in this study were organoleptic characteristics (appearance, texture, aroma, and taste) based on the panelists' preference level using hedonic test, chemical analysis (peroxide number test) and storage for 28 days in an incubator at 25°C and 45°C for each treatment. The data from the chemical test were analyzed descriptively, while the organoleptic test was analyzed using the Friedman test. Based on the Friedman food test, it was found that HDPE plastic packaging with the addition of cayenne pineapple was the preferred treatment compared to other treatments. Based on the Bayes test, taste is the most important criterion compared to appearance, texture, and aroma. The results of the peroxide value analysis for each type of packaging treatment during storage for 28 days at 25°C showed results with a peroxide value of 0 meq/kg and 45°C showed results with a weekly value of 0 meq/kg every week.

Key words: Shredded, aluminium foil, peroxide analysis, HDPE, LDPE, organoleptic

1. INTRODUCTION

Fish is one of the foodstuffs that is quite easy to obtain in Indonesia, given that the potential of our sea is so vast coupled with many sources of freshwater aquaculture, one of which is red tilapia (*Oreochromis niloticus*) which is one of the aquaculture commodities that is favored by its development. Ministry of Marine Affairs and Fisheries (KKP 2010). Fish production per capita in Indonesia from year to year has increased significantly, namely in 2019, the fish consumption figure reached 54.49 kilograms/capita. This figure increased when

compared to 2018 which reached 50.69 kg/capita. This figure is slightly higher than the target of 50.65 kg/capita.

According to Kusumayanti (2011) red tilapia (*Oreochromis niloticus*) a freshwater fish species was chosen as raw material because it has thick, compact flesh and is easily separated from the bones and spines. Red tilapia has a fat content of 4.1% including low or medium fat fish (Astawan 2003). Fat in meat also contributes to flavor because it is very necessary, from that thought, many business actors started to make processed fishery products made from red tilapia because it is known to have several advantages, one of which is fast fish growth, thick and compact meat compared to black tilapia, high tolerance for the environment and ease of cultivation, and red tilapia are less susceptible to disease when cultivated when compared to black tilapia. Therefore, it is necessary to use efforts to encourage fishery products, especially red tilapia as raw material for making shredded fish.

Shredded fish is a processed fishery product made from fish meat, through a combination of processing processes, namely the process of steaming, grinding and frying with the addition of spices. Shredded fish is a dry processed product, in its manufacture it uses ingredients such as oil, coconut milk and seasonings so that it is sensitive to air, prone to rancidity during storage, even though it is pressed to remove the oil content, but not all of it can be removed (Tridiyani 2012). During storage, shredded fish will still experience a decrease in quality due to chemical and physical changes that occur in the shredded fish during storage. According to Sudarmadji et al. (2003) during storage of food products containing fat or oil will usually experience a process of rancidity during the storage process.

According to Eko et al. (2011), abon which is processed by deep frying has a rancidity value that tends to increase significantly compared to shredded pan frying during 29 days of storage at room temperature. This shows that the oxidation process in the abon that is processed by deep frying runs faster than that of the abon that is processed by pan frying. This is due to the fact that the amount of oil used in the deep-fried shredded meat is more than the pan frying method so that the more the number of double bonds in the oil, the faster the oxidation rate. The more double bonds there are, the more likely it is for hydroperoxides to form. The use of cayenne pineapple solution in the manufacture of red tilapia shredded is intended as a natural preservative as an antioxidant substance to produce good quality shredded fish products.

Pineapple (*Ananas comosus* (L) Merr) is the most economical and highly nutritious food ingredient which has a high concentration of vitamin C, phenolic compounds and -carotene as natural antioxidants (Preeti et al 2018). According to Edi et al. (2015) the addition of cayenne pineapple juice provided an increase in antioxidant activity against corn cob extract, the increase was due to the increase in the total phenolic content contained in the extract and had a good ability to ward off free radicals.

The shelf life of a food product is influenced by packaging and environmental temperature. During the storage period, the room temperature for shredded tilapia products ranged from 25°C-27°C. The function of the packaging of a product is to maintain the quality of freshness of food products, provide convenience for storage, inhibit contamination from the water and surroundings by microorganisms, and provide attractiveness to consumers (Sauvage 1996). Packaging issues are very important and need to be considered, because packaging has a huge influence on the smooth marketing of packaged goods. Therefore, in order to maintain the quality of shredded red tilapia, the type of packaging and storage temperature must receive special attention (Afrianto 2005).

Packaging is one way or method to provide protection for food that has been produced either in the form of packages or placing the product in a container. This is done so that the product can be protected from pollution (chemical and microbial compounds), physical damage (friction and impact), environmental compounds (oxygen, water vapor) and animal disturbances such as insects, so that product safety quality is maintained and stored for a period of time. the longer one. HDPE is a plastic material that is safe to use, because HDPE is able to prevent chemical reactions between plastic made from HDPE and the food or beverage it packs. HDPE has material properties that are stronger, harder, opaque, more resistant to high temperatures and can be used for frozen storage up to -50°C (Harumningtiyas 2010).

The basic packaging materials used for shredded products in small industries are High Density Polyethylene (HDPE) plastic and aluminum foil. Food packaging made of plastic and aluminum foil functions as a packaging material for a wet and dry product as a protective material for the product from damage, contamination from free air, which causes a decrease in the quality of a product. The use of plastic and aluminum foil as packaging for a food product is due to their light weight, multipurpose, strong, easy to shape, and able to ward off water absorption. The use of the type of packaging material is of course adapted to the properties of the packaged material, each type of packaging material will have a different effect on the packaged product (Makinde et al 1976). Based on this, the authors are interested in conducting research on the effect of different types of

packaging on the quality of shredded red tilapia with added cayenne pineapple solution during storage.

2. MATERIALS AND METHODS

2.1 Time and Place

The research was carried out from March to April 2021 at the Fishery Product Processing Laboratory, Faculty of Fisheries and Marine Sciences, Padjadjaran University for the manufacture of shredded tilapia and organoleptic testing. The peroxide value test was carried out at the Food Technology Laboratory, Faculty of Engineering, Pasundan University.

2.2 Materials and Tools

The materials used in the study included 1500 g of red tilapia fish fillet obtained from the Ciroyom market in Bandung, 300 ml of cayenne pineapple, spices, 800 ml of cooking oil, HDPE plastic, LDPE plastic and aluminum foil. The tools used are knives, cutting boards, basins, spinners, blenders, pans, strainers, analytical scales, pans, stoves, incubators, measuring cups and spatulas.

2.3 Research Procedure

2.3.1 Preparation of Cayenne Pineapple Fruit Solution

Choose a ripe pineapple then the pineapple solution can be made by smoothing the pineapple flesh using a blender for 5 minutes then separate the pulp and the solution using gauze so that you get a pineapple solution.

2.3.2 Shredded Making Procedure

Preparation of tools and materials needed during the making of abon then the contents of the stomach, gills, and scales of tilapia are removed and washed clean after the tilapia is filled, put it in the pan to be steamed for 30 minutes on the stove at a temperature of 100-150°C after steaming the fish is cooled, separated between the meat, bones and skin of the fish so that the fish meat can be shredded. The spices, cayenne pineapple and fish meat are mixed and then sautéed until half cooked for 10-15 minutes at 100-150°C then 800 ml of oil is heated at 100-150°C and the batter is shredded tilapia fried until brown. The cooked tilapia shreds are drained and the oil is removed using a spinner until the texture is like meat fibers, then the shredded meat is re-shredded and packaged after it has cooled.

2.4 Research Method

The research method used is the experimental method. Data analysis was carried out with non-parametric statistics with 3 treatments and 20 semi-trained panelists as a test to determine the level of panelists' acceptance of the type of packaging on shredded red tilapia added with a 20% solution of cayenne pineapple. The treatments applied in this study were:

- 1) Treatment A: aluminum foil packaging
- 2) Treatment B: HDPE plastic packaging
- 3) Treatment C: LDPE plastic packaging

Red tilapia shreds were stored in an incubator at 25°C and 45°C for 28 days of storage: Observations were made every 7 days, on the 1st day to determine the effect of different packaging on temperature with the addition of a solution of cayenne pineapple on (7th day; 14th day; 21st; 28th day) the parameters observed included organoleptic tests (appearance, texture, aroma, and taste) based on the panelists' preference level and chemical analysis (peroxide number test) on shredded for each treatment.

2.5 Data Analysis

The results of the organoleptic test were analyzed using non-parametric, namely using the Friedman test to determine the level of preference of the panelists on the organoleptic characteristics of shredded. According to Daniel (1989) if the Friedman test shows significant results, a further test is carried out, namely multiple comparisons to find out the difference between each treatment using the Bayes method by considering the weight of the criteria and the median value. Meanwhile, the peroxide value test was analyzed descriptively.

3. RESULT

3.1 Organoleptic Characteristics of Shredded

Organoleptic test observations were carried out using a hedonic (levels of pleasure) scale with observation parameters in the form of appearance, aroma, texture, and taste to determine the panelist preference for red tilapia shredded packaged in this type of packaging as a treatment with the addition of cayenne pineapple solution.

3.1.1 Appearance

Appearance is one of the organoleptic parameters that has an important role for panelists to assess the quality of a product. Organoleptic characteristics of the appearance of red tilapia shredded on day 1 without incubator storage to determine the effect of different packaging on temperature with the addition of

cayenne pineapple fruit solution on the appearance characteristics of shredded. The first thing that the panelists noticed when making observations was the color and surface of the shredded. The average value of the hedonic test on the appearance of shredded day 1 is presented in Table 1.

Table 1. Value of Appearance of Red Tilapia Shredded on Day 1

Treatment	Median	Average appearance
A (Alumunium Foil)	9	8,1 a
B (Plastik HDPE)	9	8,2 a
C (Plastik LDPE)	9	8,1 a

Note: The number followed by the same letter on the average treatment shows that there is no significant difference according to the multiple comparison test with a 5% test level.

Friedman test results at the test level of 5%. showed that the application of the type of packaging with the addition of a solution of pineapple cayenne did not give a significant effect on the appearance characteristics of red tilapia abon with a median value ranging from 9 on day 1 (organoleptic test) in each treatment with color appearance. specific type of shredded chocolate, homogeneous fiber and brilliant color.

The following average values of hedonic tests on the appearance of shredded 7th day, 14th day, 21st day, and 28th day of incubator storage at 25°C and 45°C for 28 days are presented in Table 2.

Table 2. Appearance Value of Red Tilapia Shredded Storage 28 Days

Day Treatment	A (Alumunium Foil)				B (HDPE Plastic)				C (LDPE Plastic)			
	Median		Average		Median		Average		Median		Average	
	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C
7	9	8	8,0a	7,9a	9	9	8,2a	8,1a	9	8	8,1a	7,9a
14	9	8	8,0a	7,7a	9	8	8,1a	8,0a	9	8	8,1a	7,9a
21	8	8	8,0a	7,5a	9	8	8,1a	7,9a	9	7	8,1a	7,7a
28	7	7	7,1a	6,2a	8	7	7,8a	7,0a	8	7	7,7a	6,5a

Note: The number followed by the same letter on the average treatment shows that there is no significant difference according to the multiple comparison test with a 5% test level.

Friedman test results at the 5% test level showed that the application of the type of packaging with the addition of a solution of pineapple cayenne fruit was not significantly different, which means that all treatments of aluminum foil, HDPE plastic, and LDPE plastic packaging did not have a significant effect on the appearance characteristics of red tilapia shredded with a median value. The highest temperature was at 25°C and 45°C ranging from 9 and 8 in each packaging treatment with the results of the appearance of the shredded brown color being specific for the type of shredded, homogeneous fiber and brilliant color. The value of appearance in each packaging treatment is generated because of the Maillard reaction. The Maillard reaction is a reaction between the amino group of a free amino acid peptide chain residue or protein with the carbonyl group of a carbohydrate when both are heated or stored for a long time (Lakshmi 2014).

3.1.2 Scent

Scent is the smell of food products caused by volatile compounds from food entering the nasal cavity and being felt by the olfactory system so as to form a response. Aroma can determine the delicacy of food and affect the level of acceptance (Winarno 2002 in Tribaditia 2016). Organoleptic characteristics of red tilapia shredded aroma on day 1 without incubator storage to determine the difference in packaging to temperature with the addition of chili pineapple on the characteristics of shredded aroma. The average value of the hedonic test on the first day of shredded aroma is presented in Table 3.

Table 3. Value of Scent of Red Tilapia Shredded on Day 1

Treatment	Median	Average appearance
A (Alumunium Foil)	9	8,1 a
B (HDPE Plastic)	9	8,2 a
C (LDPE Plastic)	9	8,1 a

Note: The number followed by the same letter on the average treatment shows that there is no significant difference according to the multiple comparison test with a 5% test level.

The results of the Friedman test at the 5% test level showed that the application of the type of packaging with the addition of a solution of cayenne pineapple did not have a significant effect on the aroma characteristics of red tilapia abon with a median value of around 9 on day 1 (organoleptic test) in each treatment with the results having Specific scent of abon type and no fishy smell of fish.

The following average values of hedonic tests on the scent of shredded meat on the 7th, 14th, 21st, and 28th days of incubator storage at 25°C and 45°C for 28 days are presented in Table 4.

Table 4. Scent Value of Red Tilapia Shredded Storage 28 Days

Day Treatment	A (Alumunium Foil)				B (HDPE Plastic)				C (LDPE Plastic)			
	Median		Average		Median		Average		Median		Average	
	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C
7	9	8	8,1a	7,8a	9	8	8,2a	8,0a	9	8	8,1a	7,7a
14	9	8	8,0a	7,8a	9	8	8,1a	7,8a	9	8	8,1a	7,7a
21	8	7	8,0a	6,6a	9	8	8,1a	6,7a	8	7	8,1a	6,5a
28	8	7	8,0a	6,2a	9	8	8,1a	6,6a	8	7	8,1a	6,3a

Note: The number followed by the same letter on the average treatment shows that there is no significant difference according to the multiple comparison test with a 5% test level.

Friedman test results showed that the application of the type of packaging with the addition of a solution of cayenne pineapple fruit was not significantly different, which means that all treatments of aluminum foil, HDPE plastic, and LDPE plastic did not have a significant effect on the aroma characteristics of red tilapia abon with the highest median value at 25°C. C and 45°C ranged from 9 and 8 in each packaging treatment with the results of a specific shredded type of shredded aroma and no fishy smell of fish.

The scent value in each packaging treatment was still preferred according to the panelist's assessment because the formulation of making shredded in each packaging was the same, and during the frying process, bay leaves, lemongrass stalks, and lime leaves were added which are known to be able to remove the thick fishy smell of fish, then when By steaming, fish protein will be hydrolyzed into amino acids, namely glutamic acid, glycine, alanine and lysine, giving rise to an aroma flavor and a delicious savory taste (Winarno 2002).

The level of assessment of the aroma characteristics of the panelists did not decrease drastically because during the storage of red tilapia abon there was no oxidation process caused by the lipid peroxide reaction which caused a rancid odor. Food damage can also occur by the environment so that it can be controlled with packaging and natural preservatives that can be used, for example mechanical damage, changes in the water content of ingredients, and interactions

with oxygen, therefore the type of packaging that can control and is often used to package shredded is polyethylene packaging. This is because polyethylene packaging has a high density, is resistant to temperature and humidity, and has a low water absorption capacity so as to protect the quality of shredded meat (Ahmad et al. 2016).

Handling to prevent rancidity in the manufacture of red tilapia shredded using natural preservatives such as the addition of a solution of cayenne pineapple as a natural preservative and antioxidant to produce shredded fish products that have a specific flavor of shredded, because cayenne pineapple is the most economical and highly nutritious food ingredient. which has a high concentration of vitamin c, phenolic compounds and -carotene as natural antioxidants (Preeti et al 2018).

3.1.3 Texture

Texture is an organoleptic assessment parameter that affects consumer acceptance of a product. Texture assessment can be done by feeling the texture of the product using a touch of the skin surface or by using the fingertips (Azka et al 2020). Organoleptic characteristics of red tilapia shredded texture on day 1 without incubator storage to determine the effect of different packaging on temperature with the addition of cayenne pineapple fruit solution on the texture characteristics of shredded. The average value of the hedonic test on the 1st day of shredded texture is presented in Table 5.

Table 5. Value of Texture of Red Tilapia Shredded on Day 1

Treatment	Median	Average appearance
A (Aluminium Foil)	9	8,1 a
B (HDPE Plastic)	9	8,2 a
C (LDPE Plastic)	9	8,1 a

Note: The number followed by the same letter on the average treatment shows that there is no significant difference according to the multiple comparison test with a 5% test level.

Friedman test results at the 5% test level. showed that the application of the type of packaging with the addition of a solution of pineapple cayenne did not have a significant effect on the texture characteristics of red tilapia shredded with a median value of around 9 on day 1 (organoleptic testing) in each treatment with the results having a soft fibrous texture and not clumping.

The following average values of hedonic tests on the shredded texture on the 7th day, 14th day, 21st day, and 28th day of incubator storage at 25°C and 45°C for 28 days are presented in Table 6.

Table 6. Texture Value of Red Tilapia Shredded Storage 28 Days

Day Treatment	A (Alumunium Foil)				B (HDPE Plastic)				C (LDPE Plastic)			
	Median		Average		Median		Average		Median		Average	
	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C
7	9	8	8,0a	7,8a	9	8	8,1a	7,8a	9	8	8,1a	7,7a
14	9	8	8,0a	7,8a	9	8	8,1a	7,8a	9	8	8,1a	7,7a
21	8	7	8,0a	6,8a	8	7	8,1a	6,8a	8	7	8,1a	6,5a
28	8	7	8,0a	6,2a	8	7	8,1a	6,6a	8	7	8,1a	6,3a

Note: The number followed by the same letter on the average treatment shows that there is no significant difference according to the multiple comparison test with a 5% test level.

Friedman test results showed that the application of the type of packaging with the addition of cayenne pineapple solution was not significantly different, which means that all treatments of aluminum foil, HDPE plastic, and LDPE plastic packaging did not have a significant effect on the texture characteristics of red tilapia with the highest median value at 25°C. and 45°C ranging from 9 and 8 in each packaging treatment with the results of a soft fibrous shredded texture and not clumping.

Texture values in each packaging treatment were produced because there was no change in texture decline in each packaging treatment by the activity of microorganisms that degrade proteins into simpler compounds. According to Nur (2009) the decrease in the level of product texture assessment during storage is caused by the water activity of the food material with the storage environment, therefore the type of packaging that is often used to package shredded is polyethylene packaging. This is because polyethylene packaging has a high density, is resistant to temperature and humidity, and has a low water absorption capacity so as to protect the quality of shredded meat (Ahmad et al. 2016).

3.1.4 Flavor

Taste is a parameter that greatly determines the consumer's final decision to accept or reject a food product, if other parameters are considered good but the taste is not liked then the product will be rejected (Nirmala et al. 2016). Organoleptic characteristics of red tilapia shredded flavor on day 1 without incubator storage to determine the effect of different packaging on temperature with the addition of cayenne pineapple fruit solution on the characteristics of shredded taste. The average value of the hedonic test on the 1st day of shredded taste is presented in Table 7.

Table 7. Value of Flavor of Red Tilapia Shredded on Day 1

Treatment	Median	Average appearance
A (Alumunium Foil)	9	8,1 a
B (HDPE Plastic)	9	8,2 a
C (LDPE Plastic)	9	8,1 a

Note: The number followed by the same letter on the average treatment shows that there is no significant difference according to the multiple comparison test with a 5% test level.

The results of the Friedman test at the 5% test level showed that the application of the type of packaging with the addition of a solution of cayenne pineapple did not have a significant effect on the taste characteristics of red tilapia shredded with a median value of around 9 on day 1 (organoleptic testing) in each treatment with the results having a good taste. Specific shredded fish is very pronounced with a savory taste that tends to be sweet.

The following average values of hedonic test on shredded taste on 7th day, 14th day, 21st day, and 28th day of incubator storage at 25°C and 45°C for 28 days are presented in Table 8.

Table 8. Flavor Value of Red Tilapia Shredded Storage 28 Days

Day Treatment	A (Alumunium Foil)				B (HDPE Plastic)				C (LDPE Plastic)			
	Median		Average		Median		Average		Median		Average	
	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C	25°C	45°C
7	9	8	8,1a	8,0a	9	9	8,2a	8,1a	9	8	8,1a	8,0a

14	9	8	8,1a	7,9a	9	9	8,2a	8,1a	9	8	8,1a	7,9a
21	8	7	8,0a	7,6a	8	8	8,1a	8,0a	8	7	8,1a	7,9a
28	8	7	8,0a	7,2a	8	8	8,1a	8,0a	8	7	8,1a	7,5a

Note: The number followed by the same letter on the average treatment shows that there is no significant difference according to the multiple comparison test with a 5% test level.

Friedman test results showed that the application of the type of packaging with the addition of a solution of pineapple cayenne fruit was not significantly different, which means that all treatments of aluminum foil, HDPE plastic, and LDPE plastic did not significantly affect the taste characteristics of red tilapia shredded with the highest median value at 25°C. C and 45°C ranged from 9 and 8 in each packaging treatment with the result having a specific taste of shredded fish that was very pronounced with a savory taste that tends to be sweet.

The value of taste in each packaging treatment was produced because the formulation of making the floss in each package was the same and there was no loss of organic acid components present in the red tilapia floss due to storage at 25°C and 45°C so that the panelist's level of assessment of the taste did not change. decrease. Good taste is also influenced by several factors, namely chemical compounds, temperature, concentration and interactions with other flavor components.

According to Meity (2016), the process of spreading heat quickly with high temperatures will help speed up the process of releasing volatile compounds so that the unwelcome flavor of fresh fish such as the fishy smell of fish will be reduced. This is in accordance with Herliani (2008) statement that the taste image can be influenced by heating or processing that is carried out so that it results in the degradation of the preparation of the taste image and the physical properties of food ingredients, therefore the level of assessment of taste characteristics is still preferred according to the panelist assessment. Handling of the specific taste

resistance of abon, it is best to use polyethylene packaging to package the shredded meat, with this packaging having advantages, namely the characteristics of water-resistant packaging and a good degree of density (Jayadi 2016).

3.2 Decision Making Using Bayes Method

Bayes method is a tool to measure the probability of every decision taken by changing the pairwise comparison with a set of numbers that can represent the priority of the relative weight values of the criteria and alternatives. According to Siregar (2014) decision making using the Bayes method requires information in the form of probabilities for each alternative which will produce an expected value as the basis for decision making. The average value of the shredded Bayes test on the type of packaging with the addition of different concentrations of red ginger essential oil is presented in Table 9.

Table 9. Decision Matrix for Valuation of Shredded Tilapia with Bayes Method

Treatment	Criteria				Value	Value
	Appearance	Scent	Texture	Flavor	Alternative	Priority
Alufo	7,00	7,00	8,00	7,00	7,03	0,30
HDPE	8,00	7,00	8,00	8,00	8,36	0,36
LDPE	9,00	9,00	7,00	7,00	8,11	0,35
Criterion weight	0,20	0,36	0,03	0,42	23,50	1,00

Based on the Bayes test table 9 above, it can be concluded that taste is the most important criterion compared to appearance, aroma and texture. The use of this type of packaging on shredded with the addition of a solution of nana cayenne fruit obtained the highest alternative value of 8.36. Based on the preference test parameters that have been carried out on HDPE plastic treatment on shredded with the addition of a solution of pineapple cayenne fruit is the best treatment and is still preferred by the panelists.

4.3 Peroxide Number Analysis

The peroxide value is a measure of the damage or rancidity of an oil. The higher the peroxide number, the more rancid the oil is. The results of the calculation of the peroxide value of red tilapia in the type of packaging treatment with the addition of a solution of cayenne pineapple are presented in Table 10

Table 10. Results of Peroxide Number Analysis on Red Tilapia Shredded

No	Day Storage	Alumunium Foil (meq/kg)	HDPE (meq/kg)	LDPE (meq/kg)
1.	1	0,00	0,00	0,00
2.	7	0,00	0,00	0,00
3.	14	0,00	0,00	0,00
4.	21	0,00	0,00	0,00
5.	28	0,00	0,00	0,00

Based on Table 10. The results of the analysis of the peroxide value obtained on red tilapia shredded in storage on the 7th day, 14th day, 21st day, 28th day at a temperature of 25°C and 45°C in each different packaging treatment. namely treatment A (aluminum foil), treatment B (HDPE plastic), and treatment C (LDPE plastic) was 0 meq/kg which means that there is no peroxide value in red tilapia shredded products every week. This happens because of the type of packaging and the addition of natural preservatives when making shredded which can control the interaction of oxygen which causes the oxidation process caused by the lipid peroxide reaction which causes a rancid odor, one of the treatments is by using the type of packaging that is often used to package the shredded meat. polyethylene (HDPE) packaging. This is because polyethylene packaging has a high density, is resistant to temperature and humidity, and has a low water absorption capacity so as to protect the quality of shredded meat (Ahmad et al. 2016). In contrast to aluminum foil packaging, according to Julianti (2007), the thicker the aluminum foil used, the higher it is permeability of the packaging to water vapor and oxygen, while the thickness of the aluminum foil used in this study was 0.05 mm so that the packaging has a fairly high permeability to oxygen and water vapor. High packaging permeability can lead to oxidation and hydrolysis of the fat contained in shredded flying fish (Eka 2010).

According to the statement of Maulana et al (2016) that polyethylene plastic (HDPE) is able to produce the lowest peroxide value every week of 9.33 meq/kg on the 10th day of storage, 13.95 meq/kg on the 20th, 20th day of storage, 02 meq/kg on the 30th day of storage and 24.39 meq/kg on the 40th day of storage at room temperature compared to LDPE plastic and aluminum foil, the presence of this peroxide number causes bad smell and taste and causes rancidity of the oil, the greater the value of the peroxide number means the more peroxide contained in the sample (Ketaren 1986). Subsequent treatment with the addition of natural preservatives used in this study as an antioxidant substance such as cayenne

pineapple solution because cayenne pineapple is the most economical and highly nutritious food ingredient which has a high concentration of vitamin C, phenolic compounds and -carotene as natural antioxidants. Preeti et al. 2018), so that red tilapia shredded packaged in a different type of packaging with the addition of a cayenne pineapple solution did not have a peroxide value and was good for consumption.

Conclusion

Based on the results of the research, it can be concluded that red tilapia abon packaged using HDPE packaging with the addition of a solution of cayenne pineapple is the preferred treatment by panelists compared to other treatments based on the Friedman test. Based on the Bayes test, taste is the most important criterion compared to appearance, texture, and aroma. The results of the peroxide value analysis for each type of packaging treatment during 28 days of storage at 25°C showed results with a peroxide value of 0 meq/kg and 45°C showed results with a peroxide value of 0 meq/kg per week.

KEPENTINGAN BERSAING

Tidak ada kepentingan bersaing.

REFERENSI

1. Afrianto E, Liviawaty E. 2005. Food Preservation and Fish Processing. Canisius. Yogyakarta.
2. Ahmad, A.F., Maflahah I, & Rahman A. 2016. The Effect of Type of Packaging and Storage Time on Product Quality of Gambus Nugget. *AGROINTEK*, 10(2):70–75.
3. Azka, A. and Mujiyanti, A. 2020. Effect of Addition of Sodium Tripoliphosphat on Preference Level of Kamoboko Malong Fish (*Muraenesox cinerus*). *Aurelia Journal*. 1(2): 129-136.
4. Edi Suryanto, Indrika Utami Hayat and Jemmy.2015. Effect of Pineapple Juice on Antioxidant Activity of Corn Cob Extract. Udayana University. Bali.
5. Eko Nurcahya Dewi, Ratna Ibrahim and Nuzulia Yuniva. 2011. Shelf Life of Shredded Red Tilapia (*Oreochromis niloticus* trewavas) Processed With Different Frying Method. *Journal of Fisheries Science*. 6 (1) : 6-12.
6. Eka Kurniawati. 2010. Oxidation of Beef Shredded Fat During Storage At Room Temperature. Thesis. Bogor Agricultural Institute. Bogor
7. Herawati, Heny. 2008. Determination of Shelf Life in Food Products. *Agricultural Research and Development Journal*. 27(4): 124-130.
8. Hardoko., Herdanto, Lina., Marsillam, S. 2010. Utilization of Purple Sweet Potato (*Ipomoea Batatas* L. Poir) as a Partial Substitute for Wheat

- Flour and Source of Antioxidants in Bread. *Journal of Food Technology and Industry*, 20:1-2
9. Handayani, A., Alimin., and Rustiah W. O. 2014. Effect of Storage at Low Temperatures on Water Content and Fat Content in Lemuru Fish. *Al-Kimia Journal*. 2(1): 64-75.
 10. Jayadi, Queue., Anwar, Badaruddin and Sukainah, Andi. 2016. The Effect of Storage Temperature and Type of Packaging on the Quality of Flying Fish Shredded, *Journal of Agricultural Technology Education*, 2: 62-69
 11. Junianto. 2016. *Management of Fishery Products Industry*. Unpad Press. Jatinangor
 12. Makinde, M. A., Gilbert, S. G & Lacahee, P.1976. Nutritional Implication of Packaging System Food Product. *J.Food Packaging*, 9:112-123.
 13. Nurmainah, Mimi 2003. *Research Journal of Various Plastic and Paper Packaging and Their Effects*. Department of Agricultural Technology. North Sumatra.
 14. Rismarini. 2015. Estimation of Shelf Life of Skipjack Tuna (*Katsuwonus Pelamis*). Using the Arrhenius Method. Essay. Faculty of Agriculture: Hasanudin University. Macassar.
 15. Sauvage, F. 1996. *The Marketing aspect Of Packaging*. Dalam. Bureau, G & Multon, J.L (pnyt). *Food Packaging. Technology*. Hlm. 33-41 VCH publisherbinc. Amerika.
 16. Sudarmadji, S., B. Haryono and Suhardi. 2003. *Analysis of Foodstuffs and Agriculture*. Gadjah Mada University Inter-University Center. Liberty. Yogyakarta.
 17. Suryani A., Erliza H. and Encep H. 2007. *Making Various Abon*. Self-Help Spreader. Jakarta.
 18. Tridiyani, A. 2012. *Changes in Quality of Shredded Marlin Fish (Istiophorus sp) Vacuum-Non-Vacuum Packaging at Various Storage Temperatures and Estimating Their Shelf Life*. Thesis. Faculty of Fisheries and Marine Sciences, Bogor Agricultural University. Bogor.
 19. Tribaditia, R. 2016. *Determination of Optimization Value Of Organoleptic Characteristics Aroma And Taste Of Corn Hair Tea Products With The Addition Of Lime And Honey*. *Journal of Agrosience*. 6(1) : 20-29.
 20. Widati, A.S., Mustakim and Sri Indriana. 2007. Effect of Liming Time on Water Content, Protein Content, Calcium Level, Growth Power and Organoleptic Quality of Cowhide Rambak Crackers. *Journal of Animal Products Science and Technology*. 2(1) : 47-56.
 21. Winarno, F.G. 2002. *Flavor for the Food Industry*. M-Brio Press. Jakarta