

## Physicochemical Parameters and Metals Distribution in Selected Shell Fishes Along the Oपुरo-Ama Creek in Rivers State, Niger Delta, Nigeria.

### ABSTRACT

Heavy metals concentration in selected shell fishes (*U. tangeri*, *C. amnicola*, *T. fuscatus*, *P. monodon*), Sediment and Water were collected from a fishing port along the Oपुरo-ama in Creek, Asari Toru Local Government area of Rivers State, Nigeria and were analysed for Manganese (Mn), Iron (Fe), Cobalt (Co) and Zinc (Zn) using *Atomic Absorption Spectrophotometer*. The order of metal accumulation in these shellfishes, sediment and water were; *C. amnicola* (Fe > Zn > Cu > Co > Mn); *U. tangeri* (Fe > Zn > Cu > Co > Mn); *T. fuscatus* (Fe > Cu > Zn > Co > Mn); *P. monodon* (Fe > Co > Zn > Cu = Mn); Sediment (Fe > Zn > Co > Cu > Mn); Water (Fe > Cu > Mn > Co = Zn). There was statistical difference ( $p < 0.05$ ) in metals concentration in the soft tissues in all the shellfish. From the results, the concentration of the metals in the tissues of all the shell fishes, the interstitial water and the sediment sampled from the creek were within the permissible limit recommended by the [10; 13 and 9] except for Fe which was above the recommended limit. The physicochemical parameters of the interstitial water such as pH, Temperature, Conductivity, Total Dissolved Solids (TSS) Biological Oxygen Demand (BOD), Dissolved Oxygen (DO), and Salinity were within the acceptable levels in the guidelines for drinking water by [10; 13 and 9]. Although Chemical Oxygen Demand (COD) and the Total Suspended Solids (TSS) were slightly higher than the recommended limit of these metals by [10]. This indicates that if these metals are not checked, they may increase the potential for bioaccumulation in the shellfishes and the creek as well. Therefore, there is need for regular monitoring of this creek to avoid future deterioration

**Keywords:** Physicochemical parameters, heavy metals, Shellfishes, Oपुरo-ama creek.

## 1. INTRODUCTION


Environmental pollution, natural or man-made is assuming enormous proportions, increasing considerably by the day, with the advent of civilization, industrialization and urbanization [1]. Most of the world's air, water and land are now mostly contaminated by chemical wastes from domestic, industrial processes, including those of crude oil and gas [2] Consequently, the aquatic ecosystems act as one of the major receptacle ecosystems for various contaminants generated through the unregulated release of effluents from mines, smelters, industries, excessive usage of agrochemicals, and from aerial deposition [3]. In Nigeria, the input of environmental pollutants in aquatic systems is a common phenomenon. It is a common believe that aquatic systems have an unlimited capacity to absorb increasing amount and variety of waste substances and energy from our civilization [4]. Aquatic environment pollution as a result of heavy metals has been a critical issue all over the world, this is because they are indestructible and most of them have toxic effects on organisms [5]. Fin and shell fishes have been widely used as bio-indicators to monitor heavy metals concentrations in the coastal environment, due to their wide range of distribution, and also their important position in the food chain. Among the different kinds of pollutants, heavy metal pollution needs to be dealt with on top priority considering both its fast rate of accumulation in the aquatic medium and the impact of its toxicity on the organisms along the food chain and its biomagnification impacts [6]. Giving the potential environmental exposure of the Opuro-ama creek to heavy metal contamination by virtue of its position as recipient creek to many other tributaries in the Niger Delta region, it is pertinent to evaluate the likely effect of its pollution on the inhabitants and communities which depend on it for their livelihood in terms of fishery resources and other services.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The study was carried out from three stations along the Opuro-ama creeks in Asari-Toru Local Government Area of Rivers State ((Figure 1). The Creeks is a joint tributary of the Sombraro Estuary, Located Southeast of the Niger Delta between longitude N04<sup>0</sup> 48' 14.0" and latitude E006<sup>0</sup> 50' 16. It shears boundaries with Sa-ama and Abalama Creek. These creeks system consists of the main channel and associated feeder creeks linking Opuro-ama, Sa-ama, Abalama, Te-ama, Sangama Degema, Krakrama and other riparian communities. Lutjanidae, Clupeidae, Cichlidae and Claroteidae are among the main fish family in this ecosystem but the Claroteidae and Cichlidae most common are the Claroteidae (silver catfish) (tilapias) [7]. Whereas the tidal mudflats species include the gobies, periwinkles, crabs, mudskippers, only to mention a few.

The mangrove vegetation of the area comprises *Rhizophora racemosa*, *R. mangle* and *R. harrisonii* (red mangrove). Other mangrove species in the area are *Avicennia germinans* (white mangrove), *Laguncularia racemosa* (black mangrove) and *Conocarpus erectus* (button wood). There is also presence of mangrove associates such as *Acrostichum aureum* (mangrove fern) and *Paspalum vaginatum* (mangrove sedge). Other human threats to these creeks mangrove swamp include artisanal channelization, Illegal refining and bunkering activities, sand dredging and deliberate cutting of mangroves at the community landing corridors for 'security reasons etc.



**Figure 1:** Showing the Sampling Area

## 2.2 Collection of Samples

A total of twenty representative samples per station at each sampling campaign were obtained within a sampling period (January – June, 2018) to provide for statistical variation. The healthy species were collected from the creeks.

## 2.3 Shell Fish

The swimming crabs (*Callinectes amnicola*) was caught using a drag net while the Mud flat crab (*Uca tangeri*) was collected randomly from their hide out holes in the mud flat using creek using a crab trap. Periwinkle (*Tympanotonus fuscatus*) was handpicked from the mud flat too. while the Tiger shrimp (*Penaeus monodon*) was caught using a drag net. Sampling was carried out from three different sample stations across the creek. The samples collected was persevered in an ice pack before transported to the laboratory for heavy metal concentration using atomic absorption Spectrometer.

## **2.4 Sediment Sample**

A bottom sediments samples were collected in a composite form from three different stations using an Ekman grab sampler and kept in a plastic container which had been previously treated with 10% nitric acid for 24 hours and rinsed with de-ionized water.

## **2.5 Water Sample**

Three pits were dogged randomly on the intertidal flat of each sample station and allow for the accumulation of the interstitial water. After fifteen minutes, the interstitial water would be collected in-situ using water collection bottle.

## **2.6 Determination of Heavy Metals**

The Cobalt (Co), Copper (Cu), Iron (Fe), Zinc (Zn), Magnesium (Mn) were determined using the Atomic Absorption Spectrophotometer (AAS) method.

## **2.7 Determination of physicochemical parameters**

The interstitial water samples were collected using Schott glass bottles. The pH, interstitial water Temperature, Salinity, Conductivity, Total Suspended Solids (TSS), Total Dissolved Solids (TDS) of the water were measured using an in-situ Handheld Multi-meter (Milwaukee Model pH600 and Laboratory Benchtop meter 860033-model). Dissolved Oxygen (DO) was measured using a Milwaukee Dissolved oxygen meter (MW 600 Model). Biochemical Oxygen Demand (BOD) was determined by the 5-day BOD test (APHA, 2005). Chemical Oxygen Demand (COD) was determined using the Closed Reflux Method 5220C with a higher concentration of potassium dichromate solution. All parameter determined were method recommended by APHA 2340C (1995) standards.

## **2.8 Statistical Analysis**

Data will be analyzed using a one-way ANOVA and Duncan's multiple range and results will be

tested for statistically significant differences at the 0.05.

### 3. RESULTS AND DISCUSSION

Results of physicochemical parameters, metals distribution and bioaccumulation patterns from the study are presented in Table 1 to 7 and Figures. 2-6. The bioaccumulation factors are presented in Table 3 and 4.

**Table 1: Physicochemical Quality of the Interstitial Water Samples.**

Parameters	Min-Max	Mean -SE	WHO (2011)	DPR (2002)	EPA (2017)
Temperature (°C)	29.70 - 30	30.0±.057	24.8-30	22.32-25	27 - 30
pH	6.40 - 6.80	6.57±0.06	6.5-8.5	6.5-8.5	6.5 - 8.5
DO (Mgl <sup>-1</sup> )	3.0 - 3.2	3.2±0.058	6	5	7.5
Salinity (ppt)	11.50 - 11.70	11.69±0.09	120	600	-
BOD (Mgl <sup>-1</sup> )	0.53 - 0.87	0.68±0.033	10	10	5 - 7
Conductivity (µS/ cm <sup>-1</sup> )	19.6 - 22.7	21.62±0.06	400	-	200 - 1000
Total Dissolved Solids (µS/cm)	16.0 - 17.43	16.61±0.10	2000	2000	1000 - 2000
Total Suspended Solids (µS/cm)	94.0 - 140.7	111±0.58	500-1000	30	30
Chemical Oxygen Demand (µS/cm)	52.0 - 73	64.6±0.88	40	10	-

**Table 2: Heavy Metals Composition in Some Selected Shell Fishes along the Opuro-ama in Creek**

Heavy Metals (µg/g)	<i>C. amnicola</i>	<i>U. tangeri</i>	<i>T. fuscatus</i>	<i>P. monodon</i>	Sediment	Water	WHO (2011)	EPA (2017)
Cu	0.34±0.01 <sup>b</sup>	0.35±0.02 <sup>b</sup>	0.43±0.02 <sup>a</sup>	0.02±0.02 <sup>c</sup>	0.04±0.02 <sup>c</sup>	0.03±0.02 <sup>c</sup>	2.0	3.0
Zn	0.43±0.02 <sup>a</sup>	0.45±0.02 <sup>a</sup>	0.40±0.02 <sup>a</sup>	0.04±0.02 <sup>c</sup>	0.12±0.02 <sup>b</sup>	0.01±0.02 <sup>c</sup>	3.0	3.0
Co	0.06±0.02 <sup>a</sup>	0.05±0.02 <sup>b</sup>	0.06±0.02 <sup>a</sup>	0.05±0.02 <sup>b</sup>	0.06±0.02 <sup>a</sup>	0.01±0.01 <sup>c</sup>	-	-
Fe	2.77±0.02 <sup>cd</sup>	3.20±0.02 <sup>b</sup>	2.80±0.02 <sup>c</sup>	3.10±0.02 <sup>b</sup>	3.70±0.02 <sup>a</sup>	2.57±0.02 <sup>d</sup>	0.1	0.5
Mn	0.02±0.01 <sup>a</sup>	0.02±0.02 <sup>a</sup>	0.02±0.01 <sup>a</sup>	0.02±0.01 <sup>a</sup>	0.02±0.01 <sup>b</sup>	0.02±0.01 <sup>a</sup>	0.05	0.02

\* In each column, mean with a common letter are not significantly different (P>0.05).

**Table 3: Patterns of heavy metals concentrations in the Shellfishes, Sediment and water**

Medium	metal accumulation pattern
<i>U. tangeri</i>	Fe>Zn>Cu>Co>Mn
<i>T. fuscatus</i>	Fe>Cu>Zn>Co>Mn
<i>P. monodon</i>	Fe>Co>Zn>Cu=Mn
<i>C. amnicola</i>	Fe>Zn>Cu>Co>Mn
Sediment	Fe>Zn> Co> Cu>Mn
Water	Fe>Cu>Mn>Co=Zn

**Table 4:** Concentration factors of metals in the shell fishes with reference to water.

Medium	Cu	Zn	Co	Fe	Mn
<i>C. amnicola/water</i>	11.3	43.0	1.0	1.1	1.0
<i>U. tangeri/water</i>	11.6	45.0	5.0	1.25	1.0
<i>T. fuscatus/water</i>	14.3	40.0	6.0	1.08	1.1
<i>P. monodon/water</i>	0.7	4.0	5.0	1.21	1.0

**NB:** Concentration in (mg kg<sup>-1</sup>)

**Table 5:** Bioaccumulation patterns of heavy metals in the shell fishes with reference to water.

Medium	metal accumulation pattern
<i>C. amnicola</i>	Zn>Cu> Fe>Co=Mn
<i>U. tangeri</i>	Zn>Cu>Co> Fe>Mn
<i>T. fuscatus</i>	Zn>Cu>Co>Fe>Mn
<i>P. monodon</i>	Co>Zn> Fe>Cu=Mn

**Table 6:** Concentration factors of metals in the shell fishes with reference to sediment.

Medium	Cu	Zn	Co	Fe	Mn
<i>C. amnicola/ sediment</i>	8.5	3.58	1.0	0.75	1.0
<i>U. tangeri/ sediment</i>	8.75	3.75	0.83	0.86	1.0
<i>T. fuscatus/ sediment</i>	10.75	3.3	1.0	0.75	1.0
<i>P. monodon/ sediment</i>	0.5	0.3	0.83	1.2	1.0

**NB:** Concentration in (mg kg<sup>-1</sup>)

**Table 7:** Bioaccumulation patterns of heavy metals in the shell fishes with reference to sediment.

Medium	metal accumulation pattern
<i>C. amnicola</i>	Cu>Zn>Co=Mn>Fe
<i>U. tangeri</i>	Cu>Zn>Mn>Fe>Co
<i>T. fuscatus</i>	Cu>Zn>Co = Mn>Fe
<i>P. monodon</i>	Fe>Mn > Co>Cu>Zn




Fig 2: Comparative values of Cu in Some Selected Shell Fishes from a fishing Port along the Opuro-ama in Creek




Fig 3:Comparative values of Zn in Some Selected Shell Fishes from a fishing port along the Oपुरo-ama in Creek




Fig 4:Comparative values of Co in Some Selected Shell Fishes from a fishing Port along the Oपुरo-ama in Creek




Fig 5:Comparative values of Fe in Some Selected Shell Fishes from a fishing Port along the Oपुरo-ama in Creek




Fig 6: Comparative values of Mn in Some Selected Shell Fishes from a fishing port along the Oपुरo-ama in Creek.

### 3.1 Physicochemical Parameter of the Interstitial Water

The physicochemical parameters of the interstitial water samples collected from the Oपुरo-ama creek during the study period are shown in Tables 1. The mean pH concentration ranges from 6.40–6.80 with a mean value of  $6.57 \pm 0.06$ . The overall water temperature observed at the period of study ranged from 29.70–30(°C) with a mean value of  $30.0 \pm 0.057$ (°C) while the conductivity was between 19.6–22.7 ( $\mu\text{S}/\text{cm}$ ) with a mean value of  $21.62 \pm 0.06$ ( $\mu\text{S}/\text{cm}$ ) and TDS concentration ranged from 16.0–17.43( $\mu\text{S}/\text{cm}$ ) with a mean value  $16.61 \pm 0.10$ ( $\mu\text{S}/\text{cm}$ ). The Total Dissolved Solids (TSS) concentration ranged from 94.0–140.7( $\mu\text{S}/\text{cm}$ ) with a mean value of  $111 \pm 0.58$  Solids ( $\mu\text{S}/\text{cm}$ ) while the mean values of BOD observed was between 0.53–0.87 ( $\text{Mg l}^{-1}$ ) mean value of  $0.68 \pm 0.033$ ( $\text{Mg l}^{-1}$ ). The low DO values were between 3.0–3.2( $\text{Mg l}^{-1}$ ) with a mean value of  $3.2 \pm 0.058$ ( $\text{Mg l}^{-1}$ ) and the Salinity values observed was between 11.50 –11.70 (ppt) with a mean value of  $11.69 \pm 0.09$ (ppt) during the study. The Chemical Oxygen Demand (COD) values range from 52.0–73( $\mu\text{S}/\text{cm}$ ) with a mean value of  $64.6 \pm 0.88$ ( $\mu\text{S}/\text{cm}$ ).

The mean temperature value measured was within the range of 27–30°C recommended [8]. This was also within the acceptable range by [9 and 10]. This was also similar to results reported by [11] and [12] in of Lagos Lagoon, Lagos, Nigeria. The overall mean pH measured was 6.57

within the range of 6.5 – 8.5 as being ideal for supporting aquatic life by [9; 10 and 13]. [14] reported similar results from Ogunpa River at Bodija. The reported in the study, which is within the acceptable limit for freshwater bodies, was similar to those reported by [1] along the Tin Can Island in Lagos, Nigeria. The mean DO measure in station one (3.2 mg/l) was below the recommended level of 6-7.5mg/L for the culture of freshwater organisms. This could be due to the high organic contents from anthropogenic activities peculiar to the study area. Similar value was reported by [15] in the interstitial water quality of soft-bottom flats at Bodo creek. the mean DO value was within the recommended levels [12]. The BOD was also within the recommended values of ideal to supporting aquatic life by [9 and 13]. This result did not agree with the assertion by [16] who reported that the higher the concentration of BOD, the lower the concentration of DO. Total Dissolved Solids ( $\mu\text{S}/\text{cm}$ ), Conductivity ( $\mu\text{S}/\text{cm}^{-1}$ ) and Salinity (ppt) were also within the permissible limit of [9; 10 and 13]. Similar observations were reported in the adjoining creeks and lagoons in Lagos and its environments by [17 and 18]. Although Chemical Oxygen Demand (COD) and the Total Suspended Solids (TSS) were slightly higher than the recommended limit by [9 and 10]. The high COD values were indication of the presence of organic and inorganic pollutants, respectively. This could be attributed to the domestic sewage from various activities of human residing around the studied area, reception of effluents containing [19]. Similar result was reported by [20] in southeast coast of India.

### **3.2 Heavy Metals Composition**

Tables 2 Shows the heavy metals concentration in some selected Shellfishes samples from Opuro-ama Creek. The level of Coper in the shellfishes was highest in *T. fuscatus* ( $0.43\pm 0.02$  mg/l) followed by *U. tangeri* ( $0.35\pm 0.02$  mg/l) and *C. amnicola* ( $0.34\pm 0.01$  mg/l) and the least was observed in *P. monodon* ( $0.02\pm 0.02$  mg/l). There was significant variation ( $p < 0.05$ ) across

the different medium. Although, no significant difference  $p > 0.05$  was observed between *U. tangeri* and *C. amnicola*; and between surface water ( $0.03 \pm 0.02$  mg/l) and Sediment ( $0.04 \pm 0.02$  mg/l). *U. tangeri* ( $0.45 \pm 0.02$  mg/l) recorded the highest level of Zinc concentration while *P. monodon* ( $0.02 \pm 0.02$  mg/l) recorded the least. There was significant difference  $p < 0.05$  between *P. monodon* and the other shellfishes, although no significant difference  $p > 0.05$  was observed between *U. tangeri*, *T. fuscatus* and *C. amnicola*. The highest values of Cobalt were recorded in *C. amnicola* ( $0.06 \pm 0.02$  mg/l), *T. fuscatus* ( $0.06 \pm 0.01$  mg/l) and sediment ( $0.06 \pm 0.01$  mg/l) respectively while the least value was recorded in the water ( $0.01 \pm 0.01$  mg/l). *P. monodon*, Water and *U. tangeri* were significant difference  $p < 0.05$  from *C. amnicola*, *T. fuscatus* and sediment. The concentration of Iron in the shellfishes was highest in sediment ( $3.70 \pm 0.02$  mg/l) followed by *U. tangeri* ( $3.5 \pm 0.02$  mg/l) and the least was recorded in water ( $2.57 \pm 0.02$  mg/l) and there was significant difference  $p < 0.05$  across the different medium. The highest concentration of Manganese was recorded in *U. tangeri* ( $0.03 \pm 0.02$  mg/l) while the least were recorded in *C. amnicola*, *T. fuscatus*, *P. monodon*, sediment and water ( $0.02 \pm 0.01$  mg/l). There was no significant difference  $p > 0.05$  was observed across the different media.

From the study, the heavy metals recorded varied from species to species. The species *T. fuscatus* and *T. coronata* were found to have higher values of Cu while the least was recorded in the interstitial water. This assertion agrees with the findings of [21], who compared heavy metals in *Anadara granosa* and *periwinkle* in Malaysia. Similar result was reported by [4] attributed higher heavy metals in periwinkle to its shape and life history of some commercially important shell in Port-Harcourt. Mean values of Zinc recorded in this study were highest *U. tangeri* and least in the interstitial water. The values from all the shell fish species including the sediment and water were within the acceptable limits of [9 and 13]. The values from this study were lower when

compared to the values obtained in some shell fishes from lagoon systems in the Southern Gulf of Mexico which were higher than the values of Zinc obtained in some shell fishes considered by [22]. The high concentration on cobalt which was observed *C. amnicola*, *T. fuscatus* and the sediment could be attributed to the water chemistry, duration of exposure to contaminants to the sediment leading to the increase in the concentrations of contaminants in their habitat, feeding habits or season of sampling [23]. High concentration on cobalt which was observed *C. amnicola*, *T. fuscatus* could be related to their specific and feeding habits and the bioconcentration capacity of each species [24]. Iron (Fe) levels observed in these shell fishes, sediment and water obtained in this work were all above acceptable limits of acceptable limits of for human consumption of 0.1 and 0.5µg/g as reported by [9 and 13]. The result was not in accordance with [25], who stated that the median iron concentration in rivers was reported to be 0.7 mg/litre. He also stated that in anaerobic groundwater where iron is in the form of iron (II), concentrations will usually be 0.5–10 mg/litre. Concentrations of iron in drinking-water are normally between 0.1-0.5mg/litre [9 and 13]. Although higher in countries where various iron salts are used as coagulating agents in water-treatment plants and where cast iron, steel, and galvanized iron pipes are used for water distribution [26].The accumulations of metals were generally found to be different shell fish species were higher than the value recorded in the sediment and the interstitial water except for manganese was not significantly different across board. The concentration on manganese which was observed in *C. amnicola*, *T. fuscatus*, *T. fuscatus*, *P. monodon*, sediment and the water were within the acceptable limits of [9 and 13].

### **3.3 Concentration factors of Heavy Metals in Sediments and Water**

The shell fishes have been reported to be good accumulators of organic and inorganic pollutants which are later transferred to other animals including man through the food web chain. The

concentration factors of metals in shell fishes accumulation of metals was in the order of magnitude; *C. amnicola*: Zn>Cu> Fe>Co>Mn, *U. tangeri*: Zn>Cu>Co>Fe>Mn, *T. fuscatus*: Zn>Cu>Co>Mn>Fe, *P. monodon*: Co> Zn >Fe>Mn> Cu. Zn and Cu were better accumulated in *C. amnicola*, *U. tangeri* and *T. fuscatus*. However, some of the metals such as Fe and Co were better bioaccumulated in *P. monodon*.

Bioaccumulation of these metals was generally higher in *C. amnicola*, *U. tangeri* and *T. fuscatus* when water was the medium of evaluation with metals such as Zn and Cu. The magnitude of bioaccumulating in these shellfish may be higher in the organisms as a result of interaction of the fishes with higher metal concentrations in water which typifies their filter feeding habit. The order of magnitude of metal accumulation in these shellfishes from sediment was in the sequence; *C. amnicola*: Cu > Zn > Co = Mn >Fe, *U. tangeri*: Cu > Zn > Mn > Fe > Co, *T. fuscatus*: Cu > Zn > Co = Mn > Fe, *P. monodon*: Fe > Mn > Co > Cu > Zn. The bioaccumulation patterns observed in the present study shows that even in similar media, different organisms may accumulate of different contaminants and there may be specific factors influencing the selective bioaccumulation of these metals in the organisms. However, according to [27], the rate of bioaccumulation of heavy metals in aquatic organisms depends on the ability of the organisms to digest different metals and the concentration of the metals in the river. The Bioaccumulation factor may have to do with the concentration of the heavy metal in the surrounding sediments as well as their feeding habits [28]. Other factors such as lipid content in the tissue of the fishes, age of fish and mode of feeding may be significant factors that affect the accumulation of these heavy metals in shellfishes.

Similar studies have reported similar bioaccumulation patterns of metals in different organisms in the Bonny/New Calabar River Estuary [27; 29; 28] and elsewhere in the Niger Delta [1]. Heavy

metals have multiple effects in aquatic ecosystems depending on the oxidation state, formation of complexes and biotransformation of elemental species. Health effects of heavy metals such as Fe, Mn, Co, Cu and Zn in humans have been demonstrated in acute toxicity, neurotoxicity and nephrotoxicity [7]. The concentration factors result of heavy metals revealed in this study shows that there was bioaccumulation in the different media but at a low rate. Bioaccumulation Factor value of Zn and Cu were better accumulated in *C. amnicola*, *U. tangeri* and *T. fuscatus*. However, some of the metals such as Fe and Co were better bioaccumulated in *P. monodon*. The study indicated that sediment accumulated more heavy metals than water. This is in line with results reported works by [30]. They stated that in most cases, sediment is reported to be the main sink of metals, with more than 99% of the total amount of metals present in the aquatic body (Eja *et al.*, (2003). This value was not in line with reported by [32] in *Tympanotonus fuscatus var radula* obtained from the upper reaches of the Bonny estuary.

The concentration factors in some of the shellfish species were  $<1$  for all the studied heavy metal. Likewise, heavy metal bioaccumulation from water (BAF) showed  $<1$  for all except Zn ( $8.09$  to  $11.71 \text{ mg kg}^{-1}$ ) and as ( $2.00 \text{ mg kg}^{-1}$ ). This study observed that these trace metals were accumulated in the tissue of the fish more than they are actually present in the water. This singular potential should qualify the mudskipper to be categorized in the league of some well-known bioindicator species like the bivalves, crabs and shrimps identified as standard bioindicators of aquatic pollution owing to their capability to bioaccumulate and bioconcentrate organic pollutants in tissues in addition to target organs at levels higher than background concentrations [33]. Although tissue may not be an active site for bio-accumulation, the findings in this study may imply that at chronic exposures in minute concentrations, fish tissues could concentrate heavy metals that exceed the permissible limits for human consumption with

severe health implications. Zn concentration was below [34] permissible limit of 40 mg kg<sup>-1</sup> in the muscle. The tissue of fish is the edible part, and the concentrations of the heavy metal in it are of health concern. The selective accumulation of some metals as observed in this study implies that the affinity for the accumulation of heavy metal in tissues varies from fish to fish and may be metal-specific. Research on temporary variation in *Periophthalmus papillion* obtained from Azuabe Creek in the upper Bonny Estuary observed heavy metal concentrations in the flesh of mudskippers to levels above [34] permissible limits and concluded that mudskippers have the ability to bioaccumulate and bio-magnify metal pollutants without physical stress. These make them potentially dangerous to the consumers, and by extension, all other tolerant commercial species fished from these water bodies.

Some reported effect of these metals (Fe, Zn, Cu, Co, Mn) in aquatic environments include tissue damage, genotoxicity and growth reduction. Mollusks and crustaceans are more sensitive than other organisms [35]. Although the results of surface water concentrations of heavy metals observed in the present study agrees with the general opinion of low-level heavy metal concentrations in surface water in the study area and Niger Delta [36] metals such as Fe was higher than stipulated limits by [9 and 13] and require continuous monitoring to detect uncertain increases as a result of anthropogenic activities and avert possible health implications of these metals on consumers of the seafood from the water body in this study area.

#### **4. Conclusion**

The results showed that the concentration of the metals in the tissues of all the shell fishes, the interstitial water and the sediment sampled from the creek were within the permissible limit recommended by [13 and 9]. Except for Fe which was above the recommended limit. There was an overall statistical difference ( $p < 0.05$ ) in metals concentration in the soft tissues in all the

shellfish. The physicochemical parameters of the interstitial water were also within the acceptable levels in the guidelines for drinking water by Department of Petroleum Resources [10; 13 and 9].

## **5. Recommendation**

There is need for regular monitoring of the metallic level in this creek to avoid future deteriorations should therefore be periodically checked to avoid increase in the potential for bioaccumulation of these metals in the shellfishes and the creek water as well.

## **Compliance With Ethical Standards**

The research complied with all ethical standards.

## **Reference**

1. Davies IC, Nkeeh DK, Akoko S, Ariri IP. Spatial Variation of Heavy Metals in Blue Crab *Calinectes amnicola* Harvested from The Water Front of Some Fishing Communities in River State, Nigeria. *International Research Journal of Applied Sciences, Engineering and Technology*.2021;7(11):33-49
2. Akankali JA, Davies IC. Analysis of heavy metals concentration in different media of Iwofe Creek, Niger Delta, Nigeria; *International Journal of Scientific and Research Publications*, 2020;10(04): 2250-3153.
3. Davies IC, Okonkwo SE. Analyses of Some Heavy Metal and Nutrients of Water Samples from Ajegunle Creek in Lagos. *International Journal of Environment and Pollution Research*.2021; Vol.9, No.1 pp.7-26.
4. Akinrotimi OA, Amachree D, Gentle CG. Metal concentrations in some commercially important shell in Port Harcourt, Rivers State, Nigeria.*International Journal of Fisheries and Aquatic Studies*. 2019;7(2): 134-138.
5. Zhou YW, Peng YS, Li XL, Chen GZ. Accumulation and partitioning of heavy metals in mangrove rhizosphere sediments. *Environmental Earth Sciences*, 2011;64(3), 799-807.
6. Okere MC, Davies IC, Onyena AP. Variation of The Physicochemical Parameters, Nutrients and Some Selected Heavy Metals Around the Waters of the Tincan Island in Lagos, Nigeria. *British Journal of Environmental Sciences*,2021;9(4), 1-17.
7. Davies IC, Ekperusi AO. Evaluation of Heavy Metal Concentrations in Water, Sediment and Fishes of New Calabar River in Southern Nigeria. *Journal of Limnology and Freshwater*

*Fisheries Research*, (2021);7(3): 207-218. doi: 10.17216/LimnoFish.816030.

8. FEPA (Federal Environmental Protection Agency) National Guidelines and Standards for Industrial Effluents and Water Quality Tests FEPA (Nigeria) Official Gazette, Nigeria 1991.

9. World Health Organization. Guidelines for drinking-water quality. *WHO chronicle*, 2011;38(4), 104-108.

11. Tosin OV, Oloyede TL, Iortim M. The effect of water renewal on growth of *Clarias gariepinus* fingerlings. *Croatian Journal of Fisheries: Ribarstvo*, 2016;74(1), 25-29.

12. Lawson EO. Physicochemical parameters and heavy metal contents of water from the Mangrove Swamps of Lagos Lagoon, Lagos, Nigeria. *Advances in biological research*, 2011;5(1), 8-21.

14. Okonkwo, S. E; Davies I. C; Okere M.C. (2021): Assessment of Physicochemical Characteristics and Phytoplankton of a Polluted Tidal Creek In Ajegunle, Lagos. *British Journal of Environmental Sciences* 9(1): 51-69.

15. Zabbey N. Spatial and temporal variability in interstitial water quality of soft-bottom flats at Bodo creek, eastern lower Niger Delta, Nigeria. *Tropical Freshwater Biology*, 2012;21(1), 83.

16. Atobatele OE, Morenikeji OA, Ugwumba OA. Spatial variation in physical and chemical parameters of benthic macroinvertebrate fauna of River Ogunpa, Ibadan. *The Zoologist*, 2005;3, 58-67.

18. Onyema IC, Popoola RT. The physico-chemical characteristics, chlorophyll a levels and phytoplankton dynamics of the east mole area of the Lagos Harbour, Lagos. *Journal of Asian Scientific Research*, 2013;3(10), 995-1010.

19. Okere MC, Davies IC, Okonkwo SE. Seasonal Variation of the Hydro-Environmental Factors and Phytoplankton Community around Waters in Tincan Island, Lagos State, Nigeria. *Journal of Applied Sciences & Environmental management*. 2020;24(10) 1739-1746.

20. Eswari YNK, Ramanibai R. Estuarine copepod abundance and diversity in relation to environmental variables, southeast coast of India. *Journal of the Marine Biological Association of India*, 2004;46(1), 10-20.

21. Min KS, Kee KP, Tien SK. Proximate Analysis and Heavy Metal Concentrations of Tissues of Cockles (*Anadara granosa*) from Several Cockle Farms in Peninsular Malaysia; Analisis Proksimat dan Logam Berat dalam Tisu Kerang (*Anadara granosa*) di Beberapa Tapak Akuakultur di Semenanjung Malaysia. *Sains Malaysiana*, 2011;40(2):139-146.

22. Mohammad M, Sary AA, Khodadadi M. Determination of heavy metals in two barbs, *Barbus grypus* and *Barbus xanthopterus* in Karoon and Dez Rivers, Khoozestan, Iran. *Bull Environment Contain Toxicology*. 2011; 87:158-162.

23. Ross SM. Toxic Metals in soil – plant systems. Wiley, Chichester. UK. 2014;653pp.
24. Tuzen M. Determination of heavy metal in fish samples of middle black sea (Turkey) by graphite furnace atomic absorption spectrometry. *Food Chemistry*. 2003;80:119-173.
25. Vitamin, H. (2001). Mineral Requirements. *Report of a joint FAO/WHO expert consultation Bangkok, Thailand*, 181-94.
26. National Research Council (NRC),1979. Committee on Medical, Biologic Effects of Environmental Pollutants. Subcommittee on Iron, Assembly of Life Sciences (US). Subcommittee on Iron, National Research Council, National Research Council (États-Unis). Committee on Medical, Biologic Effects of Environment, ... & Biologic Effects of Environmental Pollutants. *Iron*. University Park Press.
27. Babatunde B, Bsikoki FD, Onojake MC, Akpiri RU, Akpuloma D. Heavy Metal Profiles In Various Matrices of The Bonny/New Calabar River Estuary, Niger Delta, Nigeria *Global Journal Of Environmental Sciences* 2013;Vol. 12:1-11.
28. Akankali JA, Davies IC, Oyema IC. Evaluation of some physicochemical Parameters of the Tin Can Island Creek, Lagos, Nigeria, *Nigerian Journal of Fisheries*, 2019;16(2):1783-1793.
29. Davies, I. C., Agarin, O. J., and Onoja C. R. (2021). Study On Heavy Metals Levels and Some Physicochemical Parameters of a Polluted Creek Along the Tin Can Island in Lagos. *International Journal of Environment and Pollution Research*.9(2), 25-39.
30. Jamabo N, Chinda A. Aspects of the ecology of *Tympanotonusfuscatus var fuscatus* (Linnaeus, 1758) in the mangrove swamps of the upper Bonny River, Niger Delta, Nigeria. *Curr. Res. J. Biol Sci*. 2010;2, 42-47.
31. WHO. Guidelines for drinking-water quality (electronic resource). Incorporating first addendum - 3rd ed. Volume 1. Recommendations. World Health Organization, Geneva, 2006.
32. Onojake MC, Frank O. Assessment of heavy metals in a soil contaminated by oil spill: A case study in Nigeria, *Chemistry and Ecology*,2012;DOI:10.1080/02757540.2012.717619.
33. Chindah AC, Braide AS, Sibeudu OC. Distribution of hydrocarbons and heavy metals in sediment and a crustacean (*Penaeusnotialis*) from the Bonny River/New Calabar River Estuary, Niger Delta. *African Journal Environmental Assessment and Management*. 2004;(9): 1-17.