

### **G and P Genotypes of Group A Rotavirus in Diarrheic Calves in Costa Rica**

#### **ABSTRACT**

Rotavirus infections are the most common causes of infectious diarrhea in young children and animals worldwide. We performed a genotyping of rotavirus strains by RT-PCR for G (VP7) and P (VP4), in diarrhea samples from calves, in several locations in Costa Rica. Combinations of genotypes G (VP7) and P (VP4) were identified in cattle: 4 (26%) corresponding to G8 P [11], 4 (24%) to G6P [11], 3 (19%) to G8P [7], 2 (13%) to G10P [1], 3 (19%) to G10P [11], and 1 (6%) to G8P [1] and mixed infections. This research represents the first report of bovine rotavirus strains (calves) in Costa Rica and in the Central American region.

**Key words:** rotavirus, diarrhea, calves, VP7 and VP4.

Rotavirus is considered the main etiological agent in diarrhea in young animals and children around the world [1].

Group A Rotaviruses are classified using a binary system, according to the genetic and antigenic variation of two surface proteins (VP7 and VP4). So far, 36 types G (VP7) and at least 51 types P (VP4) of rotavirus group A circulating in humans and animals [2] have been reported, particularly in cattle, 14 different G (VP7) genotypes (G1 -G6, G8, G10-G12, G15, G18, G21, and G24 and 11 P

(VP4) genotypes (P [1], P [3], P [5] -P [7], P [11], P [14], P [17], P [21], P [29], and P [33]) [3, 4] that affect animal populations causing great economic losses for livestock producers, due to vaccination expenses, treatment, weight loss of affected animals and mortality.

Risk factors that contribute to the appearance of diarrhea caused by rotavirus are failures in the transfer of passive immunity especially observed in heifers, presence of seasonal diarrhea, lack of hygiene in artificial rearing systems, leading to economic losses for livestock producers, due to vaccination, treatment and weight loss of affected animals and mortality regions [5].

Knowledge of the epidemiology of rotavirus is essential for the control of this disease; especially since it has been observed that it presents differences between developed and developing countries, as well as between temperate and tropical climate regions [6].

In Costa Rica, in the Central American region and in the Dominican Republic we have carried out characterizations of rotavirus strains in children, where we report common and unusual genotypes [7].

This research represents the first genotyping study of rotavirus G (VP7) and P (VP4) strains detected in cattle in Costa Rica and in the Central American region.

Samples of feces from pigs under 1 month of age were taken from 17 dairy farms (bovine-calves) located in Río Cuarto de Grecia, Pital, Poás, Zarcero, Sucre and Pacayas in 2008 in Costa Rica. All calf stool samples were analyzed in the virology laboratory, at the School of Veterinary Medicine at the National University of Costa Rica (UNA), to determine the presence of rotavirus using the immunochromatography technique (CorisBioConcept, Belgium) and the RT-PCR technique (VP7) and (VP4) described by Genstch JR et al [8] and Gouvea V et al [9].

From the 200 samples (calves feces), a total of 38 samples were ELISA positive for rotavirus and 25 samples were sent for characterization of the G (VP7) and P (VP4) genotype.

By using the RT-PCR technique, the combinations of genotypes G (VP7) and P (VP4) were identified in cattle: 4 (26%) corresponding to G8 P [11], 4 (24%) to G6 P [11], 3 (19%) to G8 P [7], 2 (13%) to G10 P [1], 3 (19%) to G10P [11], and 1 (6%) to G8P [1]. In 3 samples, it was not

possible to identify the G (VP7) genotype with the primers used. Mixed infections were found in 6 samples (Table 1).

The distribution of genotypes G (VP7) and P (VP4) of Group A bovine rotavirus in the dairy farms of the different locations in Costa Rica, were different. Namely, G10 was the most prevalent genotype in the Poás, Zarcero and Sucre area, while the G6 genotype prevailed in the Río Cuarto de Grecia, Poás and Pital de San Carlos areas. Finally, G8 was identified mainly in the Pacayas, Poás, Cedral de San Carlos and Zarcero areas. In the P (VP4) types, P [11] was identified as the main genotype in Río Cuarto de Grecia, Pital de San Carlos, Poás, Cedral de San Carlos and Pacayas, P [7] was found only in Río Cuarto de Grecia, whereas P [1] was identified in Zarcero and Sucre. Mixed infections were identified in Pacayas and Poás (Table 2).

There are reports in the literature of infections in cattle caused by various genotypes of Group A Rotavirus such as G1, G2, G3, G6, G8, G10 and G11 [4, 10, 11, 12]. In the present study, the (VP7) G6, G8 and G10 genotypes were identified, with the G10 genotype prevailing, which was identified in 10 samples, followed by G6 and G8 identified in 8 and 7 fecal samples, respectively. Our findings coincide with reports in Brazil, Argentina, and other countries in the world where high frequencies of these genotypes were reported and considered common of epidemiological importance in cattle [10, 11, 12]. Regarding the genotypes P (VP4), bovines reported as frequent worldwide we have the P [1], P [5] and P [11] [10, 11, 12]. In this work, the genotype P [11] was the most frequent, followed by P [1] identified in 2 positive samples and P [7]; in addition, mixed infections corresponding to P [11 + P1] and P [11 + P7] were identified. For example, Da Silva Medeiros et al [10], found P [11] as the most frequent genotype in calf diarrhea in Brazil. Among the findings from different locations in Costa Rica, a variation of P groups (VP4) is reflected; but despite the variety of P groups found in different areas of the country, they are similar to those described in other parts of the world [ 11, 12, 13].

Within the different combinations of genotypes G (VP7) and P (VP4) of Group A Rotavirus in bovines, G6 P [11] was identified as the most frequent detected in 4 of the 16 positive samples, G8 P [7] was the second most frequent combination identified in 3 of the 16 positive samples, followed by G8 P [11] and G10 P [11] found in 2 samples; G8 P [1] and G10 P [1] were identified in one sample. Our findings differ partially from the combinations of G6 P genotypes [5] reported as the most frequent in some investigations in South America [10,11]. Studies carried out in children have

also revealed the presence of the G8, G10 genotype in America and other continents [14,15,16]. Although we also identified mixed infections in this investigation; studies have shown that rotaviruses produce “reassortant” and these genetic changes contribute to the evolution of rotaviruses, resulting in a diversity of genotypes and mixed infections, which is characteristic of rotavirus infections [1, 3, 13, 14]. A review carried out in the countries of the American continent reveals the emergence of new strains of rotavirus [14]. Studies made on the phylogenetics of rotavirus have demonstrated differences in the lineage and sub lineage of genotypes, highlighting the great genetic variability of rotavirus [1]. A small percentage of samples analyzed in this study could not be typified with the use of specific primers, which may be due to the accumulation of point mutations and diversity of reported lineages. We consider that additional experiments and sequencing analysis of these genotypes should be carried out in future investigations, because studies carried out in Nicaragua, Guatemala, Honduras, and the Dominican Republic have reported the presence of rotavirus strains of bovine origin, pigs, and bats in rotavirus samples, excreted from children with diarrhea, showing possible zoonosis [17, 18, 19, 20]. Furthermore, studies carried out in Uruguay have detected human rotavirus genes in calf diarrhea samples [13]. There is a report on the detection analysis of rotavirus strains in bats in Costa Rica. This investigation also included taking samples of bat feces from other countries in the world, revealing strains of rotavirus never reported before, even some of them have similarities with rotavirus strains found in other mammals [21]. Currently, commercial vaccines are available for the prevention of neonatal diarrhea in calves and children that have shown varying degrees of effectiveness. These vaccines are manufactured with the genotypes most frequently found and reported in different parts of the world. In this context, the immunogenicity and efficacy of rotavirus vaccines may be challenged by the evolution of the rotavirus viral genome.

## **CONCLUSION**

Therefore, and adding that research reveals that rotaviruses cause deaths in children, cattle and pigs, we consider urgent the need to maintain an epidemiological surveillance of rotavirus strains in children, domestic animals, wild animals, cattle and pigs in the country, as well as the reinforcement

in the maintenance and consolidation in the unification of detection and characterization protocols of rotavirus in laboratories in Central America.

#### COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

**Table 1. Distribution of G (VP7) P (VP4) of rotavirus strains isolated from bovine in Costa Rica.**

Genotype	P [11]	P[11]+P[7]	P[7]	P[1]	P[11]+P[1]	Total
<b>G8</b>	2(13)	2(13)	3(19)	1 (6)	0	8(51)
<b>G6</b>	4(24)	0	0	0	0	4(24)
<b>G10</b>	2(13)	0	0	1(6)	1(6)	4(25)
<b>Total</b>	8(50)	2(13)	3(19)	2 (12)	1(6)	16 (100)

No. (%) positive (n=16)

**Table 2. Distribution of genotypes G (VP7) and P (VP4) of Group A Bovine Rotavirus in 17 dairy farms sampled in the year 2008.**

Farm	Location	Samples Collected	Positive Samples	G (VP7)	P (VP4)
1	Río Cuarto de Grecia	22	6	G6	P[11]
2	Poas	21	7	G6 , G10	P[11]
3	Pital San Carlos	11	2	G6	P[11]
4	Poas	13	0		
5	Poas	8	1	G8	
6	Poas	7	0		
7	Poas	9	0		
8	Poas	8	0		
9	Poas	7	2	G10	P[11], P[1]
10	Poas	10	1		P[11], P[1]
11	Poas	8	0		
12	Pacayas, Cartago	18	4	G8	P[11], P[7]
13	Cedral, San Carlos	5	1	G8	P[11]
14	Zarcero	10	1	G8	P[1]
15	Sucre	16	1	G10	P[1]

	16	Zarcero	12	1	G10
	17	Sucre	15	1	G10
Total			200	28	

## References:

1. Luchs A, TimenetskyMdo C. Group A rotavirus gastroenteritis: post-vaccine era, genotypes and zoonotic transmission. *Einstein (Sao Paulo)*. 2016; 14(2):278-87.
2. RCWG, Rotavirus Classification Working Group, Newly Assigned Genotypes - Update May 29th 2018, (2018) (Accessed 2 July 2018), <https://rega.kuleuven.be/cev/viralmetagénomics/virus-classification/rcwg>.
3. Martella V, Bányai K, Matthijnsens J, Buonavoglia C, Ciarlet M. Zoonotic aspects of rotaviruses. *VetMicrobiol*. 2010;140(3-4):246-55.
4. Matthijnsens J, Ciarlet M, McDonald SM, Attoui H, Bányai K, Brister JR, et al. Uniformity of rotavirus strain nomenclature proposed by the Rotavirus Classification Working Group (RCWG). *Arch Virol*. 2011; 156(8):1397-413.
5. Teodoroff T, Tsunemitsu H, Okamoto K, Katsuda K, Kohmoto M, Kawashima K, et al. Predominance of porcine rotavirus G9 in japanese piglets with diarrhea: close relationship of their VP7 genes with those of recent human G9 strains. *Journal of Clinical Microbiology*. 2005; 43(3): 1377-1384.
6. De Oliveira LH, Danovaro-Holliday MC, Andrus JK, de Fillipis AM, Gentsch J, Matus CR, Widdowson MA; Rotavirus Surveillance Network. Sentinel hospital surveillance for rotavirus in Latin American and Caribbean countries. *J Infect Dis*. 2009; 200 Suppl 1:S131-9.
7. Bourdett-Stanziola L, Ortega-Barria E, Espinoza F, Bucardo F, Jimenez C, Ferrera A. Rotavirus genotypes in Costa Rica, Nicaragua, Honduras and the Dominican Republic. *Intervirology*. 2011; 54(1):49-52.
8. Gentsch JR, Glass RI, Woods P, Gouvea V, Gorziglia M, Flores J, et al. Identification of group A rotavirus gene 4 types by polymerase chain reaction. *J ClinMicrobiol*. 1992; 30(6):1365-73.
9. Gouvea V, Santos N, TimenetskyMdo C. Identification of bovine and porcine rotavirus G types by PCR. *J ClinMicrobiol*. 1994; 32 (5):1338-40.
10. da Silva Medeiros TN, Lorenzetti E, Alfieri AF, Alfieri AA. G and P genotype profiles of rotavirus A field strains circulating in beef and dairy cattle herds in Brazil, 2006-2015. *Comp ImmunolMicrobiol Infect Dis*. 2019; 64:90-98.

11. Badaracco A, Garaicoechea L, Rodríguez D, Uriarte EL, Odeón A, Bilbao G, et al. Bovine rotavirus strains circulating in beef and dairy herds in Argentina from 2004 to 2010. *Vet Microbiol.* 2012; 158(3-4):394-9. A.
12. Cashman O, Lennon G, Sleator RD, Power E, Fanning S, O'Shea H. Changing profile of the bovine rotavirus G6 population in the south of Ireland from 2002 to 2009. *Vet Microbiol.* 2010; 146(3-4):238-44.
13. Castells M, Caffarena RD, Casaux ML, Schild C, Miño S, Castells F, et al. Phylogenetic Analyses of Rotavirus A from Cattle in Uruguay Reveal the Circulation of Common and Uncommon Genotypes and Suggest Interspecies Transmission. *Pathogens.* 2020; 9(7):570.
14. Bourdett-Stanziola L, Centeno E, Cuevas-Abrego M, Durant-Archibold A. A, Ortega-Barría E, Bucardo F. The Emergence of New Rotavirus Strains in America. Review. *South Asian Journal of Research in Microbiology.* 2021; 11(1), 46-62. doi.org/10.9734/sajrm/2021/v11i130244.
15. Esona MD, Page NA, Akran VA, Armah GE, Steele AD. Characterization of 2 human genotype G10 rotavirus strains, 3008CM and 1784/CI/1999, isolated in Cameroon and Cote d'Ivoire during the 1999-2000 rotavirus season. *J Infect Dis.* 2010; 202 Suppl: S212-9.
16. Zuccotti G, Meneghin F, Dilillo D, Romanò L, Bottone R, Mantegazza C, et al. Epidemiological and clinical features of rotavirus among children younger than 5 years of age hospitalized with acute gastroenteritis in Northern Italy. *BMC Infect Dis.* 2010; 10: 218.
17. Bányai K, Esona MD, Mijatovic S, Kerin TK, Pedreira C, Mercado J, et al. Zoonotic bovine rotavirus strain in a diarrheic child, Nicaragua. *J Clin Virol.* 2009; 46(4):391-3.
18. Quaye O, Roy S, Rungsisuriyachai K, Esona MD, Xu Z, Tam KI, et al. Characterization of a rare, reassortant human G10 P [14] rotavirus strain detected in Honduras. *MemInst Oswaldo Cruz.* 2018; 113(1):9-16.
19. Gautam R, Mijatovic-Rustempasic S, Roy S, Esona MD, Lopez B, Mencos Y, et al. Full genomic characterization and phylogenetic analysis of a zoonotic human G8P [14] rotavirus strain detected in a sample from Guatemala. *Infect Genet Evol.* 2015; 33:206-11.
20. Bourdett-Stanziola L, Centeno E, Nordgren J, Durant-Archibold A, Ortega-Barría E and Bucardo F. Potential Bat-like rotavirus in Hospitalized Children with Diarrhea from the Dominican Republic. *As J Res Infec Dis.* 2021; 8(1):1-7. doi.org/10.9734/ajrid/2021/v8i130225.
21. Simsek C, Corman VM, Everling HU, Lukashev AN, Rasche A, Maganga GD, et al. At Least Seven Distinct Rotavirus Genotype Constellations in Bats with Evidence of Reassortment and Zoonotic Transmissions. *mBio.* 2021;12(1):e02755-20.