

Polio Vaccination and Chronic Fatigue Syndrome

ABSTRACT

Previous research has suggested that enteroviruses may be implicated in the development and persistence of Chronic Fatigue Syndrome (CFS). One method of investigating this topic has been to use a polio vaccination challenge, and a previous study showed that CFS patients had more shedding than healthy controls. There was no effect of the vaccination on the clinical condition or wellbeing of the CFS patients. In the previous study, the control group were more likely to have had a recent booster vaccination. This was controlled in the present study, where 18 CFS patients were randomly assigned to vaccination or placebo conditions. Nine healthy volunteers were also given the polio vaccination. The results confirmed that vaccination had no negative effects on the CFS group. Although there was more virus shedding in the CFS polio group than in the control polio group, this difference was not significant. This study confirms that polio vaccination is not contraindicated in CFS patients but could not confirm that they are more susceptible to enterovirus infection.

Keywords: Chronic Fatigue Syndrome (CFS); Polio vaccination; Enteroviruses.

1. INTRODUCTION

There has been considerable research on enteroviruses and Chronic Fatigue Syndrome (CFS). The early studies led to conflicting results [1-3]. Enteroviruses can be broadly categorised into coxsackie viruses A and B, echoviruses, the polioviruses (types 1, 2, 3), and individual enterovirus serotypes. Early descriptions of CFS suggested that it resembled poliomyelitis [4]. Increased neutralising antibodies and IgM antibodies specific to coxsackie B viruses were found in CFS patients compared to healthy controls [5]. Enteroviruses were also more frequent in the faecal samples of those with CFS than controls [3]. This led to the suggestion that CFS may be caused by persistent enteroviral infection [6]. Early results provided mixed support for this view. In one study, CFS patients were found to show more enteroviral persistence in their muscles than healthy controls [7]. However, a later, larger study found no differences between the groups [8].

More recent studies have also led to conflicting results (see [9, 10] for reviews). Several studies have obtained results that argue against the persistence of enteroviruses in CFS, and this persistence being the cause of the pathogenesis of the syndrome [11-16]. Other research supported chronic enterovirus infection in CFS patients [17-23].

An alternative approach to CFS and enteroviruses has used the polio vaccine paradigm [24]. This approach measured the virological, immunological, symptomatic and behavioural effects of vaccination of CFS patients with an oral poliomyelitis vaccine (Sabin). This paradigm allows one to examine enterovirus infection after administration of a live booster dose. All of the CFS group had been given a primary vaccination with no adverse effects. It has previously been suggested that there is an association between vaccination and CFS [25]. The research examined the possible effects of live poliomyelitis re-vaccination on the symptoms and behaviour of CFS patients and healthy controls. In addition, differences in the two groups immune responses to the vaccine challenge were also investigated. Vaccination did not change the clinical condition of the CFS patients, and their T- and B-cell polio-specific responses were no different from the healthy volunteers. However, the CFS patients did shed higher poliovirus levels, as identified by direct isolation, compared to the healthy controls. This study showed that live poliovirus vaccination in CFS patients is not contraindicated, which argues against the view that CFS is exacerbated by vaccinations. In addition, it is unlikely that a specific immune defect in response to enteroviruses can account for the pathogenesis of CFS. However, the increased poliovirus shedding in the CFS patients requires further study, and the underlying mechanisms still need to be identified.

Many of the findings linking enteroviruses to CFS have been criticised in terms of poor methodology. In the original polio vaccine study, the control group had been given “booster” vaccinations more recently than the patients, which could plausibly account for the increased virus shedding. The present study addressed this by using a healthy control group that was matched with the CFS group in terms of the timing of the primary polio vaccination. The aims of the study were identical to the original one, namely, to determine whether polio vaccination of CFS patients led to any adverse effects and to examine virus shedding after vaccination.

2. METHODS

The study was approved by the local, regional ethical committee and carried out with the informed consent of the participants.

2.1 Participants

Eighteen patients with CFS, diagnosed according to the Oxford criteria, were recruited randomly from a panel of those who had in the past or were currently attending the University of Wales College of Medicine (UWCM) CFS outpatient clinic. The demographic characteristics and illness history of these patients can be shown in Tables 1 and 2. In general, the profile of this sample was consistent with those of the typical patients attending the clinic from which they were recruited and of other CFS populations reported in the literature. A further nine individuals without CFS were recruited from the partners of the CFS patients taking part.

Table 1. Demographic characteristics of CFS patients and controls

	CFS patients (vaccine)	CFS patients (placebo)	Controls (vaccine)
Sex (%)	57 Female 43 Male	75 Female 25 Male	25 Female 75 Male
Age Range	38 – 65 years	46 – 54 years	39 – 65 years
Marital status (%)			
Single	14	-	25
Married	86	100	75
Education level (%)			
Primary education only	7	-	-
Left school before 16	36	25	38
Completed ‘O’ levels	29	25	12
Completed ‘A’ levels	-	25	12
At least one year at the University	-	25	12
BSc or BA	21	-	12
PhD, MD or other	7	-	12

Table 2. Clinical profile of the CFS patients and controls

	CFS patients (vaccine)	CFS patients (placebo)	Controls (vaccine)
Mean duration of illness (months)	126.3	107.5	N/A
Mean time since diagnosis (months)	73.8	52.0	N/A
Current severity (%)			
Worse than at any stage	14	0	N/A
Bad	14	50	
Bad with some recovery	43	50	
Recovering with relapses	29	0	
Completely recovered	0	0	
Symptoms (%) – at time of study			
Physical weakness (Yes)	79	100	37
Excessive fatigue	93	100	25
Legs feeling heavy	79	75	25
Muscle pain	71	100	-
Pain in chest	43	25	-
Painful joints	79	100	12
Nausea	29	25	-
Indigestion	43	50	12
Bloated stomach	50	75	12
Wind	57	50	25
Sore throat	57	50	-
Headache	57	50	12
Earache	36	50	-
Sore eyes	50	50	12
Sensitive to noise	71	75	-
Sensitive to light	64	100	12
Feeling hot/cold	86	75	12
Sweating	71	75	-
Shivering	43	50	-
Swollen glands	79	50	-
Racing heart	43	75	-
Insomnia	36	100	12
Depression	50	75	25
Anxiety	57	50	-
Loss of concentration	79	100	-
Loss of memory	79	75	25
Allergies	29	0	-

2.2 Study design

Eighteen patients with CFS and nine individuals without CFS were recruited into a 28-day double-blind study to determine any effects of poliovirus vaccination in individuals with CFS. On day 0, the healthy

controls received the poliomyelitis vaccine (SmithKline Beecham), a live (Sabin) polio vaccine containing a mix of attenuated poliovirus types 1,2 and 3 (6 log₁₀ poliovirus 1, 5 log₁₀ poliovirus 2 and 5.5 log₁₀ poliovirus 3) while the patients with CFS were randomly allocated into either the placebo (sterile saline) or vaccine treatment groups and both the patients and experimenters who conducted the investigation were blind to which individuals received the vaccine. All individuals were followed up on four occasions thereafter (days 2, 7, 14, 28 post-vaccination).

2.3 Virology

Pan-enterovirus and poliovirus-specific assays were developed and utilised for this part of the study. A method based on nucleic acid sequence-based amplification (NASBA) was found to be suitable for the analysis of poliovirus shedding [26](Fox et al., 2002). NASBA has been found to be a suitable alternative to RT-PCR for the detection of enterovirus sequences. These kit-based reagents have enabled a wide range of laboratories to use molecular-based diagnostic procedures to identify RNA viruses.

2.4 Clinical assessment

Subjective and objective assessments of the physical wellbeing of the sample were undertaken throughout the investigation by:

- completion of a physical symptoms index at the start and a questionnaire completed at week 0 and week 4 of the study, which examined fatigue-related and somatic symptoms.
- The objective assessments were undertaken by the clinician responsible for the CFS patient group, who arranged a consultation session for each patient on days 0 and 28. The clinician was blind to the treatment group into which each patient had been allocated.

2.5 Psychosocial assessment

A series of questionnaires were administered to examine demographic and illness characteristics, current symptoms, and psychosocial measures. The second series of questionnaires were administered during the study: these examined: symptoms [24], mood [27], depression [28], state anxiety [29], emotional difficulties, cognitive difficulties, somatic symptoms, and fatigue [30].

2.6 Cognitive assessment

Subjective ratings of cognitive function [31] and objective assessments of cognitive functioning (simple reaction time and cognitive vigilance [32]) were undertaken during the study.

3. RESULTS

3.1 Virus shedding results

More patients than controls shed virus following vaccine challenge, but this effect was not significant (see Table 3).

Table 3. Poliovirus shedding in patients and controls

	Controls	CFS Vaccine	CFS Placebo
Day 2 (%)	12	21	0
Day 7	37	36	0
Day 14	12	21	0
Day 28	12	14	0

3.2 Effects of vaccination on mood, symptoms, and performance

There were no significant effects of vaccine challenge on subjective reports of health and wellbeing or objective measure (see Tables 4 and 5).

Table 4. Baseline and post-vaccination questionnaire scores (means, s.d.s in parentheses)

	Controls		CFS Vaccine		CFS placebo	
	Day 0	Day 28	Day 0	Day 28	Day 0	Day 28
Positive Mood (2)	38.62 (7.07)	38.87 (7.36)	31.07 (9.18)	28.46 (7.79)	31.75 (11.73)	33.33 (2.31)
Negative Mood (1)	8.50 (5.48)	6.62 (4.21)	20.71 (9.53)	21.16 (6.59)	18.50 (8.70)	17.67 (11.93)
Centre for Epidemiologic Studies Depression Scale (1)	27.12 (8.04)	28.12 (8.13)	37.50 (9.29)	38.69 (9.02)	35.00 (9.56)	35.00 (9.54)
State Anxiety (1)	29.50 (6.19)	32.75 (6.63)	39.93 (9.92)	42.31 (12.82)	34.75 (7.89)	37.00 (6.56)
Emotional Distress (1)	20.50 (5.01)	22.00 (7.13)	45.28 (22.88)	42.08 (15.06)	35.00 (19.34)	31.00 (19.47)
Fatigue (1)	25.12 (17.87)	21.62 (12.18)	60.28 (15.39)	62.38 (12.93)	61.75 (11.95)	48.67 (4.16)
Cognitive Difficulty (1)	17.12 (11.44)	15.50 (4.69)	48.43 (16.46)	48.31 (14.53)	40.25 (17.82)	38.00 (17.00)

Somatic Stress (1)	19.87 (3.27)	20.12 (4.26)	55.93 (21.93)	57.23 (19.53)	53.25 (30.43)	53.00 (19.31)
Beck Depression Inventory (1)	5.37 (4.66)	5.25 (4.65)	12.86 (6.61)	13.08 (5.56)	11.25 (9.21)	8.33 (3.78)

(1) High scores = greater impairment ; (2) Low scores = greater impairment

Table 5. Baseline and post-vaccine scores for the performance tests (means and s.d.s)

	Controls		CFS Vaccine		CFS Placebo	
	Day 0	Day 28	Day 0	Day 28	Day 0	Day 28
SRT – mean RT msec	359.62	321.12	383.57	411.69	480.50	488.00
Repeated digits vigilance task – Mean Hits	14.13	16.50	12.57	11.31	12.50	13.67

4. DISCUSSION

The present study was part of a larger programme on viral infections and CFS. In the first part of the research, prospective studies of upper respiratory tract illnesses (URTIs) were conducted [33, 34]. The

that this was not due to a reporting bias. Indeed, similar virus identification rates were obtained in patient and control groups with clinical illnesses suggesting that identical mechanisms were in operation but that the CFS patients had more infections and illnesses. The results also showed that the CFS patients had more sub-clinical infections, which again supports the view that this group are particularly susceptible to acute infections.

The present study was unable to replicate our previously reported finding that CFS patients are more likely to shed virus following polio vaccine challenge [24]. In the original study, the control group had been given “booster” vaccinations more recently than the patients, which plausibly accounts for our earlier results. The study did confirm that vaccination had no detrimental effect on the patients, which supports our earlier view that it is unlikely to cause the type of problem suggested by some sources.

CONSENT

The research was carried out with the informed consent of the participants.

ETHICAL APPROVAL

The research was carried out with the approval of the local regional ethics committee.

REFERENCES

1. Jones JF, Ray CG, Minnich LL, Hicks MJ, Kibler R, Lucas, DO. Evidence for active Epstein-Barr virus infection in patients with persistent, unexplained illnesses: Elevated anti early antigen antibodies. *Ann. Intern. Med.* 1985; 102: 1-7.
2. Wakefield D, Lloyd A, Dwyer J, Salahuddin SZ, Ablashi DV. Human herpesvirus 6 and myalgic encephalomyelitis. *Lancet.* 1988; 334: 1059.
3. Yousef GE, Bell EJ, Mann GJ, Bell EJ, Murugesan V, McCartney RA, Mowbray, JF. Chronic enterovirus infection in patients with postviral fatigue syndrome. *Lancet.* 1988; 334:146-150.
4. Behan PO, Bakheit AMO. Clinical spectrum of post-viral fatigue syndrome. *Br. Med. Bull.* 1991; 47: 793-808.
5. Behan PO, Behan WMH, Bell, EJ. The post-viral fatigue syndrome: An analysis of the findings in 50 cases. *J. Infect.* 1985; 10: 211-222.
6. Cunningham L, Bowles NE, Archard, LC. Persistent virus infection of muscle in postviral fatigue syndrome. *Br. Med. Bull.* 1991; 47: 852-871.
7. Gow JW, Behan WMH, Clements GB, Woodall C, Riding M, Behan, PO. Enteroviral sequences detected by polymerase chain reaction in muscle biopsies of patients with the postviral fatigue syndrome. *BMJ.* 1991; 302: 692-696.
8. Gow JW, Behan WMH, Simpson K, McGarry F, Keir S, Behan, PO. Studies on enterovirus in patients with chronic fatigue syndrome. *Clin. Infect. Dis.* 1994; 18 (Suppl. 1): s126-s129.

9. Chia JK. The role of enterovirus in chronic fatigue syndrome. *J Clin Pathol*. 2005 Nov;58(11):1126-32. doi: 10.1136/jcp.2004.020255.
10. O'Neal AJ, Hanson MR. The Enterovirus Theory of Disease Etiology in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Critical Review. *Front Med (Lausanne)*. 2021 Jun 18;8:688486. doi: 10.3389/fmed.2021.688486.
11. Swanink CM, Melchers WJ, van der Meer JW, Vercoulen JH, Bleijenberg G, Fennis JF, Galama JM. Enteroviruses and the chronic fatigue syndrome. *Clin Infect Dis*. 1994 Nov;19(5):860-4. doi: 10.1093/clinids/19.5.860.
12. Buchwald D, Ashley RL, Pearlman T, Kith P, Komaroff AL. Viral serologies in patients with chronic fatigue and chronic fatigue syndrome. *J Med Virol*. 1996 Sep;50(1):25-30.
13. Manian FA. Simultaneous measurement of antibodies to Epstein-Barr virus, human herpesvirus 6, herpes simplex virus types 1 and 2, and 14 enteroviruses in chronic fatigue syndrome: is there evidence of activation of a nonspecific polyclonal immune response? *Clin Infect Dis*. 1994 Sep;19(3):448-53. doi: 10.1093/clinids/19.3.448.
14. Lindh G, Samuelson A, Hedlund KO, Evengård B, Lindquist L, Ehrnst A. No findings of enteroviruses in Swedish patients with chronic fatigue syndrome. *Scand J Infect Dis*. 1996;28(3):305-7. doi: 10.3109/00365549609027178.
15. Milton JD, Clements GB, Edwards RH. Immune responsiveness in chronic fatigue syndrome. *Postgrad Med J*. 1991 Jun;67(788):532-7. doi: 10.1136/pgmj.67.788.532.
16. McArdle A, McArdle F, Jackson MJ, Page SF, Fahal I, Edwards RH. Investigation by polymerase chain reaction of enteroviral infection in patients with chronic fatigue syndrome. *Clin Sci (Lond)*. 1996 Apr;90(4):295-300. doi: 10.1042/cs0900295.
17. Chia JK, Chia AY. Chronic fatigue syndrome is associated with chronic enterovirus infection of the stomach. *J Clin Pathol*. 2008 Jan;61(1):43-8. doi: 10.1136/jcp.2007.050054.
18. Chia J, Chia A, Voeller M, Lee T, Chang R. Acute enterovirus infection followed by myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) and viral persistence. *J Clin Pathol*. 2010 Feb;63(2):165-8. doi: 10.1136/jcp.2009.070466.
19. Galbraith DN, Nairn C, Clements GB. Phylogenetic analysis of short enteroviral sequences from patients with chronic fatigue syndrome. *J. Gen. Virol*. 1995; 76: 1701-1707.
20. Nairn C, Galbraith DN, Clements GB. Comparison of coxsackie B neutralisation and enteroviral PCR in chronic fatigue patients. *J Med Virol*. 1995 Aug;46(4):310-3. doi: 10.1002/jmv.1890460404.
21. Lane RJ, Soteriou BA, Zhang H, Archard LC. Enterovirus related metabolic myopathy: a postviral fatigue syndrome. *J Neurol Neurosurg Psychiatry*. 2003 Oct;74(10):1382-6. doi: 10.1136/jnnp.74.10.1382.
22. Clements GB, McGarry F, Nairn C, Galbraith DN. Detection of enterovirus-specific RNA in serum: The relationship to chronic fatigue. *J. Med. Virol*. 1995; 45: 156-161.
23. Douche-Aourik F, Berlier W, Féasson L, Bourlet T, Harrath R, Omar S, Grattard F, Denis C, Pozzetto B. Detection of enterovirus in human skeletal muscle from patients with chronic

- inflammatory muscle disease or fibromyalgia and healthy subjects. *J Med Virol*. 2003 Dec;71(4):540-7. doi: 10.1002/jmv.10531.
24. Vedhara K, Llewelyn M, Fox JD, Jones R, Clements GB, Wang ECY, Smith AP, Borysiewicz LK. Consequences of live poliovirus vaccine administration in Chronic Fatigue Syndrome. *Journal of Neuroimmunology*. 1997; 75: 183-195.
25. Canada Communicable Disease Report. Report of the Working Group on the Possible Relationship between Hepatitis B Vaccination and the Chronic Fatigue Syndrome. 1993; Vol. 19: pp. 25-28.
26. Fox JD, Han S, Samuelson A, Zhang Y, Neale ML, Westmoreland D. Development and evaluation of nucleic acid sequence based amplification (NASBA) for diagnosis of enterovirus infections using the NucliSens Basic Kit. *J Clin Virol*. 2002 Feb;24(1-2):117-30. doi: 10.1016/s1386-6532(01)00241-4.
27. Zevon MA, Tellegen A. The structure of mood change: An idographic nomothetic analysis *J Pers. Soc. Psychol*. 1982; 43: 111-122.
28. Radloff L. The CES-D scale: A self-report depression scale for use in the general population. *Appl. Psychosoc. Measures* 1977; 1: 385-401.
29. Spielberger C, Gorsuch R, Lushene RE. *STAI Manual for the State-Trait Anxiety Inventory*. Consulting Psychologists Press, Palo Alto. 1977.
30. Ray C, Weir WRC, Phillips S, Cullen S. (1992) Development of a measure of symptoms in chronic fatigue syndrome: The profile of fatigue-related symptoms (PFRS). *Psychology and Health*. 1992;7: 27-43.
31. Broadbent DE, Cooper PJ, Fitzgerald PF, Parkes, KR. The cognitive failures questionnaire (CFQ) and its correlates. *Br. J. Clin. Psychol*. 1982; 21: 1-6.
32. Smith A, Whitney H, Thomas M, Perry K, Brockman P. (1995) Effects of regular alcohol intake and stress on mental performance, mood and cardiovascular function. *Hum. Psychopharmacol*. 1995; 10: 423-431.
33. Smith AP, Thomas M, Borysiewicz L, Llewelyn M. Chronic fatigue syndrome and susceptibility to upper respiratory tract illnesses. *British Journal of Health Psychology* 1999, 4, 327-335.
34. Smith AP, Thomas MA. Chronic fatigue syndrome and increased susceptibility to upper respiratory tract infections and illnesses. *Fatigue: Biomedicine, Health & Behavior* 2015, 3(3), 156-163. <http://dx.doi.org/10.1080/21641846.2015.1033271>