

Skin Irritation Associated with Hand Hygiene Practice and Use of Face Mask during COVID-19 pandemic among the people in Gazipur, Bangladesh

Abstract:

Background: In COVID-19 pandemic increasing the frequency of hand washing and disinfection is the most important preventive measure of this disease which is causing in deleterious impact on skin among general population. So the aim of the study was to evaluate whether these measures correlate with development of skin damage. *Methods:* This was a descriptive type of cross-sectional, observational study conducted over a period of six months from April 2021 to September 2021 including the 1166 skin patients coming to some private clinics of Gazipur. Skin diseases were diagnosed by dermatologist on basis of clinical morphology with appropriate investigations. Data were collected by structured questionnaire and analyzed using SPSS. *Results:* Out of 1166 patients, 35.68% reported skin changes or symptoms over hands (ScH), and 13.98% reported skin changes on their face (ScF). Around 87.39% and 86.53% of the participants reported a change in hand washing habits and sanitizer using habits during the COVID-19 pandemic. There were significant higher percentage (57.21%) of skin conditions in females (ScH: 45.28% vs ScF: 17.30%), 43.82% individuals were working in environments requiring frequent hand washing (ScH: 41.88% vs. ScF: 18.59%), 39.88% working in facilities where they have to interact with people during the pandemic (ScH: 42.58% vs. ScF: 19.57%), 5.75% those encountering COVID-19 patients (ScH: 49.25% vs. ScF: 25.64%), 15.78% those exposed to chemicals (ScH: 52.72% vs ScF: 25.82%), and 9.17% healthcare workers (ScH: 57.00% vs ScF: 31.77%). Almost 35.22% reported skin dryness, 9.44% reported changes in the texture, 13.42% reported scaling, 16.52% reported itchiness, 4.46% reported changes in skin color, 14.52% reported redness, and 4.62% reported pain/ burning, while 1.80% reported skin ulcers. *Conclusion:* Our study suggests that the general population's skin was negatively affected by COVID-19 precautions where increased frequency of hand washing and the use of alcohol-based sanitizers, overuse of soaps were contributing factors for skin disorders.

Keywords: COVID-19 pandemic, precautions, questionnaire, general population, skin, hand hygiene

Introduction:

The health of general population and health workers is currently threatened by spread of the novel corona virus disease (COVID-19) pandemic with millions of cases and deaths¹. This infection can be spread from person to person through the inhalation of small droplets produced as a result of sneezing, coughing, or talking by infected persons or indirectly through contact with contaminated surfaces with hands². Using surface disinfectants, regularly washing hands, using soaps, hand sanitizers, covering the mouth and face with face masks, and using personal protective equipment (PPE), social distancing, avoiding touching the face wearing disposable gloves can prevent and reduce spread of this infection is³.

Use of soaps and water or an alcohol-based sanitizer is the effective, simple, available method for maintaining hand hygiene against COVID-19 infection^{4,5}. To prevent the spread by half, WHO recommends hand hygiene at the right times in a proper way.^{3,6}

Higher concentrations of alcohol mixed with other organic/inorganic acids increases antiviral and antibacterial activity of sanitizers^{7,8}. Sanitizers can prevent the spread of many pathogens⁹. The sebaceous glands of the skin secrete oils have antiviral properties¹⁰, skin is dehydrated by the continual use of alcohol-based hand sanitizers which washes away these oils, resulting in fissures and erosion, making easy access to microbial infection³. Skin diseases, most commonly irritant and allergic contact dermatitis can be caused by excess use of soap which might disturb the normal skin flora and natural protective skin barrier.¹¹,

Increased skin sensitivity to any physical and chemical agents^[12], also contact cutaneous xerosis, flaking and eczema.^[13,14] Some countries reported an increased incidence of skin problems during this pandemic.^[15-18]

To our knowledge, no studies have been locally conducted regarding the prevalence of hand skin diseases among the general population during COVID-19 in Gazipur. In this study, skin damage among the general population might be observed with increased frequency due to increased hand hygiene measures. Thus, we aimed to evaluate prevalence and associated factors of skin disorders during the COVID-19 pandemic in Gazipur.

Methodology:

Study Design and study population: This was a descriptive, cross-sectional, observational study, conducted over a period of six months from April 2021 to September 2021 including the 1166 skin patients coming to some private clinics of Gazipur, Bangladesh. Skin diseases were diagnosed by dermatologist on basis of clinical morphology with appropriate investigations. The consent was taken from patients and approval from proper authority.

Data Collection: Data were collected in printed proforma. The questionnaire was developed after an in-depth literature review and was reviewed by dermatologists. Eight questions for socio-demographic data, seven for hand washing and eight for sanitizing habits, two for gloves, three for mask usage were included in questionnaire.

Statistical Analysis. A descriptive analysis was conducted, A chi-squared test was conducted. A p value of 0.05 or less was considered significant. Data are reported in tabular and graphical form. SPSS version 25 was used to conduct the analysis.

Results:

Data from 1166 individuals were collected. Of the total number of participants, 57.21% were females, 76.59% were single, 26.41% were employed or self-employed, and 73.58% were students. (Table 1).

When asked about their hand washing practices, 43.82% of the employees reported that they work in settings that require frequent hand washing, and 39.88% reported that they work in settings that necessitate encounters with customers. Additionally, 15.78% reported working in settings that require the use of irritating materials like chemicals, soaps, or other detergents. Only 5.75% of participants were working in settings in which there was direct contact with COVID-19 patients, and 9.17% were healthcare workers (Table 1).

There is a significantly higher prevalence of skin conditions on patients' hands and face in the following demographics: females (skin condition on hands (ScH): 45.28% and skin condition on face (ScF): 17.30%), married individuals (ScH: 49.82% and ScF: 17.95%), those who were employed (ScH: 40.58% and ScF: 19.48%), individuals working in environments that require hand washing several times (ScH: 41.88% vs. ScF: 18.59%), those who were working in facilities where they have to interact with people during the pandemic (ScH: 42.58% vs. ScF: 19.57%), those who were encountering COVID-19 patients (ScH: 49.25% vs. ScF: 25.64%), those working in facilities where they have to work with chemicals (ScH: 52.72% vs. ScF: 25.82%), and healthcare workers (ScH: 57.00% vs. ScF: 31.77%) [Table 1].

Table 1: Socio-demographics and hand hygiene practices during the COVID-19 pandemic, and the prevalence of skin conditions

Skin conditions	Total = 1166 (%)	Skin conditions on hand (n=416)	P value	Skin conditions on face (n=163)	P value
Gender					
Male	499 (42.79)	114 (22.84)	<0.001	47 (9.42)	<0.001
Female	667 (57.21)	302 (45.28)		116 (17.30)	

Marital Status					
Single	893 (76.59)	280 (31.35)	<0.001	114 (12.76)	0.19
Married	273 (23.41)	136(49.82)		49(17.95)	
Employment					
Employed	308 (26.41)	125 (40.58)	<0.001	60 (19.48)	0.004
Housewife/Student	858 (73.58)	291 (33.92)		103 (12.00)	
Healthcare worker					
Yes	107(9.17)	61 (57.00)	<0.001	34 (31.77)	<0.001
No	1059 (90.82)	355 (33.52)		129 (12.18)	
Hand washing required					
Yes	511 (43.82)	214 (41.88)	<0.001	95 (18.59)	0.022
No	655 (56.17)	202 (30.84)		68 (10.38)	
Encounter with people					
Yes	465 (39.88)	198 (42.58)	<0.001	91 (19.57)	<0.001
No	701 (60.12)	218(31.09)		72 (10.27)	
Encounter with COVID-19 patients					
Yes	67 (5.75)	33 (49.25)	0.003	17 (25.64)	0.006
No	1099 (94.25)	583 (34.85)		146 (13.28)	
Use of chemicals					
Yes	184 (15.78)	97 (52.72)	<0.001	47 (25.82)	<0.001
No	982 (84.22)	319 (32.48)		116 (11.81)	

More than two-thirds (87.39%) of the participants re-ported that their hand washing habits changed during the COVID-19 pandemic. During the pandemic, the frequency of hand washing increased drastically more than 10 times for 55.57% of individuals. Nearly two-thirds (69.04%) reported washing their hands for less than 1 minute everytime, while 26.41% reported washing it for 1–2 minutes. 81.47% reported using warm water when washing their hands. Nearly half reported using antiseptic soaps for washing their hands, 53.60% reported using regular soaps, 51.20% reported using perfumed soaps, and 85.51% reported using liquid soaps (Table2).

Regarding the usage of hand sanitizers, 86.53% reported that their habit of using hand sanitizers has changed during the pandemic, 80.19% were using them more than 10timesperday, and 75.38% werenot using perfumed sanitizers. 68.78% reported using hand sanitizers after coming in to contact with any person or surface, while 22.29% were using it every one or two hours. In addition, 63.55% used it every time they entered their homes after being outside, and 49.48% reported using it before and after eating (Table2). In regard to glove usage, 49.48% were not using any gloves, with 9.78% of participants being reported not using gloves because of skin problems. Out of those who did wear gloves, 53.83% used 1–2 pairs per day (Table2).

Table 2: Determinants of skin diseases on hands in period of COVID-19 infection

Variables	Totaln:1166(%)	<u>Yes(%)</u>	<u>pvalue</u>
Altered Hand cleaning practice			
Yes	1019 (87.39)	403(39.55)	<0.001
No	147(12.61)	13(8.84)	
Hand cleaning frequency			
≤10times	648 (55.57)	289(44.60)	<0.001
>10times	518(44.42)	127(24.52)	
Washing duration			
>1min	805(69.04)	277(34.41)	
1–2min	308(26.41)	118 (38.31)	0.040
<2min	53(4.55)	21(39.62)	

<i>Temperature of water</i>			
Tepid/warmish	950 (81.47)	328(34.55)	
Cold	121(10.38)	29(23.97)	0.004
Hot	95(8.15)	59(62.10)	
<i>Variety of soap</i>			
Regular	625 (53.60)	220(35.20)	0.375
Antiseptic	541 (46.34)	196(36.23)	
<i>Fragranced soap</i>			
Yes	597 (51.20)	206(34.51)	0.644
No	569 (48.79)	210(36.91)	
<i>Consistency of soap</i>			
Solid	169(14.41)	56(33.14)	0.395
Liquid	997 (85.51)	360(36.12)	
<i>Altered hand sanitization practice</i>			
Yes	1009 (86.53)	382(37.86)	<0.001
No	157(13.46)	34(21.66)	
<i>Frequency of hand sanitization</i>			
≤10times	935(80.19)	376(40.21)	0.023
>10times	231(19.81)	40(17.32)	
<i>Alcohol concentration</i>			
>60%	108(9.26)	44(40.74)	
<60%	261(22.38)	124(47.51)	<0.001
Do not know	797 (68.35)	248(30.74)	
<i>Fragranced sanitizer</i>			
Yes	287(24.61)	94(32.75)	0.538
No	879 (75.38)	322(36.63)	
<i>Sanitizing when touching person/surface</i>			
Yes	802 (68.78)	306(38.15)	<0.001
No	364 (31.22)	110 (30.22)	
<i>Each 1 or 2 hours</i>			
Yes	260(22.29)	108(41.54)	0.003
No	906(77.70)	308(33.99)	
<i>After entering home from exterior</i>			
Yes	741 (63.55)	278(37.52)	0.044
No	425(36.45)	138(32.47)	
<i>Before and after eating</i>			
Yes	577 (49.48)	229(39.69)	<0.001
No	589(50.51)	187(31.75)	
<i>Use of hand gloves daily</i>			
Do not wear	550 (47.17)	173(31.45)	0.010
<1pair	616 (52.83)	243(39.45)	
<i>Quit wearing gloves</i>			
Yes	114(9.78)	65(57.02)	<0.001
No/donotknow	1052(90.22)	351(33.36)	
<i>Had hand dermatitis</i>			
Yes	192(16.47)	104(54.17)	<0.001
No	974(83.53)	312(32.03)	
<i>Household work</i>			
≤2hours/week	729(62.52)	237(32.51)	0.027
>2hours	437(37.48)	179(40.96)	

Skin conditions on hands were significantly more common among specific groups, including individuals who did not change their hand washing habits during the pandemic (37.1%), those who washed their hands several times per day (43.1%), those who washed more than 2 minutes (38.7%), or those who

washed hands with cold water (38.9%). Skin conditions were also reported to be significantly more common among individuals who altered their usage of hand sanitizers during the pandemic (36.3%), those who used sanitizers several times per day (39.2%), those who used sanitizers with a greater alcohol concentration (46.1%), those who sanitized every one to two hours (40.8%), those who sanitized after coming from outside (36.4%), and those who sanitized before and after eating (38.7%) (Table2).

Table 3. Determinants of skin diseases on face in period of COVID-19 infection

Variable	Total n : 1166	Yes (%)	p value
<i>Use of face mask</i>			
Surgical	865 (74.18)	133 (15.37)	0.018
Others	225 (19.30)	19 (8.44)	
Do not use	76 (6.52)	11 (14.47)	
<i>Duration of face mask usage</i>			
<2 hours	620 (53.17)	82 (13.22)	0.635
>2 hours	341 (29.25)	52 (15.25)	
Do not use	205 (17.8)	29 (14.15)	
<i>Frequency of mask use</i>			
1-2 times	760 (65.18)	110 (14.47)	0.830
> 2 times	167(14.32)	26(15.57)	
Do not use	239(20.50)	27(11.30)	

With respect to face masks, 74.18% reported using surgical/medical face masks; 53.17% of the participants used the masks for half an hour to 2 hours; and 65.18% reported changing it 1–2 times per day. However, while skin conditions on the face were related to wearing face masks, the duration of wearing face masks and the number of face masks changed per day did not affect skin conditions (Table3).




Figure 1: Skin Conditions on hand during the pandemic

Of the total number of respondents, 35.68% (416 individuals) reported skin changes or symptoms during COVID-19, of whom 35.22% reported skin dryness, 9.44% reported changes in the texture, 13.42%

reported scaling, 16.52% reported itchiness, 4.46% reported changes in skin color, 14.52% reported redness, and 4.62% reported pain/burning, while 1.80% reported skin ulcers (Figure 1).

Discussion:

This population-based study was conducted to determine the prevalence and associated factors of hand skin disorders in Gazipur during the ongoing COVID-19 pandemic. In the study, most participants were females, single and unemployed, while 9.17% were healthcare workers. 55.57% practiced frequent hand washing. 87.39% and 86.53% of the participants reported a change in hand washing habits and sanitizer using habits respectively, with their frequency increasing to more than 10 times a day. As a result, 35.68% reported skin changes over hands and 13.98% over the face.

Dermatitis is an inflammatory response of the skin caused by allergens, irritant substances, or both.^[19] As seen in this study, frequent hand washing caused skin changes, mostly skin dryness 35.22%, with some cases of redness, scaling, pain, itching, or even ulcers. A review by Cristina Beiu highlighted frequent hand washing for COVID-19 prevention can cause hand dermatitis^[13]. A study in India found overzealous hand hygiene during the COVID 19 pandemic causing an increased incidence of hand eczema among general population^[15]. Skin irritation is significantly more among the healthcare workers (57.00%) and individuals who have contact with COVID-19 patients (49.25%) in this study, as they follow strict hygiene practices and wear PPE for a longer period of time. Similarly, a Chinese study reported a 74.5% prevalence^[16], while another German study reported a 90.2% incidence and of eczema among health care workers^[17]. Moreover, a study conducted in Milan, Italy, reported an increased frequency of hand eczema^[18].

The skin's permeability to various agents can be increased by disruption of the skin's outer layer by constant and prolonged use of soap water in humid environments^[20]. Many diseases including atopic dermatitis can be increased by stress, atmospheric aspects, quarantine, and lockdowns.^[21] Irritant contact dermatitis can be triggered by wet work and gloves occlusion causes skin barrier impairment, when combined with exposure to soaps or sanitizers.^[22] Healthcare workers are prone to adverse skin reactions working in COVID-19 wards wearing PPE for several hours.^[23] Other studies have shown that development of contact dermatitis due to use of gloves.^[5, 24], another study showed that dermatitis increased three times with the use of >5 pairs of gloves.^[25] It was also found in our study that dermatitis was significantly prevalent in 39.45% of individuals who used gloves (>1 pair per day).

In our study the skin changes (47.51%) were significantly present in participants who used sanitizers with an alcohol concentration of more than 60%. Alcohol-based hand rubs with alcohol concentration of >60% are the most appropriate alternative, when one's hands are not visibly dirty or when soaps and water are not available according to the WHO's recommendation.^[6] Similarly, in another study, washing hands >10 times a day and using alcohol-based gel caused skin damage.^[18] Since coronavirus is enveloped in a lipid bi-layer, an alcohol-based sanitizer is undoubtedly effective. The major problems are the existence of substandard products in the markets as well as the emergence of alcohol tolerance, antimicrobial resistance (AMR), opportunistic infection, and product toxicity.^[26-28] Recent published studies also reported high prevalence of eczema in healthcare workers.^[5, 29]

In this study, some of the participants also faced skin changes on their faces due to the excessive use of face masks and PPE. A recent study reported that 97% of the skin damage was due to enhanced protective measures, and these include 83% of the nasal bridge lesion^[30]. Other studies reported similar findings in addition to reporting other dermatologic side effects like pressure injury, urticaria, dryness of the skin, allergic contact dermatitis, and aggravation of underlying dermatosis. In all of these conditions, occlusion and friction were the main contributing factors^[31-34]. These results were more commonly found among healthcare workers, who wear protective gear for prolonged times.^[16]

However, the major strength of the study is that it covers both the general population and healthcare workers, in addition to using a relatively large sample size.

Conclusion:

In conclusion, our study suggests that COVID-19 infection prevention measures creating deleterious impact on the general population's and health workers's skin. A significant association between more hand washing frequency (>10 times) and duration (>2 min), sanitization with alcohol concentration (>60%) and development of skin damage during the pandemic was detected. Concurrent use of balanced preventive measure with maintaining skin integrity is of utmost importance. Therefore, we recommend spreading awareness of skin protective modalities and the use of regular skin moisturization for hand protection. Proper awareness, appropriate knowledge and good practice can prevent skin changes during this pandemic, with regular hydration of the skin.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

References

1. "Coronavirus disease (COVID-19) situation report-206," 2020, https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200813-covid-19-sitrep-206.pdf?sfvrsn=bf38f66b_6
2. Y. Bai, L. Yao, T. Wei, F. Tian, D.-Y. Jin, and L. Chen, "Presumed asymptomatic carrier transmission of COVID-19," *JAMA*, vol. 323, no. 14, pp. 1406-1407, 2020.
3. D. Pradhan, P. Biswasroy, P. Kumar Naik, G. Ghosh, and G. Rath, "A review of current interventions for COVID-19 prevention," *Archives of Medical Research*, vol. 51, no. 5, pp. 363–374, 2020.
4. E. E. Sickbert-Bennett, L. M. DiBiase, T. M. Willis, E. S. Wolak, D. J. Weber, and W. A. Rutala, "Reduction of healthcare-associated infections by exceeding high compliance with hand hygiene practices," *Emerg Infect Dis*, vol. 22, pp. 1628–1630, 2016.
5. N. Lotfinejad, A. Peters, and D. Pittet, "Hand hygiene and the novel coronavirus pandemic: the role of healthcare workers," *Journal of Hospital Infection*, vol. 105, no. 4, pp. 776-777, 2020.
6. WHO, Infection Prevention and Control during Health Care when Novel Coronavirus (nCoV) Infection is Suspected. Interim Guidance, WHO, Geneva, Switzerland, 2020, <https://apps.who.int/iris/rest/bitstreams/1266296/retrieve%202020>.
7. K. M. Cottingim, H. Verma, P. E. Urriola, F. Sampedro, G. C. Shurson, and S. M. Goyal, "Feed additives decrease survival of delta coronavirus in nursery pig diets," *Porcine Health Management*, vol. 3, no. 1, p. 5, 2017.
8. G. Ionidis, J. Hübscher, T. Jack, B. Becker, B. Bischoff, and D. Todt, "Development and virucidal activity of a novel alcohol-based hand disinfectant supplemented with urea and citric acid," *BMC Infectious Diseases*, vol. 16, no. 1, p. 77, 2016.
9. J. de Witt Huberts, K. Greenland, W.-P. Schmidt, and V. Curtis, "Exploring the potential of antimicrobial hand hygiene products in reducing the infectious burden in low-income countries: an integrative review," *American Journal of Infection Control*, vol. 44, no. 7, pp. 764–771, 2016.
10. P. W. Wertz, "Lipids and the permeability and antimicrobial barriers of the skin," *Journal of Lipids*, vol. 2018, Article ID 5954034, 2018.
11. Guertler A, Moellhoff N, Schenck TL, et al. Onset of occupational hand eczema among healthcare workers during the SARS-CoV-2 pandemic: comparing a single surgical site with a

- COVID-19 intensive care unit. *Contact Dermatitis*. 2020;83(2):108–114. doi:10.1111/cod.13618
12. R. Jindal and D. Pandhi, “Hand hygiene practices and risk and prevention of hand eczema during the COVID-19 pandemic,” *Indian Dermatol Online Journal*, vol. 11, no. 4, pp. 540–543, 2020.
 13. C. Beiu, M. M. Mihai, L. Popa, L. Cima, and M. Popescu, “Frequent hand washing for COVID-19 prevention can cause hand dermatitis: management tips,” *Cureus*, vol. 12, 2020.
 14. R. Darlenski and N. Tsankov, “COVID-19 pandemic and the skin: what should dermatologists know?” *Clinics in Dermatology*, 2020, In press.
 15. M. Singh, M. Pawar, A. Bothra, and N. Choudhary, “Over-zealous hand hygiene during the COVID 19 pandemic causing an increased incidence of hand eczema among general population,” *Journal of the American Academy of Dermatology*, vol. 83, no. 1, pp. e37–e41, 2020.
 16. J. Lan, Z. Song, X. Miao, H. Li, Y. Li, and L. Dong, “Skin damage among health care workers managing coronavirus disease-2019,” *Journal of the American Academy of Dermatology*, vol. 82, no. 5, pp. 1215-1216, 2020.
 17. A. Guertler, N. Möllhoff, T. Schenck, C. Hagen, B. Kendziora, and R. Giunta, “Onset of occupational hand eczema among healthcare workers during the SARS-CoV-2 pandemic-comparing a single surgical site with a COVID-19 intensive care unit,” *Contact Dermatitis*, vol. 83, no. 2, pp. 108–114, 2020.
 18. S. Giacalone, P. Bortoluzzi, and G. Nazzaro, “The fear of COVID-19 infection is the main cause of the new diagnoses of hand eczema: report from the frontline in Milan,” *Dermatologic Therapy*, vol. 33, no. 4, p. e13630, 2020.
 19. J. D. Johansen, K. Aalto-Korte, T. Agner, K. E. Andersen, A. Bircher, and M. Bruze, “European Society of contact dermatitis guideline for diagnostic patch testing,” *Recommendations on Best Practice*, vol. 73, no. 4, pp. 195–221, 2015.
 20. C. Patruno, G. Fabbrocini, L. Stingeni, and M. Napolitano, “The role of occupational dermatology in the COVID-19 outbreak,” *Contact Dermatitis*, vol. 83, no. 2, pp. 174-175, 2020.
 21. C. Patruno, S. P. Nistico, G. Fabbrocini, and M. Napolitano, “COVID-19, quarantine, and atopic dermatitis,” *Medical Hypotheses*, vol. 143, Article ID 109852, 2020.
 22. D. Tiedemann, M. L. Clausen, S. M. John, I. Angelova-Fischer, S. Kezic, and T. Agner, “Effect of glove occlusion on the skin barrier,” *Contact Dermatitis*, vol. 74, no. 1, pp. 2–10, 2016.
 23. P. Lin, S. Zhu, and Y. Huang, “Adverse skin reactions among healthcare workers during the coronavirus disease 2019 outbreak: a survey in Wuhan and its surrounding regions,” *British Journal of Dermatology*, vol. 183, no. 1, pp. 190–192, 2020.
 24. G. Cavanagh and C. Wambier, “Rational hand hygiene during COVID-19 pandemic,” *Journal of the American Academic of Dermatology*, vol. 82, no. 6, p. e211, 2020.
 25. T. H. Mekonnen, D. G. Yenealem, and B. M. Tolosa, “Self-report occupational-related contact dermatitis: prevalence and risk factors among healthcare workers in Gondar town, Northwest Ethiopia, 2018—a cross-sectional study,” *Environmental Health and Preventive Medicine*, vol. 24, no. 1, p. 11, 2019.
 26. D. K. Mal and B. Mukherjee, “Educational motivation and legislative approaches for safe and effective hand washes and hand rubs in hand health care during COVID-19 Pandemic,” *Asia Pacific Journal of Public Health*, 2020.
 27. S. J. Pidot, W. Gao, A. H. Buultjens, I. R. Monk, R. Guerillot, and G. P. Carter, “Increasing tolerance of hospital *Enterococcus faecium* to handwash alcohols,” *Science Translational Medicine*, vol. 10, p. 452, 2018.
 28. T. Sandle, “Coronavirus pandemic shortages and the risks of using ineffective hand sanitisers in cleanrooms,” *Clean Air Journals*, 2020.
 29. L. D. Moore, G. Robbins, J. Quinn, and J. W. Arbogast, “The impact of COVID-19 pandemic on hand hygiene performance in hospitals,” *American Journal of Infection Control*, vol. 20, p. 30805, 2020.
 30. D. M. Elston, “Occupational skin disease among health care workers during the coronavirus

(COVID-19) epidemic,” *Journal of the American Academy of Dermatology*, vol. 82, no. 5, pp. 1085-1086, 2020.

31. J. Donovan, I. Kudla, L. D. Holness, S. Skotnicki-Grant, and J. R. Nethercott, “Skin reactions following use of N95 facial masks,” *Dermatitis*, vol. 18, no. 2, p. 104, 2007.
32. Y. Yan, H. Chen, L. Chen, B. Cheng, P. Diao, and L. Dong, “Consensus of Chinese experts on protection of skin and mucous membrane barrier for healthcare workers fighting against coronavirus disease 2019,” *Dermatologic Therapy*, vol. 33, no. 4, p. e13310, 2020.
33. S. Den Boon, C. Vallenas, M. Ferri, and S. L. Norris, “Incorporating health workers’ perspectives into a WHO guideline on personal protective equipment developed during an Ebola virus disease outbreak,” *F1000Research*, vol. 7, p. 45, 2018.
34. B. Bhoyrul, K. Lecamwasam, M. Wilkinson, F. Latheef, S. J. Stocks, and R. Agius, “A review of non-glove personal protective equipment-related occupational dermatoses reported to EPIDERM between 1993 and 2013,” *Contact Dermatitis*, vol. 80, no. 4, pp. 217–221, 2019.

UNDER PEER REVIEW