

Positive Peritoneal Lavage Cytopathology: Prevalence and Relation to Clinico-Pathological Characteristics of Colonic Adenocarcinoma: a Cohort Study

Key words: *Peritoneal carcinomatosis; Colorectal cancer; Free intraperitoneal cancer cells; Peritoneal Lavage Cytopathology.*

Abstract

Background Positive peritoneal lavage cytopathology (PPLC) is an established poor prognostic factor in CRC that has been proven in many studies and was associated with an increase in peritoneal recurrence, overall recurrence and mortality. The incidence of PPLC among CRC patients varied in different studies between 2.1-52%. Similarly, different variables were reported to correlate with peritoneal lavage positivity in published studies. The aim of this trial was to study the prevalence of free intraperitoneal malignant cells at the time of radical resection of colon cancer and to assess its relation to different clinico-pathological characteristics.

Patients and methods This cohort study was conducted at the gastrointestinal surgery unit, general surgery department and the emergency hospital at Tanta University during the period from January 2020 ending in December 2020. Forty patients with non-metastatic primary colonic carcinoma were included in the study. Peritoneal lavage fluid was collected before radical resection of the tumour. One hundred milliliters of warm normal saline solution was installed into the peritoneal cavity and at least 80 ml of the lavage fluid was collected for cytopathological examination. After preparation, the sample was stained with Haematoxylin and Eosin and examined under a light microscope by an experienced cytopathologist.

Results Eighteen smears (45%) were found positive for malignant cells. Among variables tested, a positive correlation was found between PPLC and a low BMI ($< 25 \text{ kg/m}^2$), the histopathological tumour type, an advanced pT stage, a high lymph node ratio and an advanced overall TNM stage.

Conclusion In non-metastatic AJCC stage II and III colon cancer, the prevalence of PPLC detected by conventional cytopathology was found to be 45%. A low BMI $< 25 \text{ kg/m}^2$, the presence of intraperitoneal free fluid at the time of the operation, a high lymph node ratio, an advanced T stage, histopathological type and an advanced overall TNM stage correlated positively with PPLC.

*CRC: colorectal cancer *PM: peritoneal metastases * PC : peritoneal carcinomatosis *PLC : peritoneal lavage cytopathology , *PPLC: positive peritoneal lavage cytopathology , *NPLC : negative peritoneal lavage cytopathology.

Introduction

Colorectal cancer (CRC) is a leading cause of cancer-related death, with peritoneal recurrence occurring in 25-35% of patients (1). Currently, patient selection for adjuvant therapy in CRC does not include routine evaluation of peritoneal lavage fluid for tumour cells (2).

Peritoneal lavage cytology (PLC) has been considered to be a useful tool for predicting an individual prognosis of some malignancies. *Keettel and Elkins, (1956)* (3) introduced the technique of intraoperative peritoneal lavage cytology in ovarian cancer patients for the first time. The presence of free malignant cells in the peritoneal fluid in non-gynecological cancers, especially gastric, colorectal and pancreatic adenocarcinomas, is associated with a poor survival and peritoneal recurrence (4-6). In a prospective study, *Noura et al, 2011* (7) showed an improvement in peritoneal recurrence and overall survival in patients with positive lavage who underwent intraperitoneal chemotherapy with mitomycin C compared to those who did not.

Although free cancer cells can be detected in peritoneal fluid at the time of surgery, peritoneal lavage is not used in routine CRC surgery(8). Several studies, including systematic reviews and meta-analyses, had proven that PPLC is a poor prognostic factor associated with a higher mortality, peritoneal recurrence and overall recurrence in CRC (5, 7, 9).

Testing for free intraperitoneal malignant cells involves examining peritoneal lavage for malignant cells or markers of such cells, using a variety of laboratory methods. Techniques used to date include histological examination (cytopathology), immunocytochemistry and molecular techniques polymerase chain reaction (PCR) (10). The incidence of PPLC among CRC patients varied in different studies between 2.1-52%. Similarly, different variables were reported to correlate with peritoneal lavage positivity in published studies. Factors most commonly associated with PPLC included the depth of tumour invasion (T stage) and the presence of metastases to lymph node, liver or the peritoneum. Controversial factors less commonly reported include lymphatic invasion, venous invasion, tumour stage, tumour grade, tumour necrosis, gross picture of the tumour, and elevated preoperative CA 19.9. No significant association was found between the age, gender or tumor site (5, 11). The aim of this trial was to study the prevalence of free intraperitoneal malignant cells at the time of radical resection of colon cancer and to assess its relation to different clinico-pathological characteristics.

Patient and methods

This cohort study was conducted on 40 patients with colonic carcinoma at the Gastrointestinal Surgery Unit, General Surgery Department and the Emergency Hospital at Tanta University during the period from January 2020 ending in December 2020.

Inclusion criteria: Adult patients with primary non-metastatic colonic carcinoma eligible for radical resection, either emergency or elective, were included in the study.

Exclusion criteria: Patients with any of the followings were excluded from the study:

- Evidence of distant metastasis,
- Evidence of malignant ascites,
- Perforated colonic carcinoma,
- Recurrent colonic carcinoma,
- History of receiving neoadjuvant chemotherapy and/or radiotherapy for malignancy including cancer rectum,
- History of previous abdominal malignancy.

Technique of peritoneal lavage and specimen preparation

In the current study, peritoneal lavage was performed immediately after gaining access to the abdominal cavity through laparotomy incision and before touching the tumour. One hundred milliliter (ml) of warm normal saline solution (temp 37°C) were installed into the peritoneal cavity including the tumor area with the table in the flat position. After gentle hand stirring, as much as possible of this fluid was collected from dependent areas at the Douglas and Morrison pouches. A minimum of 80 ml (80% retrieval) of the lavage fluid was adequate for analysis. If ascites was present, ascitic fluid was aspirated instead of performing a lavage. The specimen was immediately delivered to the pathology department for cytopathological examination. Once received by the pathology department, the sample was centrifuged in Cytospin 2® (Shanton; Athens, Greece; 1200 rotations/min) for 10 minutes. The cell pellet was then smeared on slides, fixed with cytospray, stained with Haematoxylin and Eosin stain (H&E) and examined under a light microscope by an experienced cytopathologist who was blinded to the clinical data. Smears were classified according to their cytopathological features, as follows:

1. *Positive for malignant cells (PPLC):* malignant cells were recognized under light microscope using standard criteria, namely by presence of large, hyperchromatic nuclei, coarse granular nuclear chromatin, prominent nucleolus, distorted nuclear outline, increased nuclear/cytoplasmic ratio, intracytoplasmic mucin vacuoles, or heterogeneity of

cells with clumping of groups. Cells are mainly arranged in loose clusters with the occasional presence of floating cells.

2. *Negative for malignant cells (NPLC)*: normal cells present or cells showing only milder changes of chromatin. Cytopathological findings were graded from class I to V according to The International System (TIS) for reporting serous effusion cytopathology(12).

End points

The primary endpoint was to study the prevalence of PPLC and its relation to the different clinicopathological characteristic in selected colonic carcinoma patients.

Statistical analysis

Collected data were tabulated and analyzed statistically using the appropriate tests, including the mean, standard deviation and were fed to the computer and analyzed using IBM SPSS software package version 20.0. Qualitative data were described using the number and percent. The Kolmogorov-Smirnov test was used to verify the normality of distribution Quantitative data were described using range (minimum and maximum), mean, standard deviation and median. The significance of the obtained results was judged at the 5% level. The used tests were:

1. Chi-square test for categorical variables, to compare between different groups.
2. Fisher's Exact or Monte Carlo correction for Chi-square when more than 20% of the cells have expected count less than 5.
3. Student t-test for normally distributed quantitative variables, to compare between two studied groups

Ethical issues

An informed written consent was obtained from every patient before being enrolled in the study which was approved by the research ethics committee of the Faculty of medicine, Tanta University.

Results

Clinical data:

There were 18 males (45%) and 22 females (55%). The patients' ages ranged between 29 and 85 years with a median of 56.0 (49.0–65.0) years. The BMI ranged between 23 kg/m² and 45 kg/m² with a median of 30.0 kg/m². Five patients (12.5%) had positive family history of malignancy and 2 of them had positive family history of CRC. CEA was elevated in 16 patients (40%) while CA19.9 was elevated in 14 patients (35%). All operations were performed via open approach.

Pathological data:

- **Location of the tumor** 14 patients (35%) had their tumors located in the right side and 24 patients (60%) in the left side. Only one patient had his

tumour located at the transverse colon (2.5%) while another patient had synchronous lesions at the ascending colon and splenic flexure (2.5%). Thirteen patients (32.5%) had malignant large bowel obstruction and were operated in the emergency hospital, while the remaining 27 patients (67.5%) had elective surgery table (1).

- **Size of the tumours:** the maximum tumor size ranged from 1 cm to 14 cm with median 5.0 cm (4.0–7.50).
- **Morphologic types:** 14 patients (35%) had fungating mass, 12 patients (30%) had malignant ulcer and 8 patients (20%) had circumferential (annular) lesions and other 6 patients (15%) had mixed gross patterns.
- **Histopathological types:** conventional adenocarcinoma was found in 26 patients (65%), adenocarcinoma with mucoid differentiation in 10 patients (25%), adenocarcinoma with signet ring differentiation in 3 specimens (7.5%) and one patient had an adenocarcinoma with neuroendocrine differentiation.
- **Tumour grade:** 18 patients (45%) had tumours with poor differentiation (grade III) and 22 patients (55%) had tumours of moderate differentiation (grade II).
- **pT stage:** 27 patients (67.5%) had pT3 tumours, 9 patients (22.5%) had pT4 tumours and 4 patients (10%) had pT2 tumours.
- **pN stage:** the number of harvested LNs ranged from 14 to 31 with a median number of 17 lymph nodes. Sixteen cases (40%) has no LN metastasis (N0 stage), 10 patients (25%) had N1 nodal status, while 14 patients (35%) had N2 nodal status.
- **The overall TNM stage:** 16 patients (40%) had stage II tumours, 24 patients (60%) had stage III tumours Table (2).

Peritoneal fluid analysis

Variable amounts of intraperitoneal free fluid were found in the peritoneal cavity in 7 patients (17.5%) which weren't detectable preoperatively. This fluid was aspirated and analyzed. Six out of these 7 patients (85.7%) had PPLC. The presence of intraperitoneal free fluid was found statistically significant with the positivity of peritoneal fluid for cancer cells in both univariate and multivariate analyses ($p < 0.017$). Peritoneal lavage using normal saline was performed in the remaining 33 patients, which revealed PPLC in 12 (36.5%). Collectively, 18 smears (45%) were found positive for the presence of free malignant cells in the peritoneal fluid on cytopathologic examination (PPLC) while the remaining 22 specimens (55%) had no free malignant cells in the peritoneal fluid (NPLC) (Fig 1).




Fig 1: Positive cytopathology. The cells are large columnar in shape, with moderate amount of vacuolated cytoplasm and have round, hyperchromatic nuclei with coarse chromatin (Low magnification - H&E stain).

Relation between peritoneal lavage cytopathology and different clinicopathological variables (Table 3, 4):

Comparison between patients who had PPLC and those with NPLC showed no statistically significant differences between the 2 groups regarding: gender, positive family history of CRC, clinical presentation, elevated preoperative tumor marker (CEA, CA19.9), tumor size, tumor site, morphologic type, tumour grade or lymph node involvements (pN stage).

On the other hand, comparison between both groups showed a statistically significant differences regarding the BMI, histopathological type, pT stage, the lymph node ratio and the overall TNM stage.

A. Relation between peritoneal lavage cytopathology and different clinical variables (Table 3):

(1) Gender:

PPLC was found in 11 out of 18 male patients (61%), while 7 out of 22 females (31.8%) in our study were found to have positive result (38.9%). No statistically significant difference was found between both gender ($p < 0.064$).

(2) Family history of malignancy:

Among 5 patients with positive family history of malignancy, 4 patients (80%) including the 2 patients with positive family history of CRC had PPLC and 1 patient (20%) had NPLC, no statistically significant difference was found ($p < 0.155$).

(3) BMI:

PPLC was found in 5 out of 6 patients (83.3%) with average BMI $< 25 \text{ kg/m}^2$. On the other hand, NPLC smears were found in 15 out of 21 obese patients (71.4%) BMI $> 30 \text{ kg/m}^2$, while PPLC was found in 7 out of 13

overweight patients. The difference in incidence of PPLC among the patients with different BMI was found statistically significant ($P < 0.044$), with increased PPLC in patients with BMI $< 25 \text{ kg/m}^2$.

(4) Clinical presentation:

Among the 27 patients with elective presentations, 11 patients (40.8%) had PPLC. On the other hand, 7 out of the 13 patients (59.3%) presented to the emergency department with intestinal obstruction had PPLC.

Of all 18 patients with PPLC, 11 patients (61.1%) presented as elective cases with iron deficiency anemia, change in bowel habits, bleeding per rectum, unexplained weight loss while the remaining 7 patients (38.9%) presented with intestinal obstruction. There was no statistically significant difference between both presentations ($p < 0.435$).

(5) Tumor markers:

CEA: Sixteen patients had elevated serum CEA. Ten patients (62.5%) out of these 16 had PPLC while 6 patients (37.5%) had NPLC with no statistically significant difference ($p < 0.069$).

CA19.9: Fourteen patients were found to have elevated serum CA19.9 levels, 9 patients out of them (55.6%) had PPLC, while 5 (45.4%) patients had NPLC with no statistically significant difference ($p < 0.072$).

B. Relation between peritoneal lavage cytopathology and different pathological variables (Table 4):

(1) Tumour size:

The median size of specimens in patients with PPLC was 6.0 cm while it was 5.0 cm in patients with NPLC. The difference in the tumour size between patients with PPLC and those with NPLC didn't attain statistical significance ($p < 0.106$).

(2) The morphologic type of the tumour:

Among different morphologic patterns, the fungating growth pattern had a greater positivity of peritoneal lavage (8 out of 14 patients (57.2%). On the other hand, 4 out of 12 patients with ulcerating tumours (33.3%) had PPLC, while 4 out of 8 patients with circumferential lesions (50%) had PPLC and 2 out of the 6 patients with a mixed growth pattern (33.3%) had PPLC. The difference in incidence of PPLC among the 4 growth patterns was found statistically insignificant ($p < 0.626$).

(3) The Histopathological type of the tumor:

Seven out of 10 peritoneal specimens (70%) in patients with mucoid adenocarcinoma were PPLC, while 2 out of 3 specimens (66.6%) of patients with signet ring carcinoma were found to have PPLC. On the other hand, 8 out of 26 patients with adenocarcinoma (30.8%) had PPLC, while the patient with neuroendocrine differentiation had PPLC. The difference in incidence of

PPLC among these 4 different histopathological types was found statistically significant ($p < 0.047$). The more aggressive the tumour histopathologically is, the higher the possibility of PPLC.

(4) Tumour grade:

Eleven out of 18 patients (61.1%) with poorly differentiated adenocarcinomas (Grade 3) had PPLC, while 7 out of 22 patients (31.8 %) with moderately differentiated tumours (Grade 2) had a lower probability of PPLC. Although the possibility of PPLC increased with decreased tumour differentiation, the difference in incidence of PPLC between these 2 different grades didn't achieve statistical significance ($p < 0.064$).

(5) pT-stage:

Increasing the depth of invasion of malignant cell into the wall of the colon was associated with a higher possibility of PPLC. The four pT2 tumours had a NPLC. In pT3 tumours, on the other hand, 11 (40.8%) out of 27 specimens had a PPLC, while in pT4 tumours, 7 out of 9 specimens (77.8%) with were found to have a PPLC. The difference among the 3 categories regarding the positivity of specimen was found statistically significant ($p < 0.025$).

(6) pN- stage:

Four out of 16 patients (25%) with no LN metastasis (pN0) had PPLC. Six out of 10 patients (60%) with pN1 had PPLC, while, 8 out of 14 patients (57.1%) with N2 had PPLC. The difference among the 3 categories regarding the positivity of peritoneal lavage was found statistically insignificant ($p < 0.115$).

(7) Lymph node ratio:

In patients with PPLC, the range of lymph node ratio was 0-95.83% with a mean of 27.80 ± 23.55 (median 27.8). In patients with NPLC, the range of lymph node ratio was 0-50.94% with a mean of 12.40 ± 16.21 (median 10.0). The difference between PPLC and NPLC regarding the lymph node ratio was found statistically significant ($p < 0.026$).

(8) TNM stage:

Fourteen out of 24 patients with stage III (58.3%) had PPLC, while only 4 out of 16 patients (25%) with stage II had positive cytology. The difference between the 2 stages was found statistically significant ($p < 0.038$). This meaning that the positivity of lavage analysis increased with increasing the overall TNM stage of the tumours .

(9) Lymphatic, venous and perineural invasion:

Of 39 patients with lymphatic invasion, 18 patients (42.2%) had PPLC while the remaining 21 patients (57.8%) had NPLC. The difference was found statistically insignificant ($p < 1,000$). Seventeen out of 37 patients with venous invasion (45.9%) had PPLC and 20 patients (54.1%) had NPLC. The difference was statistically insignificant ($p < 1.000$). Finally, 16 out of 37 patients with perineural invasion (47.2%) had PPLC while the remaining 19 patients (52.8%) had NPLC. The

difference was found statistically insignificant ($p < 0.613$) Table (4).

Factors associated with positivity of peritoneal lavage cytopathology

We performed univariate and multivariate analyses including 20 different independent clinical and pathological variables which could have a potential correlation to PPLC. In the univariate analysis BMI, presence of intraperitoneal free fluid at exploration, lymph node ratio and depth of tumour invasion through colonic wall (pT stage) correlated positively with PPLC. In multivariate analysis, BMI, intraperitoneal free fluid and lymph node ratio were independent variables correlating with PPLC (Table 5).

Table (1) Patients' demographic data

Demographic data	No (40)	%
Sex		
Male	18	45.0
Female	22	55.0
Age (years)		
Range	29.0 – 85.0	
Mean \pm SD.	57.90 \pm 14.72	
Median (IQR)	56.0 (49.0–65.0)	
BMI (kg/m²)		
<25 (Normal)	6	15.0
>25 – <30 (Overweight)	13	32.5
>30 (Obese)	21	52.5
Min. – Max.	23.0 – 45.0	
Mean \pm SD.	29.56 \pm 5.42	
Median (IQR)	30.0 (25.0–31.0)	
Family history of malignancy		
Negative	35	87.5
Positive	5	12.5

Table (2): Postoperative Pathological data

Pathological data	No (40)	%
Tumor size (cm)		
Range	1– 14.0	
Mean	5.76 \pm 2.89	
Median	5.0 (4.0–7.50)	
Morphologic type		
Fungating mass	14	35.0
Malignant ulcer	12	30.0
Circumferential lesion	8	20.0
Mixed	6	15.0
Histopathologic type		
Adenocarcinoma	26	65.0
Adenocarcinoma with mucoid differentiation	10	25.0
Adenocarcinoma with signet ring differentiation	3	7.5
Adenocarcinoma with neuroendocrine differentiation	1	2.5
Grade		
II	22	55.0
III	18	45.0
pT stage		
T2	4	10.0
T3	27	67.5
T4	9	22.5

pN stage		
N0	16	40.0
N1	10	25.0
N2	14	35.0
Range		14-31
Mean ± SD		16.7 ±6.23
Median		17
Overall TNM stage		
Stage II	16	(40%)
Stage III	24	(60%)
Intraperitoneal free fluid	33	82.5
Absent	7	17.5
Present		
Peritoneal lavage analysis	22	55
NPLC	18	45
PPLC		

Table (3): Peritoneal lavage cytopathology status and different clinical variables

	PLC				Test of Sig	P
	-ve (n=22)		+ve (n=18)			
	No.	%	No.	%		
Gender						
Male	7	38.9	11	61.1	$\chi^2=3.432$	0.064
Female	15	68.2	7	31.8		
BMI (kg/m²)						
Normal	1	16.7	5	83.3	$\chi^2=6.030^*$	0.044*
Overweight	6	46.23	7	53.8		
Obese	15	71.4	6	28.6		
Range	24.0 – 45.0		23.0 – 43.0		t= 1.464	0.153
Mean ± SD.	30.70 ± 4.80		28.17 ± 5.93			
Median	30.0		25.0			
Family history						
Positive	1	20	4	80	$\chi^2=2.828$	FE p=0.155
Negative	21	60	14	40		
Clinical presentation						
Elective					$\chi^2=0.609$	0.435
Emergency	16	59.2	11	40.8		
	6	46.2	7	53.8		
Tumour Marker						
CEA						
Normal	16	66.7	8	33.3	$\chi^2=3.300$	0.069
Elevated	6	37.5	10	62.5		
CA19.9						
Normal	17	65.4	9	34.6	$\chi^2=3.237$	0.072
Elevated	5	35.7	9	64.3		

χ^2 Chi square test

MC Monte Carlo

FE Fisher Exact

t: Student t-test

P: p value for comparing between the two studied categories

*: Statistically significant at $p \leq 0.05$

PLC Peritoneal lavage cytopathology, CEA: carcinoembryonic antigen, CA19.9: Cancer Antigen, BMI: body mass index

Table (4): Peritoneal lavage cytopathology status and different pathological variables

	PLC				Test of Sig.	P
	-ve (n=22)		+ve (n=18)			
	No.	%	No.	%		
Morphologic type						
Fungating mass	6	42.8	8	57.2	$\chi^2=1.940$	MC p=0.626
Malignant ulcer	8	66.7	4	33.3		
Circumferential lesion						
Mixed	4	50	4	50		
	4	66.7	2	33.3		
Tumor size cm (max)						
Range	1.0 – 12.0		1.80 – 14.0		U=131.50	0.106
Mean ± SD.	5.10 ± 2.72		6.54 ± 2.97			
Median	5.0		6.0			
Grade						
II	15	68.2	7	31.8	$\chi^2=3.432$	0.064
III	7	38.8	11	61.2		
Histopathological type						
• Adenocarcinoma	18	69.2	8	30.8	$\chi^2=6.317^*$	MC p=0.047*
• Adenocarcinoma with mucoid differentiation	3	30.0	7	70.0		
• Adenocarcinoma with signet ring	1	66.6	2	33.3		

differentiation						
• Adenocarcinoma with neuroendocrine differentiation	0	0.0	1	100.0		
T-stage					$\chi^2=$	MC p=
T2	4	100.0	0	0.0	7.384*	0.025*
T3	16	59.2	11	40.8		
T4	2	22.2	7	77.8		
N stage					4.329	0.115
N0	12	75.0	4	25.0		
N1	4	40.0	6	60.0		
N2	6	42.9	8	57.1		
Lymph node ratio					U=	0.026*
Range	0 – 50.94		0 – 95.83		116.0	
Mean	12.40 ± 16.21		27.80 ± 23.55			
Median	10		27.78			
Stage						
II	12	75.0	4	25.0	$\chi^2=$	0.038*
III	10	41.7	14	58.3	4.313	
Lymphatic invasion	21	53.8	18	46.2	0.839	1.000
Venous invasion	20	54.1	17	45.9	0.178	1.000
Perineural invasion	19	52.8	17	47.2	0.718	0.613

χ^2 : Chi Square test; MC: Monte Carlo U: Mann Whitney test; FE: Fisher Exact p: p value for comparing between two categories; *: Statistically significant at $p \leq 0.05$

PLC Peritoneal lavage cytopathology

Table (5) Factors associated with positive peritoneal lavage cytopathology

Independent variable	Univariate analysis		Multivariate analysis	
	OR (95% CI)	P value	OR (95% CI)	P value
Sex	3.367 (0.913–12.417)	0.109		
Age	0.554 (0.157–1.952)	0.356		
BMI	4.286 (1.135–16.182)	0.022*	5.634 (2.547–11.647)	0.037*
Family history of CRC	2.375 (1.636–3.448)	0.109		
CEA level	0.300 (0.080–1.124)	0.114		
CA 19.9 level	0.294 (0.076–1.145)	0.126		
Free fluid	10.500 (1.126–17.908)	0.017*	8.625 (2.637–14.527)	0.024*
Tumour location	0.764 (0.215–2.708)	0.429		
Tumor size	0.589 (0.155–2.236)	0.457		
Morphologic type	8.521 (0.627–14.523)	0.671		
Histopathologic type	3.367 (0.913–12.5417)	0.217		
Tumour grade	0.297 (0.081–.095)	0.174		
pT stage	0.461 (0.103–0.617)	0.037*	0.685 (0.216–7.521)	0.327
pN stage	0.528 (0.167–4.521)	0.318		
Lymph node ratio	0.654 (0.204–0.857)	0.019*	0.348 (0.127–0.647)	0.041*
Overall TNM Stage	0.851 (0.318–7.521)	0.294		
Lymphatic invasion	0.526 (0.389–8.712)	0.317		
Venous invasion	1.700 (0.142–20.422)	0.719		
Perineural invasion	2.684 (0.254–28.311)	0.461		

Discussion

The objective of our study was to detect those free intraperitoneal cancer cells (IPCC) by conventional cytopathologic examination of the peritoneal lavage fluid obtained from a select cohort of colonic carcinoma patients undergoing radical resection and to define both clinical and pathological variables that correlate positively with their presence.

Since the objective of our study was to detect the prevalence of PPLC in relation to clinical and pathological variables, we performed pre-resection peritoneal lavage before tumour manipulation, as described in literature (4, 13-17). Conventional cytopathology was used in our study, because it is a universal and inexpensive method that can be easily performed at any institution (11). Since no significant difference in the positivity of PLC was observed based on the type of stain used, H&E stain was the preferred stain employed in the current study.

Our study showed that the prevalence of PPLC was 45%. The reported prevalence of free IPCC on conventional cytopathology showed a wide range of positivity among published studies. *Nishikawa et al*, (11) published a meta-analysis of 18 studies, 15 studies employed conventional cytopathology as one of techniques used to detect free IPCC. In studies using conventional cytopathology as the sole method for detection of free IPCC, the detection rate ranged from 0 to 35.5%. In the meta-analysis of *Passot et al*, (8), the yield rate of positive IPCC detection by conventional cytopathology varied from 4% to 35.5%.

This wide range of outcomes can be attributed to the marked heterogeneity among studies concerning different aspect of the technique including the volume of lavage fluid, the type of lavage fluid, timing of performing lavage, the volume of retrieved fluid, the different staining techniques employed for detection of free IPCC and the varied experience of cytopathologists involved in these studies. Also, the selection criteria of the included patients showed significant heterogeneity among studies. For these reasons, comparing our results with published results is difficult and should be interpreted cautiously.

Three studies employed technical steps very close to ours including pre-resection sample retrieval and using H&E in conventional cytopathology. The prevalence of PPLC reported in these studies was 35.5% (32/90), 7.9% (15/189) and 15.8% (23/145) of patients reported by *Vogel et al*, (18), *Lee et al*, (19) and *Temesi et al*, (20) respectively. Our results showed a higher prevalence of PPLC than these studies. Looking at the pathological results in our study may explain this higher outcome, as most of the patients had unfavorable pathology; about one third had mucinous and signet ring differentiation which behave less favorably than conventional adenocarcinoma. Also, with conventional

adenocarcinoma, all patients have either moderately or poorly differentiated tumours. We have no T1 tumour and about one fourth of our patients has tumour reaching the serosal surface; T4. Moreover, 60% of our patients had advanced stage III cancer. All these pathological variables including; histopathological type, depth of tumour invasion, higher lymph node ratio and overall TNM staging revealed higher statistical significant correlation with PPLC in our study.

To the best of our knowledge, no prospective study had investigated the relation between the BMI and the presence of free IPCC in colonic carcinoma patients. Investigating this relation in our study, we found that 21 out of the 22 patients (95%) with NPLC were overweight or obese and only 1 patient (5%) had a normal BMI, while in patients with PPLC, 13 out of 18 patients (72.2%) were either overweight or obese and 5 patients (27.8%) had a normal BMI. The difference in incidence of PPLC among the patients with different BMI was found statistically significant ($P < 0.044$), with increased PPLC in patients with BMI $< 25 \text{ kg/m}^2$. This outcome was confirmed in both univariate and multivariate analysis. Thus, a higher BMI seems to play a protective role against spread of free cancer cells within the peritoneal cavity. Recently in a retrospective analysis done by *Aldaqa et al*, (21) investigating the relation between obesity and clinicopathologic characteristics and prognosis of 233 patients with CRC, they found that obese patients had the lowest incidence of stage IV disease (17 of 69 patients; 25.8%). Meanwhile, the percentage of stage IV disease increased by decreasing the BMI; 38.7%, 39.2% and 57.1% for overweight, normal and underweight patients respectively. The authors concluded that underweight patients may have the worst prognosis. The data of this study supported our observation that a higher BMI may have a protective role against spread of free cancer cells within the peritoneal cavity.

The presence of the malignant cells in the lavage fluid in the current study increased with increasing the depth of tumour invasion (pT stage) into the bowel wall ($p = 0.032$). All 4 patients with pT2 tumors had NPC while, (7/9) of patients with pT4 tumours (78%) had PPLC. In the univariate analysis, the pT-stage showed a correlation with PPLC. Our results coincided with those reported widely in the literature (4, 13-15, 22, 23) in spite of using another stain in their conventional cytopathology. Also our results agree with the similar three clinical trials (18-20).

Concerning the histopathological types in our patients, 7 out of 10 peritoneal lavage specimens (70%) in patients with mucoid adenocarcinoma had PPLC, while 2 out of 3 specimens (66.6%) of patients with signet ring carcinoma were found to have PPLC. On the other hand, 8 out 26 patients with conventional

adenocarcinoma (30.8%) had PPLC and the difference among these different histopathological types was found statistically significant ($p < 0.047$). The more aggressive the tumour histopathologically is, the higher the possibility of PPLC. On studying the correlation between tumor grade and PPLC, we found that the possibility of PPLC increased with decreased tumour differentiation (poorly differentiated adenocarcinomas (Grade 3) was (61.1%) and moderately differentiated tumours (Grade 2) was (31.8 %)), although this difference in incidence of PPLC between these 2 different grades didn't achieve statistical significance ($p < 0.064$). These outcomes in our study coincided with *Hara et al*(22), *Nishikawa et al*,(15) ; *Lee et al*(19), who found a positive correlation between higher tumour grade and positive lavage .

Looking to the pN stage, no positive relation between this variable and PPLC was found in our study ($p = 0.115$). This was in agreement with what was reported by *Hase et al*,(24), *Yamamoto et al*,(2) and *Lloyd et al*, (25). However, in multivariate analysis, *Hara et al*;(22) *Noura et al*, (4); *Katoh et al*,(17) and *Temesi et al*, (20) reported a strong correlation between this variable and PPLC . It is noteworthy, however, that in the current study, a positive correlation was found between PPLC and lymph node ratio. Patients with PPLC had a lymph node ratio of 27.80 ± 23.55 in comparison to 12.40 ± 16.21 in patients with NPLC and this difference was found statistically significant ($p < 0.026$) in both univariate and multivariate analysis. To the best of our knowledge, this variable wasn't investigated before in other studies.

In this study, 14 out of 24 patients with stage III (58.3%) had PPLC, while only 4 out of 16 patients (25%) with stage II had positive cytopathology. The difference between the 2 stages was found statistically significant ($p < 0.038$). This means that, the positivity of lavage analysis increased with increasing the overall TNM stage of the tumours. *Lee et al*, (19) reported similar results with 13.3% PPLC in stage II, 40% in stage III and 46.7% in stage IV which were statistically significant ($p < 0.001$). Also, *Kanellos et al*,(13) and *Noura et al*, (4) found this positive correlation between tumor stage and PPLC.

In the current study, 6 out of the 7 patients (85.7%) who had free fluid at exploration were found to have PPLC . The presence of free fluid in the peritoneal cavity at the time of operation showed a positive correlation with PPLC in both univariate and multivariate analyses. Our results are supported by a retrospective study recently published by *Sato et al*, (26) who looked for free malignant cells in the minimal amount of free fluid found at surgery ($n = 225$) and in the peritoneal lavage fluid ($n = 367$) in 592 clinical

stage II–IV CRC patients. The positivity rate was 17.8% versus 6.5% in peritoneal cytology and PLC groups respectively ($p < 0.001$). Patients with free fluid were younger, had more advanced and more aggressive disease with a higher incidence of emergency presentation. Our results were also quite similar to those reported by *Hase et al*,(24) and *Gozalan et al*,(14), who found that the presence of malignant cells in the peritoneal cavity can enhance fluid production by inducing peritoneal inflammation and hinder re-absorption by obstructing the stomata and lymphatic channels. Again, comparison here should be approached cautiously; we excluded patients with macroscopic metastasis discovered pre- or intra-operatively, while, these 2 cohorts included patients with gross peritoneal metastasis. *Lee et al*, (19), on the other hand, found a positive correlation between macroscopic peritoneal metastasis and PPLC, but, they didn't find correlation between peritoneal effusion and PPLC.

In our study, venous invasion, lymphatic invasion and perineural invasion didn't show correlation with PPLC. This was coincided with the results of *Wind et al*;(27) *Gozalan et al*,(14) and *Fujii et al*,(16) . *Hase et al*, (24) found lymphatic invasion to be correlated with PPLC. *Homma et al*, (23) found venous invasion to be correlated with PPLC while lymphatic invasion didn't.

The detection of free IPCC by PLC at the time of surgery has been reported to be one of the most accurate prognostic factors in some gastrointestinal and gynaecologic malignancies. It has been especially well studied in gastric adenocarcinoma (18, 28). According to the AJCC TNM classification, gastric cancer with PPLC is classified as Stage IV (28). Moreover, PPLC has been shown as a useful prognostic marker for pancreatic, esophagogastric, and gynecological malignancies (29, 30). However, in CRC, the meaning of PPLC is still controversial (23), therefore, the technique of PLC till now is not a standard procedure during resection of CRC and the presence of free IPCC does not currently influence the staging of CRC or the decision regarding the use of adjuvant therapy(11). *Bae et al*, (31) suggested that PLC must be a useful tool for selecting patients who are in need for IPC. Also *Sato et al*, (26) suggested that PLC should be a standard assessment modality for CRC that may adopt intraperitoneal chemotherapy and hyperthermic intraperitoneal chemotherapy as mandatory additional treatment after radical resection.

Conclusion

In non-metastatic AJCC stage II and III colon cancer, the prevalence of PPLC detected by conventional cytopathology was found to be 45%. BMI $< 25 \text{ kg/m}^2$,

the presence of intraperitoneal free fluid at the time of the operation, a high lymph node ratio, an advanced T stage, the histopathological type and an advanced overall TNM stage correlated positively with PPLC. During the follow-up period, PPLC was associated with a higher recurrence and mortality rates.

References

1. Gellad ZF, Provenzale D. Colorectal cancer: national and international perspective on the burden of disease and public health impact. *Gastroenterology*. 2010;138(6):2177-90.
2. Yamamoto S, Akasu T, Fujita S, Moriya Y. Long-term prognostic value of conventional peritoneal cytology after curative resection for colorectal carcinoma. *Japanese journal of clinical oncology*. 2003;33(1):33-7.
3. Keettel WC, Elkins H. Experience with radioactive colloidal gold in the treatment of ovarian carcinoma. *American Journal of Obstetrics & Gynecology*. 1956;71(3):553-68.
4. Noura S, Ohue M, Seki Y, Yano M, Ishikawa O, Kameyama M. Long-term prognostic value of conventional peritoneal lavage cytology in patients undergoing curative colorectal cancer resection. *Diseases of the colon & rectum*. 2009;52(7):1312-20.
5. Mohan HM, O'Connor DB, O'Riordan JM, Winter DC. Prognostic significance of detection of microscopic peritoneal disease in colorectal cancer: a systematic review. *Surgical oncology*. 2013;22(2):e1-e6.
6. Bosanquet D, Harris D, Evans M, Beynon J. Reply to: lack of prognostic significance of conventional peritoneal cytology in colorectal and gastric cancers: results of EVOCAPE 2 multicentre prospective study. *European Journal of Surgical Oncology: the Journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology*. 2014;40(5):596-7.
7. Noura S, Ohue M, Shingai T, Kano S, Ohigashi H, Yano M, et al. Effects of intraperitoneal chemotherapy with mitomycin C on the prevention of peritoneal recurrence in colorectal cancer patients with positive peritoneal lavage cytology findings. *Annals of surgical oncology*. 2011;18(2):396-404.
8. Passot G, Mohkam K, Cotte E, Glehen O. Intra-operative peritoneal lavage for colorectal cancer. *World Journal of Gastroenterology: WJG*. 2014;20(8):1935.
9. Rekhraj S, Aziz O, Prabhudesai S, Zacharakis E, Mohr F, Athanasiou T, et al. Can intra-operative intraperitoneal free cancer cell detection techniques identify patients at higher recurrence risk following curative colorectal cancer resection: a meta-analysis. *Annals of surgical oncology*. 2008;15(1):60-8.
10. Bosanquet D, Harris D, Evans M, Beynon J. Systematic review and meta-analysis of intraoperative peritoneal lavage for colorectal cancer staging. *Journal of British Surgery*. 2013;100(7):853-62.
11. Nishikawa T, Sunami E, Tanaka T, Tanaka J, Kiyomatsu T, Kawai K, et al. Incidence and prognostic significance of positive peritoneal lavage in colorectal cancer. *Surgery today*. 2015;45(9):1073-81.
12. Koss L, Melamed M. From Diagnostic cytology: Its origin and principles. *Koss' Diagnostic cytology and its histopathologic basis Philadelphia: Lipincott Williams and Wilkins*. 2006:3-20.
13. Kanellos I, Demetriades H, Zintzaras E, Mandrali A, Mantzoros I, Betsis D. Incidence and prognostic value of positive peritoneal cytology in colorectal cancer. *Diseases of the colon & rectum*. 2003;46(4):535-9.
14. Gozalan U, Yasti AC, Yuksek YN, Reis E, Kama NA. Peritoneal cytology in colorectal cancer: incidence and prognostic value. *The American journal of surgery*. 2007;193(6):672-5.
15. Nishikawa T, Watanabe T, Sunami E, Tsuno NH, Kitayama J, Nagawa H. Prognostic value of peritoneal cytology and the combination of peritoneal cytology and peritoneal dissemination in colorectal cancer. *Diseases of the colon & rectum*. 2009;52(12):2016-21.
16. Fujii S, Shimada H, Yamagishi S, Ota M, Kunisaki C, Ike H, et al. Evaluation of intraperitoneal lavage cytology before colorectal cancer resection.

International journal of colorectal disease. 2009;24(8):907-14.

17. Katoh H, Yamashita K, Sato T, Ozawa H, Nakamura T, Watanabe M. Prognostic significance of peritoneal tumour cells identified at surgery for colorectal cancer. *British journal of surgery*. 2009;96(7):769-77.

18. Vogel I, Kalthoff H. Disseminated tumour cells. *Virchows Archiv*. 2001;439(2):109-17.

19. Lee IK, Kim DH, Gorden DL, Lee YS, Sung NY, Park G-S, et al. Prognostic value of CEA and CA 19-9 tumor markers combined with cytology from peritoneal fluid in colorectal cancer. *Annals of surgical oncology*. 2009;16(4):861-70.

20. Temesi R, Sikorszki L, Bezsilla J, Botos Á, Kovács J, Tihanyi T. Impact of positive intraabdominal lavage cytology on the long-term prognosis of colorectal cancer patients. *World journal of surgery*. 2012;36(11):2714-21.

21. Aldaqal SM, Maqbul AA, Alhammad AA, Alghamdi AS, Alharbi BA, Alharbi MT, et al. The Impact of Body Mass Index on the Clinicopathological and Prognostic Factors of Colorectal Cancer in Saudi Arabia. *Cureus*. 2020;12(11).

22. Hara M, Nakanishi H, Jun Q, Kanemitsu Y, Ito S, Mochizuki Y, et al. Comparative analysis of intraperitoneal minimal free cancer cells between colorectal and gastric cancer patients using quantitative RT-PCR: possible reason for rare peritoneal recurrence in colorectal cancer. *Clinical & experimental metastasis*. 2007;24(3):179-89.

23. Homma Y, Hamano T, Akazawa Y, Otsuki Y, Shimizu S, Kobayashi H, et al. Positive peritoneal washing cytology is a potential risk factor for the recurrence of curatively resected colorectal cancer. *Surgery today*. 2014;44(6):1084-9.

24. Hase K, Ueno H, Kuranaga N, Utsunomiya K, Kanabe S, Mochizuki H. Intraperitoneal exfoliated cancer cells in patients with colorectal cancer. *Diseases of the colon & rectum*. 1998;41(9):1134-40.

25. Lloyd JM, McIver CM, Stephenson S-A, Hewett PJ, Rieger N, Hardingham JE. Identification of early-stage colorectal cancer patients at risk of relapse

post-resection by immunobead reverse transcription-PCR analysis of peritoneal lavage fluid for malignant cells. *Clinical Cancer Research*. 2006;12(2):417-23.

26. Sato K, Imaizumi K, Kasajima H, Kurushima M, Umehara M, Tsuruga Y, et al. Comparison of prognostic impact between positive intraoperative peritoneal and lavage cytologies in colorectal cancer. *International Journal of Clinical Oncology*. 2021;26(7):1272-84.

27. Wind P, Nordlinger B, Roger V, Kahlil A, Guin E, Parc R. Long-term prognostic value of positive peritoneal washing in colon cancer. *Scandinavian journal of gastroenterology*. 1999;34(6):606-10.

28. Kobayashi H, Kotake K, Sugihara K. Prognostic significance of peritoneal lavage cytology in patients with colorectal cancer. *International journal of clinical oncology*. 2013;18(3):411-7.

29. Ishikawa O, Wada H, Ohigashi H, Doki Y, Yokoyama S, Noura S, et al. Postoperative cytology for drained fluid from the pancreatic bed after "curative" resection of pancreatic cancers: does it predict both the patient's prognosis and the site of cancer recurrence? *Annals of surgery*. 2003;238(1):103.

30. Kuji S, Hirashima Y, Komeda S, Tanaka A, Abe M, Takahashi N, et al. The relationship between positive peritoneal cytology and the prognosis of patients with FIGO stage I/II uterine cervical cancer. *Journal of Gynecologic Oncology*. 2014;25(2):90-6.

31. Bae SJ, Shin US, Ki Y-J, Cho SS, Moon SM, Park SH. Role of peritoneal lavage cytology and prediction of prognosis and peritoneal recurrence after curative surgery for colorectal cancer. *Annals of Coloproctology*. 2014;30(6):266.

UNDER PEER REVIEW