

Assessment of Arsenic (As), Lead (Pb) and Chromium (Cr) accumulation in different organs of commercially important fish species collected from Chattogram coastal region of Bangladesh.

ABSTRACT

Aims: The purpose of this study was to evaluate whether there are any significant toxic effects of the widely exposed metals on different organs (gill, liver, kidney, and muscle) of some commercially important marine fish species (*Herpodon nehereus*, *Pampus chinensis* and *Hilsa ilisha*).

Study design: The collection of the samples was done from January to March, 2018 from Bay of Bengal near Chattogram city of Bangladesh. Analyses were performed for metals such as arsenic (As), lead (Pb) and chromium (Cr) using Atomic Absorption Spectrophotometer (AAS).

Methodology: During each month, five individuals from each three species were collected, with a total of 45 fishes in the study period.

Results: The concentrations of the metals found in the fishes varied in the following ranges: As: 0.029-0.071 ppm, Pb: 0.008-0.083 ppm and Cr: 0.0001-0.03 ppm. The obtained result revealed the highest concentrations of all three metals were recorded in *H. nehereus* fish. The organ wise lead concentration was recorded highest in kidneys and gills of examined fishes with significant variation in muscles and liver. The scenario depicted quite differently in case of chromium where concentration in gills was found to be the highest with insignificant variation in accumulation in other three organs. In case of Arsenic, kidneys and livers were the most exposed two organs in comparison to significant exposure to muscles and gills. The obtained values of arsenic accumulation were considered critical for human consumption as it was higher than the recommended values stated by WHO and FAO; but the concentration of lead and chromium were found to be safe for human consumption. The values recorded for lead were increasingly approached to the safety values.

Conclusion: So, crucial steps should be taken regarding safety and environmentally friendly discharge of arsenic and lead; high levels of pollution will not only affect aquatic life but will also invite socio-economic disasters.

Keywords: Metals; Bay of Bengal; Fish organs; Bioaccumulation; Atomic absorption spectroscopy; Pollution.

1. INTRODUCTION

In Bangladesh, fish is a popular animal-source food in the diet of millions. It has been advised that fish should be consumed two or three times weekly, because of its high levels of essential amino acids (EAA), polyunsaturated fatty acid such as omega-3 and omega-6 fatty acids [1,2]. The coastal water of Bangladesh comprises diverse fisheries resources with 475 finfish species [3]. The marine fisheries play significant role in the economy of Bangladesh. Total production of marine fisheries is 6.71 lakh MT and its contribution to total fish production is 14.90% with growth rate 1.70% [4]. In 2019-2020, fisheries and aquaculture sectors have become the second most important contributors in export earnings of Bangladesh, contributing to 3.52% in national GDP, 1.39% in export earnings and 26.37% in agriculture sector [4].

But now-a-days metal pollution of the marine environment has long been recognized as a serious environmental concern. Heavy metals are present in the aquatic environment where they bio accumulate along the food chain. Accumulation occurs in the organs of aquatic animals and may become toxic for fish and for people when it reaches a substantially high level. The organs of fish that take up heavy metals are the gills, skin, kidney, digestive tract, muscle, and liver [5]. The potential sources of metal pollution in the aquatic environment are agriculture, shipping, urban, industrial wastes, and mining [6]. Metals like arsenic, cadmium, chromium, mercury, nickel, and lead are often considered indicators of anthropogenic influence in marine environment and are themselves of potential risk to the natural environment [7].

In many studies, fishes were the subject of investigation on metal accumulations and monitoring programs in seas or fresh water, due to their importance in human nutrition [8,9,10]. Such interest aimed at ensuring the safety to the food supply and minimizing the potential hazard effect on human health. The concentration of arsenic in muscle tissue of marine fish species collected from northeastern Bay of Bengal, India was studied [11,12]. They found considerable variation of arsenic levels among the fish species which were higher than the standard value of arsenic. As a commercial hub and the industrial nerve center of Bangladesh, Chattogram coastline of Bay of Bengal is at risk of having high levels of metal pollutants due to surrounding activities. According to studies [10,13], many kinds of industries are situated on Chattogram coast, each discharging its characteristic range of effluents containing metals and other harmful pollutants into the terrestrial and aquatic ecosystems. There are some studies on the pollution level of metals in water and sediments along the Karnafuli River and its adjacent coastal area [14,15] but information on the types of metal in the Bay of Bengal coast including fish, other ecosystems, coastal areas, and rivers is poorly documented. With realizing the significance, the principal objectives of this research were: i) to determine the Arsenic (As), Lead (Pb) and Chromium (Cr) accumulation in different commercially important marine fishes collected from Chattogram coastal area. ii) to develop a comparison among different organs (Gill, Liver, Kidney and Muscle) and different species of investigated fishes in terms of those metal accumulation.

2. MATERIALS AND METHODS

2.1 Fish collection

Fishes were collected between January and March 2018 from the caught fishes by fishermen's nets from the Chattogram coast of Bay of Bengal. During each month, five individuals from each three species (*H. nehereus*, *P. chinensis* and *H. ilisha*) were collected, with a total of 45 (forty-five) fishes in the study period. Each time the fishes were transported to the Quality Control and Analytical Laboratory of Chattogram Veterinary and Animal Sciences University, Chattogram, Bangladesh. Fishes were stored in plastic bags at -20° C until dissection.

2.2 Fish dissection and preparation of sample for digestion

The collected fishes were separated according to their species and separately dissected for their muscle, gill, liver and kidney tissues. For example, 15 (5*3) muscle tissues for *H. nehereus* were collected during the three months of study period (Jan-Mar 2018). All the final sample preparation and chemical analysis for As, Pb and Cr were carried out according to the procedure described by UNEP Reference Methods [16]. The prepared samples were stored in 10% formalin solution for future analysis.

2.3 Digestion

The selected tissues were digested with concentrated nitric acid and perchloric acid (2:1 v/v) at 60 °C for 3 days and all samples were diluted with double distilled water by a microwave digester. Following acid digestion, all samples were analyzed for As, Pb and Cr by atomic absorption spectrometry (Phillips AAS with double beam and deuterium background corrector).

2.4 Arsenic (As), Lead (Pb) and Chromium (Cr) Analysis

As, Pb and Cr were analyzed in a graphite furnace (GBCGF 3000 with Zeeman background corrector) with an auto sampler. All digested samples were analyzed three times for each metal. The standard addition method was used to correct for matrix effects (If any). The instrument was calibrated with standard solutions prepared from commercial materials. Analytical blanks were run in the same way as the samples and determined using standard solutions prepared in the same acid matrix.

2.5 Data Analysis

Statistical analysis was carried out using SPSS 20. One-way analysis of variance (ANOVA) and least significant difference (LSD) test were used to assess whether metal concentrations varied significantly among species & organs. The average concentration of As, Pb and Cr among each of the muscle, gill and kidney tissues collected from each species were used. The comparative accumulation of Pb, Cr and As in each species was demonstrated by using Microsoft Excel. The level of significance of the mean values was assigned at $P < .05$.

3. RESULTS AND DISCUSSION

This study investigated As, Cr and Pb distribution in different organs (gill, liver, muscle, and kidney) of three fish species (*H. nehereus*, *P. chinensis* and *H. ilisha*) captured from Chattogram coast of Bay of Bengal. These fish species were selected, as they are most consumed and commercially important marine fishes in Bangladesh. As, Pb and Cr were chosen because these are common detrimental elements found in industrial discharges.

The average of As, Pb and Cr in different organs of the investigated species were presented in Figure 1, Figure 2 and Figure 3. The comparisons of average metal contents of the organs and the species were shown in Table 1 and Table 2.

3.1 Arsenic (As), Lead (Pb) and Chromium (Cr) Concentration in *H. nehereus*

H. nehereus economically important and they form a large part of the fish catch in the study area. The recorded values in *H. nehereus* expressed that the average value of arsenic obtained from muscle (0.039 ppm) was significantly ($P<.05$) different from the values of gill (0.053 ppm), liver (0.064 ppm) and kidney (0.071 ppm) which are higher than the recommended value of 0.01 ppm for fish [17] (Figure 1).




Fig. 1. Distribution of Arsenic (As), Lead (Pb) and Chromium (Cr) in different organs of *H. nehereus*. Bars with different letters (a and b) differ significantly ($P<0.05$)

The mean values of Chromium were the least among the three investigated metals (0.03 ppm, 0.002 ppm, 0.0005 ppm and 0.00025 ppm for gill, liver, kidney, and muscle respectively) which are far lower than the standard value (0.05-0.15 ppm) of Chromium stated by WHO and FEPA (Federal Environmental Protection Agency) [18,19]. There were however no significant difference in the chromium concentrations of different organs. The mean values of Pb for gills (0.083 ppm), liver (0.029 ppm), kidney (0.083 ppm) and muscle (0.043 ppm) were observed (Figure 1). All the values of lead were lower than the standard value (0.3 ppm) issued by European Commission [20]. From Figure 1, the accumulation pattern of metals in different organs of *H. nehereus* were: As>Pb>Cr for liver and Pb>As>Cr for kidney, gill and muscle. Concentrations of metals in muscle tissue of fish species collected from Northeast coast of India were determined [12]. They found higher concentration of Arsenic accumulation in *Harpadon nehereus* which was in agreement with this study.

3.2 Arsenic (As), Lead (Pb) and Chromium (Cr) Concentration in *H. ilisha*

The recorded values in *H.ilisha* revealed that muscle had the lowest average arsenic 0.029 ppm while kidney had the highest average arsenic 0.070 ppm which are higher than the recommended value of 0.01 ppm [17] (Figure 2).




Fig. 2. Distribution of Arsenic (As), Lead (Pb) and Chromium (Cr) in different organs of *H. ilisha*. Bars with different letters (a and b) differ significantly ($P<0.05$)

There was however no significant difference in arsenic contents of gill, liver and muscle except kidney. The observed values of Chromium in different organs were the least among the three investigated metals which are far lower than the standard value of Chromium [18,19]. The mean values of lead (0.038 ppm, 0.025 ppm, 0.076 ppm, 0.034 ppm for gill, liver, kidney and muscle respectively) were observed which are also lower than the standard value of 0.3 ppm and showed a significant ($P<.05$) variation between lead contents of liver and kidney. The accumulation patterns in the gill and liver of the *H. ilisha* revealed the following order, $As>Pb>Cr$. In the kidney and muscle, the concentrations revealed $Pb>As>Cr$ sequence (Figure 2).

3.3 Arsenic (As), Lead (Pb) and Chromium (Cr) Concentration in *Pampus chinensis*

The observed values in *P. chinensis* demonstrated that the mean values of arsenic obtained from kidney and gill (0.067 ppm and 0.029 ppm respectively) were significantly ($P<.05$) different from the values of liver and muscle (0.036 ppm and 0.032 ppm) (Figure 3) which are higher than the recommended value 0.01 ppm.




Fig. 3. Distribution of Arsenic (As), Lead (Pb) and Chromium (Cr) in different organs of *P. chinensis*. Bars with different letters (a and b) differ significantly ($P < 0.05$)

The mean values of Chromium in different organs were the least among the three investigated metals which are far lower than the standard value. There was no significant difference in the chromium concentrations of different organs. The average values of lead were also observed (0.067 ppm, 0.008 ppm, 0.083 ppm and 0.021 ppm for gill, liver, kidney and muscle respectively) which are also lower than the standard value. The accumulation levels of metals in organs of *P. chinensis* were: $Pb > As > Cr$ for gill and kidney and $As > Pb > Cr$ for liver and muscle (Figure 3).

In this study, arsenic and lead had the highest concentrations in all tissues of the investigated species while chromium was the element that had the lowest detected concentration.

3.4 Organ and species wise Arsenic (As), Lead (Pb) and Chromium (Cr) concentration

The data in Table 1 shows that there were significant differences in the metal amounts except Cr among different organs. Metal concentrations varied among different organs of a species as well as between species. Many researchers have reported that heavy metals accumulate in high concentrations in the liver, the gill and the kidney, because these organs have relatively higher potential for metal accumulation than muscle [21, 22].

Table 1. The comparison of different organs of the species with respect to the Arsenic (As), Lead (Pb) and Chromium (Cr) accumulation levels

Heavy metal	<i>H. nehereus</i>	<i>P. chinensis</i>	<i>H. ilisha</i>
Ar	$K^1 > L^1 > G^2 > M^2$	$K^1 > L^1 > M^2 > G^2$	$K^1 > L^1 > G^2 > M^2$
Pb	$K^1 = G^{1-2} > M^{1-2} > L^2$	$K^1 > G^{1-2} > M^{1-2} > L^2$	$K^1 > G^1 > M^2 > L^2$
Cr	$G^1 > L^1 > K^1 > M^1$	$G^1 > K^1 > L^1 = M^1$	$G^1 > K^1 > M^1 = L^1$

*The difference in the metal accumulation levels of the organs with a different number in the same row relating to any species is significant ($P < .05$).

K=kidney; G=gill; L=liver; M=muscle.

When we consider the organs of all three fishes, highest concentration of arsenic was estimated in kidney and second-best value was recorded in liver (Table 1). Arsenic concentrations in liver and kidney found statistically different from gill and muscle. This might be related to the feeding habit of fish species which feeds on decayed organic matter. Fishes normally endure to high pollutant exposure as they can develop pollutant-sequestering detoxifying systems which make the metal inert and not toxic to the organism [23].

In case of all three fishes, highest lead concentration was estimated in kidney and the second-best value was recorded in gill. The recorded values in Liver and Muscle were significantly different from the kidney and gill (Table 1). It was found that lead concentration more than the maximum permissible limits in gill (1.55 ppm) and liver (0.85 ppm) tissues of fish, *Harpodonnehereus* in the west coast of India during monsoon period [24]. The findings of this study are contrary to [25] who reported that lead was greater in liver as compared to the gills and kidney.

For all three marine fishes, chromium concentration was found highest in gill with very lower value in liver, kidneys and muscles which were not statistically different (Table 1). An MSDS (Material Safety Data Sheet) report showed that chromium accumulates primarily in the gill, liver, kidney, and bone in different fish species [26]. This study observation agrees with those of another study [27]. For instance, the muscle Pb contents of three investigated fish species were remarkably higher than a metabolite organ, liver (Table 1). As indicated in the literature, this finding may be the result of specific bioaccumulation characteristics, special to the species and the elements [28,29].

So, Table 1 shows that in terms of organ the metals were mostly concentrated in the kidney tissues. But the concentration in muscle was found to be the lowest which is a positive finding as we consume the muscles widely even though the concentration of arsenic in muscles was recorded higher than the safety values.

It was observed that *H. nehereus* showed the highest metal accumulation value compared to the other two species (Table 2). As *H. nehereus* occupies a higher trophic level of the food chain, the bioaccumulation process may increase the concentration of metals in their bodies.

Table 2. The comparison of the fish species with respect to the Arsenic (As), Lead (Pb) and Chromium (Cr) accumulation levels of the organs

Heavy metal	Organ	Fish species
Ar	Gill	<i>H. nehereus</i> ¹ > <i>H. ilisha</i> ¹⁻² > <i>P. chinensis</i> ²
	Liver	<i>H. nehereus</i> ¹ > <i>H. ilisha</i> ¹⁻² > <i>P. chinensis</i> ²
	Kidney	<i>H. nehereus</i> ¹ > <i>H. ilisha</i> ¹ > <i>P. chinensis</i> ¹
	Muscle	<i>H. nehereus</i> ¹⁻² > <i>P. chinensis</i> ¹ > <i>H. ilisha</i> ²
Pb	Gill	<i>H. nehereus</i> ¹⁻² > <i>P. chinensis</i> ¹ > <i>H. ilisha</i> ²
	Liver	<i>H. nehereus</i> ¹ > <i>H. ilisha</i> ¹ > <i>P. chinensis</i> ¹
	Kidney	<i>H. nehereus</i> ¹ > <i>P. chinensis</i> ¹ > <i>H. ilisha</i> ¹
	Muscle	<i>H. nehereus</i> ¹ > <i>H. ilisha</i> ¹ > <i>P. chinensis</i> ¹
Cr	Gill	<i>H. nehereus</i> ¹ > <i>P. chinensis</i> ¹ > <i>H. ilisha</i> ¹
	Liver	<i>H. nehereus</i> ¹ > <i>P. chinensis</i> ¹ = <i>H. ilisha</i> ¹
	Kidney	<i>H. nehereus</i> ¹ > <i>P. chinensis</i> ¹ = <i>H. ilisha</i> ¹
	Muscle	<i>H. nehereus</i> ¹ > <i>P. chinensis</i> ¹ = <i>H. ilisha</i> ¹

*The difference in the metal accumulation levels of the marine fish species with a different number in the same row is significant ($P<.05$).

The bio-accumulation mechanism involves uptake of the metal with food taken by marine organisms. The food is digested, assimilated, and excreted but the metal accumulates in the organs of the fishes [30]. The metal concentrations of different marine fish species have been widely studied [31,9,32,33,24]. There are similarities, as well as differences, between the results from our study and the findings from another research. It has been indicated that the levels of metals in fish depend on habitats, the durations of exposure of the fish to contaminants, their feeding habits and the age and the size of the species [34,35].

Therefore, we can conclude that the studied value of arsenic is higher than the recommended values of human intake in all the three cases of fish, indicating the low water quality of Chattogram coastal area. The Chattogram coastal water might be highly polluted with arsenic. It could be associated to effluent discharges from a major industrial and siderurgic plant near the sea and to agro-industrial activities, mainly run off from agricultural soils where fertilizers are used and other waste [10].

4. CONCLUSION

As being a widely exposed coast of the Bay of Bengal, Chattogram coast is affected by numerous numbers of metals. As this region is the principal harvesting areas of marine fishes, this metal toxicity can cause indeed a supreme problem in human health. From the findings of this study, it was concluded that the fish kidney exhibited highest accumulation of metals (As and Pb) in *H. nehereus*, *P. chinensis* and *H. ilisha*. Fish gill showed highest accumulation of Cr in three investigated fish species which is lower than the standard value of Cr. The current research has revealed that arsenic concentration is far above from the

safety levels while the concentration of lead is also approaching the acceptable limit. The immediate safety and strict regulatory actions on environment friendly discharge metal is mandatory which should be ensured by the government organizations as well as by the non-government responsible personnel and organizations.

REFERENCES

1. Gjedrem T, Robinson N, Rye M. The importance of selective breeding in aquaculture to meet future demands for animal protein: A review. *Aquac.* 2012;350-353(3): 117-129. DOI:[10.1016/j.aquaculture.2012.04.008](https://doi.org/10.1016/j.aquaculture.2012.04.008).
2. Kris-Etherton PM, Harris WS, Appel LJ. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circ.* 2002;106(21):2747–2757. <https://doi.org/10.1161/01.CIR.0000038493.65177.94>.
3. Mazid MA. Manual on culture of small and threatened indigenous fish species. Bangladesh Fisheries Research Institute, Department of Fisheries, Bangladesh Agricultural University and Ministry of Fisheries and Livestock. 2005. Accessed 25 June 2021.
4. DoF. Department of Fisheries, Yearbook of Fisheries Statistics of Bangladesh, 2019-20. Fisheries Resources Survey System (FRSS), Department of Fisheries. Bangladesh: Ministry of Fisheries and Livestock. 2020; Volume 37: 141p. Website: <http://www.fisheries.gov.bd>.
5. Annabi A, Said K, Messaoudi I. Cadmium: bioaccumulation, histopathology and detoxifying mechanisms in fish. *Am J Res Comun.* 2013;1(4):60–79
6. Gungum B, Gulsun Z, Unlu E. Heavy metal pollution in water, sediment and fish from the Tigris River in Turkey. *Chemosphere.* 1994;29(1):111-116. DOI:[10.1016/0045-6535\(4\)90094-9](https://doi.org/10.1016/0045-6535(4)90094-9)
7. Idris AM, Alqahtani F, Said TO, Fawy KF. Contamination level and risk assessment of heavy metal deposited in street dusts in Khamees-Mushait city, Saudi Arabia. *Hum Ecol Risk Assess: An International Journal.* 2018;26(2) 495-511. DOI:[10.1080/10807039.2018.1520596](https://doi.org/10.1080/10807039.2018.1520596)
8. Sioen I, Henauw SD, Verdonck FAM, Thuyne NV, Camp VJ. Development of a nutrient database and distributions for use in a probabilistic risk-benefit analysis of human seafood consumption. *J Food Compos Anal.* 2007;20(8):662–670. DOI:[10.1016/j.jfca.2006.11.001](https://doi.org/10.1016/j.jfca.2006.11.001).
9. Sujatha K, Durairajsinghu V, Silambarasan K. Determination of some heavy metals in fish, water and sediments from bay of bengal. *Pollut res.* 2016;35(4):797-799. <https://www.researchgate.net/publication/317202279>
10. Jothi JS, Anka IJ, Hashem S, Morshed S. Assessment of heavy metal concentration in edible fish muscle and water sample collected from different location in Chittagong: a public health concern. *Ukr Food J.* 2018;Volume 7. Issue 3. DOI:[10.24263/2304-974X-2018-7-3-11](https://doi.org/10.24263/2304-974X-2018-7-3-11)
11. Mukherjee DP, Kumar B. Assessment of arsenic, cadmium and mercury level in commonly consumed coastal fishes from Bay of Bengal, India. *Food Sci Qual Manag.* 2011;ISSN 2224-6088 (Paper) ISSN 2225-0557 (Online), Vol 2. www.iiste.org;
12. Kumar B, Sajwan KS, Mukherjee DP. Distribution of Heavy Metals in Valuable Coastal Fishes from North East Coast of India. *Turkish J Fish Aquat Sci.* 2012;12: 81-88. DOI:[10.4194/1303-2712-v12_1_10](https://doi.org/10.4194/1303-2712-v12_1_10)
13. Monwar M. Flow and discharge characteristics in the Karnaphuli River Estuary. Inter-term Research Paper. Institute of Marine Sciences, Chittagong University, Bangladesh. 2001.
14. Khan MU, Muhammad S, Manik RN, Khan AK, Tariq M. Heavy metals potential health risk assessment through consumption of wastewater irrigated wild plants: A

- case study. Hum Ecol Risk Assess. 2015;volume 22, issue 1.DOI:[10.1080/10807039.2015.1056292](https://doi.org/10.1080/10807039.2015.1056292)
15. Rashid H, Hasan MN, Tanu MB, Parveen R, Sukhan ZP, Rahman MS, Mahmud Y. Heavy Metal Pollution and Chemical Profile of KhiruRiver, Bangladesh. Int J Environ. 2015;2(1):57-63. Accessed 24 March 2012. Available: <http://www.BENJapan.org/IJE>.
 16. UNEP. Sampling of selected marine organisms and sample preparation for trace metal analysis: Reference Method for Marine Pollution Studies. 1984;No. 7, Rev. 2: 19 p.
 17. WHO (World Health Organization). Guidelines for drinking-water quality, 3rd. Geneva: World Health Organization, WHO Food Standards Programme, Codex Committee, Rotterdam. Reference CX/FAC 02/16. 2004.
 18. WHO (World Health Organization) vol. 1. Recommendation WHO; Geneva: Guidelines for Drinking Water Quality. 1985;p. 130.
 19. FEPA. Federal Environmental Protection Agency. Guidelines and Standards for Environmental Pollution Control in Nigeria. 2003;p. 238.
 20. European Commission, Commission Regulation (EC) No. 629/2008, Amending Regulation (EC) No. 1881/2006 setting maximum levels for certain contaminants in foodstuffs. Official Journal of the European Union. 2008; L173:6–9.
 21. Bashir F, Alhemmal E. Analysis of some heavy metal in marine fish in muscle, liver and gill tissue in two marine fish species from Kapar coastal waters, Malaysia. The Second Symposium on Theories and Applications of Basic and Biosciences. 2015;volume 2. <http://www.researchgate.net/publication/301554192>.
 22. Vijayasundaram V, Nagar A. The Bioaccumulation of Arsenic and the Efficacy of Meso2, 3-dimercaptosuccinic Acid in the Selected Organ Tissues of Labeorohita Fingerlings Using Inductively Coupled Plasma-Optical Emission Spectrometry. World Appl Sci J. 2009;6(9):1247-1254. <http://www.researchgate.net/publication/254606465>
 23. Weis JS, Weis P. Effects of environmental pollutants on early fish development. Reviews in Aquatic Sciences. 1989;1:45-73
 24. Oza A, Muralidharan L. Seasonal study on bioaccumulation of heavy metal, lead in various tissues of fish, harpodonnehereus collected from sassoon dock, mumbai coast of maharashtra, India. Int J Adv Res Sci Eng. 2018;7(4):265-272.<https://www.researchgate.net/publication/333688552>.
 25. Farombi EO, Adelowo OA, Ajimoko YR. Biomarkers of Oxidative Stress and Heavy Metal Levels as Indicators of Environmental Pollution in African Cat Fish (*Clariasgariepinus*) from Nigeria Ogun River. Int J Environ Res Public Health. 2007;4(2):158-65.DOI: [10.3390/ijerph2007040011](https://doi.org/10.3390/ijerph2007040011).
 26. MSDS. The MSDS hyperglossary heavy metal. 2006. Available online: http://www.ilpi.com/msds/ref/heavy_metal.html.
 27. Amundsen PA, Staldvik FJ, Lukin AA, Kashulin NA, Popova OA, Reshetnikov YS. Heavy metal contamination in freshwater fish from the border region between Norway and Russia. Sci Total Environ. 1997;201(3):211-224. DOI: 10.1016/s0048-9697(97)84058-2
 28. Henry F, Amara R, Courcot L, Lacouture D, Bertho ML. Heavy metals in four fish species from the French coast of the Eastern English Channel and Southern Bight of the North Sea. Environ Int. 2009;30(5):675–83. DOI: 10.1016/j.envint.2003.12.007
 29. Papagiannis I, Kagalou I, Leonardos J, Petridis D, Kalfakakou V. Copper and zinc in four freshwater fish species from Lake Pamvotis (Greece). Environ Int. 2004;30(3):357–362. <https://doi.org/10.1016/j.envint.2003.08.002>.
 30. Goel PK. Water pollution: causes, effects and control, New Age International (P) Limited, New Delhi, India. 1996;pp 97-115.
 31. Chakraborty S, Rudra T, Guha A, Ray A, Pal N, Mitra A. Spatial variation of heavy metals in Tenualosailisha muscle: A case study from the lower Gangetic delta and coastal West Bengal. Int J Inno Sci, Eng Tech. 2016;Vol. 3 Issue 4.

32. Younis A, Amin H, Alkaladi A, Mosleh Y. 2015. Bioaccumulation of heavy metals in fish, squids and crustaceans from the Red Sea, Jeddah Coast, Saudi Arabia. *Open Mar Sci.* 2015;5(4):369-378. DOI:10.4236/ojms.2015.54030.
33. Salam MA, Paul SC, Noor SNBM, Siddiqua SA, Aka TD, Wahab R and Aweng ER. Contamination profile of heavy metals in marine fish and shellfish. *Glob J Environ Sci.* 2019;5(2): 225-236. DOI:10.22034/gjesm.2019.02.08.
34. Canli M, Atli G. The relationships between heavy metal (Cd, Cr, Cu, Fe, Pb, Zn) levels and the size of six Mediterranean fish species. *Environ Pollut.* 2003;121(1):129–136. [https://doi.org/10.1016/S0269-7491\(02\)00194-X](https://doi.org/10.1016/S0269-7491(02)00194-X).
35. Rashed MN. Monitoring of environmental heavy metals in fish from Nasser Lake. *Environ Int.* 2001;27(1):27–33. DOI: [10.1016/s0160-4120\(01\)00050-2](https://doi.org/10.1016/s0160-4120(01)00050-2).

UNDER PEER REVIEW