

# **The Effect of Ramadan Fasting on Diabetic Patients With High Risk of Cardiovascular Disease**

## **Abstract:**

One of the common questions for patients with a history of cardiovascular disease is whether it is safe for them to fast during Ramadan. Yet, studies on the effects of Ramadan fasting on blood lipids, blood pressure, anthropometric parameters and other cardiovascular risk factors are scarce, and have given inconclusive results. The aim of this study is to investigate the effect of Ramadan fasting on cardiovascular risk factors, including biochemical indices, blood pressure and main anthropometric parameters. This prospective observational study was conducted at the CCU and medical wards at Madinat Zayed hospital, one of the Al Dhafra hospitals, under Seha Abu Dhabi. Fifty-six diabetic patients admitted with cardiovascular events, including 48 males and 8 females, with a previous history of cardiovascular events completed the study. Their ages ranged between 32 and 91 years with a mean of  $54.0 \pm 10$  years. A non-significant slight biochemical modification with regards to the metabolic profile pre, during and post-Ramadan period was reported: eGFR, total cholesterol, LDL, and HDL were increased, yet triglycerides and creatinine were non-significantly lower during Ramadan. The findings also showed a non-significant decrease in the post Ramadan blood pressure values that could be attributed to the metabolic switch.

**Keyword:** Ramadan; Fasting; Diabetes; Cardiovascular

## **Introduction:**

Ramadan is a month during which Muslims abstain from eating, drinking and smoking from dawn to sunset. During month of Ramadan, there are changes in the quality of food and eating patterns. It might be due to the consumption of more carbohydrates and sweet foods, mainly in the form of two large meals at dawn and sunset. It has been established that a given nutrient ingested at an unusual time can induce different metabolic effects. However, the physiological changes during Ramadan are not well known.

Fasting during Ramadan is essentially a radical change in lifestyle for the period of one lunar month that may affect cardiovascular risk in patients with coronary artery disease (CAD) and cerebro-vascular disease (CVD). The most common risk factors associated with increased risk of atherosclerotic heart disease or stroke are abnormalities in plasma lipids and some coagulation and hemostatic factors, hypertension and smoking. Cardiovascular diseases are the leading cause of death worldwide.

Lipid profiles are affected by factors such as changes in dietary habits, consuming different dietary fats, increased consumption of refined sugar, and reduced physical activity. Many studies reported a significant reduction in type 2 diabetic patients' weight during Ramadan. Others reported a non-significant reduction in weight of these patients.

Bouguerra et al. showed Ramadan fasting has a moderate effect on glycaemia and lipoprotein levels in type 2 diabetic patients when previous metabolic control was quite good, but fasting induced more deterioration when previous control was poor. Although another study in Iran showed a deterioration of glycemic control after Ramadan fasting in type 2 diabetic patients, it was more evident in patients using oral hypoglycemic medication than diet-controlled patients.

Fasting is not meant to create excessive hardship for individuals who are not able to do it, the Islamic practice exempts many people from fasting, including those who are ill and for whom fasting would be harmful. One of the common questions for patients with a history of cardiovascular disease is whether it is safe for them to fast during Ramadan. Yet, studies on the effects of Ramadan fasting on blood lipids, blood pressure, anthropometric parameters and other cardiovascular risk factors are scarce, and have given inconclusive results. In current study, the effect of Ramadan fasting on cardiovascular risk factors, including biochemical indices, blood pressure and main anthropometric parameters are investigated.

## **Methodology:**

This prospective observational study was conducted at the CCU and medical wards of Madinat Zayed hospital, one of the Al Dhafra hospitals, under Seha Abu Dhabi.

The primary outcome measure was to observe any Major Adverse Cardiovascular Events (MACE), including a combination of all-cause mortality, and myocardial infarction during the time period studied, which was assessed as one month before Ramadan, during Ramadan and one month after Ramadan, corresponding the month of Ramadan to the Gregorian calendar, as the lunar calendar is 11–12 days shorter than the solar year.

Any diabetic Muslim patient admitted to the hospital during this time between the ages of 18-80 with cardiovascular events, baseline clinical characteristics including age, gender, cardiovascular risk factors, and primary as well as secondary cardiovascular diagnoses was included in the study.

### **Data Collection**

Qualified nurses who can speak and write both English and Arabic were recruited to administer the questionnaires and perform anthropometric measurements.

A standardized questionnaire-based face-to-face interview was conducted by the nurses to fill out the questionnaires. The questionnaire was composed of:

- (a) Sociodemographic data such as age, sex, nationality, marital status, education level, occupation, height, weight, and parental consanguinity
- (b) Anthropometric data such as height and weight
- (c) Lifestyle habits such as physical activity and smoking status
- (d) Blood pressure measurements, and
- (e) Laboratory investigations, such as blood glucose, glycated hemoglobin (HbA1c), low-density and high-density lipoprotein (HDL and LDL) cholesterol, cholesterol levels, triglyceride, urea, creatinine, bilirubin, albumin creatinine ratio, etc. Data related to anthropometry and laboratory tests was filled based on actual measurements.

Anthropometric measurements like height was measured in centimeters using a height scale while the patient was standing barefoot with normal, straight posture, whereas weight was measured in kilograms using a weight scale

BMI was calculated as the ratio of weight (kg) to the square of height (m). A person was considered obese if the BMI value was at least 30kg/m<sup>2</sup> and overweight if BMI was greater than 25kg/m<sup>2</sup> and less than 30kg/ m<sup>2</sup>.

### **Blood pressure measurements**

Hypertension was defined according to the WHO, which is systolic blood pressure at least 130mmHg or diastolic blood pressure at least 85mm Hg or using antihypertensive medication. Two readings of systolic blood pressure and diastolic blood pressure were taken from the participant's left arm while seated and his/her arm at heart level, using a standard zero mercury sphygmomanometer after at least 10–15 minutes of rest. Thereafter, the average of the two readings was obtained.

### **Lifestyle habits**

Smoking habit was classified in terms of past or current smoker or nonsmoker.

Patients were classified as physically active if they reported participating in walking or cycling for more than 30 minutes per day.

All lab investigations in this study were performed in the laboratory department in Madinat Zayed Hospital – Al Dhafra region in Abu Dhabi state:

- The blood and urine specimens were collected either in phlebotomy rooms in the laboratory department or in the CCU and medical wards of MZH then sent directly to the core lab, which operates 24 hours a day.
- Blood samples were collected through venipuncture from each participant following the standard procedures of preparation and collected in BD containers (EDTA tubes, SST tubes and Li Heparin tubes).

All parameters below were measured using Auto-analyzer (cobas® 6000 analyzer-c502 and e602 modules - Roche - USA):

- Fasting blood glucose was measured by hexokinase method
- Glycosylated hemoglobin (HbA1c) levels were analyzed based on TINIA Gen.2 method with automatic calculation to report HbA1c % using formula:  $(A1/HB)*91.5+2.15$
- Fasting Lipid Panel: (Total cholesterol, Triglyceride and HDL-c) using enzymatic colorimetric tests.

- The LDL-c was calculated using Friedewald equation:  $LDL-c = Total\ cholesterol - (HDL\ cholesterol + Triglyceride * 0.46)$
- Renal function profile: U+E includes urea, creatinine, electrolytes (sodium, potassium, chloride and bicarbonate levels). The electrolytes were analyzed using ISE method.
- Urine albumin/creatinine ratio was determined by mg/mmol considering normal values  $\leq 3.00$  ; eGFR using CKD-EPI auto-calculation
- Troponin levels were performed using Trop. T high sensitive STAT kits, and for proBNP II (which is diagnostic for heart failure), the sandwich principle immunoassay was used.
- Total and direct blood bilirubin using Colorimetric diazo method.


The presence of Diabetes Mellitus was determined by the documentation in the patient's previous or current medical record of a documented diagnosis of DM that had been treated with diabetes medications.

## Results

Fifty-six volunteers, including 48 males and 8 females, with a previous history of cardiovascular event completed the study. Their age ranged between 32 and 91 years with a mean of  $54.0 \pm 10$  years. The metabolic profile of the study volunteers is shown in Table 1 and Image 1 to visualize it. The age of the participants was similar in the three groups with no statistical significance ( $p$ -value=0.38). In addition, no significant difference was seen in metabolic profile (duration of diabetes, Glycated Hemoglobin, ACR, eGFR, creatinine and blood pressure (systolic/diastolic)) means recorded at different periods. eGFR level was the highest during Ramadan compared to other periods ( $86.16 \pm 31.56$ ).

Parameters	Before Ramadan	During Ramadan	After Ramadan	P value (ANOVA)
AGE	58±14.02	53.88±11.32	52.40±13.58	0.38
DURATION OF DIABETES	4.45 ±2.30	2.64 ±1.5	6.50 ±2.5	0.91
HbA1c	8.57±2.32	8.04±2.48	7.89±1.90	0.62
eGFR	56.22±34.06	86.16±31.56	73.20±46.01	0.06
CHOLESTEROL	4.56±1.11	4.62±1.17	4.70±1.46	0.94
LDL	2.57±0.83	2.77±0.94	2.56±1.09	0.79
TG	3.06±4.13	1.96±1.09	2.61±1.85	0.54
HDL	0.95±0.19	0.99±0.21	1.02±0.23	0.69
BMI	26.57±6.93	27.44±8.26	28.49±5.79	0.67
BP(Systolic )	130.26±11.61	125.8±17.99	124.40±14.87	0.86

BP(Diastolic)	75.15±14.17	77.17±11.36	75±7.85	0.92
CREATININE	131±116.20	94.05±39.81	105.22±76.97	0.40



**Image 1:** Means of Metabolic Profile of the study subjects

### Lipid profile

The values of total cholesterol, LDL and HDL were increased but it wasn't statistically significant, but the levels for triglycerides were non-significant decreased during the time intervals studied. The calendar chart for lipid profile is presented in **Figure 2**. March and May were the highest levels of lipids with a mean of 3.12 for HDL, 4.19 for cholesterol and 2.3 for LDL (Figure 3).

**Figure 1** Calendar plot showing metabolic parameters during Ramadan




Figure 2. Calendar Plot for Lipid Profile



Figure 3: Mean and distribution of Lipid Profile (2021-2022)

## Discussion

This study evaluated the effect of holy Ramadan fasting on the metabolic health of patients with cardiovascular risk factors.

The findings revealed slight biochemical modifications with regards to the metabolic profile of the patients, but the changes were not significant during any given time interval. We showed non-significant changes in the glycated hemoglobin pre, during and post-Ramadan period. This is in accordance with the previous studies.

We also found a non-significant decrease in the post Ramadan blood pressure values, which could be attributed to the metabolic switch (in which the body switches to use ketones for energy instead of glucose after 8–12 hours of fasting) described in a recent review on intermittent fasting. Insulin drop during the metabolic switch is suggested as the key reason behind reduced blood pressure after fasting. Blood pressure is inversely correlated with parasympathetic nervous system activity and correlated with sympathetic nervous system activity. When sympathetic nervous system activity is triggered by insulin, adrenal glands produce norepinephrine, which binds to  $\alpha$ -receptors in the blood vessels. Consequently, vasoconstriction occurs. Also, insulin acts downstream and enhances renal tubular sodium reabsorption, which confers more water retained in circulation. As a result, blood pressure is raised. Other human and animal studies associate this reduction with the release of brain-derived neurotrophic factor, which elevates parasympathetic nervous system activity and improves insulin sensitivity.

As seen in this study, LDL-cholesterol was lower in the serum after Ramadan compared to levels before Ramadan. Also, HDL-cholesterol levels were higher in the serum at the end of Ramadan compared to levels before Ramadan. These results were in agreement with those reported by other authors on healthy persons. The previous data concerning HDL-cholesterol clearly explain the beneficial effect of Ramadan fasting on serum lipids of diabetics.

Serum total cholesterol and triglyceride levels decreased non significantly ( $P > 0.05$ ) towards the end of the study. The increase in the glucose levels during the fasting undoubtedly reduced the catabolism and the significant mobilization of the triglycerides from peripheral tissues to plasmatic circulation. These results may also be explained by different food habits of the populations studied and the type of food consumed during Ramadan, as there is a tendency towards an increased intake of carbohydrates and fat.

At kidney level, we noted a discrete transient increase of creatinine during Ramadan compared with the basic state. At the same time, there was a transient and significant decrease of clearance

in both the overall population and diabetic patients. The same findings were noted in a Tunisian study performed in 2013 on athletes whose decrease in kidney function during Ramadan was attributed to a hydration defect that is frequent in the summer period.

This study, however, presents some limitations such as the small sample size and the fact that the participants were recruited from a single center, which affects the generalizability of the results.

### **Conclusion**

Finally, non-significant slight biochemical modification with regards to the metabolic profile pre, during and post-Ramadan period was reported: eGFR, total cholesterol, LDL, and HDL were increased, yet triglycerides and creatinine were non-significantly lower during Ramadan. Findings also showed a non-significant decrease in the post-Ramadan blood pressure values, which could be attributed to the metabolic switch.

## References

1. Khafaji HA, Bener A, Osman M, Al Merri A, Al Suwaidi J. The impact of diurnal fasting during Ramadan on the lipid profile, hs-CRP, and serum leptin in stable cardiac patients. *Vasc Health Risk Manag.* 2012;8:7–14. doi:10.2147/VHRM.S22894
2. Kiyani MM, Memon AR, Amjad MI, Ameer MR, Sadiq M, Mahmood T. Study of human biochemical parameters during and after Ramadan. *J Relig Health.* 2017;56(1):55–62.
3. Nematy M, Alinezhad-Namaghi M, Rashed MM, et al. Effects of Ramadan fasting on cardiovascular risk factors: a prospective observational study. *Nutr J.* 2012;11:69. doi:10.1186/1475-2891-11-69
4. Tong CV, Yow HY, Noor NM, Hussein Z. Diabetes emergencies around Ramadan study (DEARS)—A multi-center study of diabetes emergencies admitted before, during and after Ramadan in Malaysia. *Diabetes Research and Clinical Practice.* 2021 May 1;175:108854.
5. Beshyah SA, Ali KF, Hafidh K, Hajjaji IM. Ramadan fasting and diabetes 2019: The year in review. *Diabetes Research and Clinical Practice.* 2021; 1;172:108593.
6. Abdelrahim D, Faris ME, Hassanein M, Shakir AZ, Yusuf AM, Almeneessier AS, BaHamman AS. Impact of Ramadan diurnal intermittent fasting on hypoglycemic events in patients with type 2 diabetes: a systematic review of randomized controlled trials and observational studies. *Frontiers in Endocrinology.* 2021; 8;12:624423.
7. Shehab A, Abdulle A, El Issa A, Al Suwaidi J, Nagelkerke N. Favorable changes in lipid profile: the effects of fasting after Ramadan. *PLoS One.* 2012;7:e47615. doi:10.1371/journal.pone.0047615
8. Azizi F. Islamic fasting and health. *Ann Nutr Metab.* 2010;56:273–282. doi:10.1159/00029584
9. Lessan N, Hannoun Z, Hasan H, Barakat MT. Glucose excursions and glycaemic control during Ramadan fasting in diabetic patients: insights from continuous glucose monitoring (CGM). *Diabetes Metab.* 2015;41:28–36. doi:10.1016/j.diabet.2014.11.004
10. Salti I, Benard E, Detournay B, et al. A population-based study of diabetes and its characteristics during the fasting month of Ramadan in 13 countries: results of the epidemiology of diabetes and Ramadan 1422/2001 (EPIDIAR) study. *Diabetes Care.* 2004;27:2306–2311. doi:10.2337/diacare.27.10.2306
11. D'Agostino RB, Vasan RS, Pencina MJ, et al. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation.* 2008;117:743–753. doi:10.1161/CIRCULATIONAHA.107.699579

12. Onyenekwu CP, Hoffmann M, Smit F, Matsha TE, Erasmus RT. Comparison of LDL-cholesterol estimate using the Friedewald formula and the newly proposed de Cordova formula with a directly measured LDL-cholesterol in a healthy South African population. *Ann Clin Biochem.* 2014;51:672–679. doi:10.1177/0004563214520750
13. Nair S, Hardy KJ, Wilding JP. The Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) formula performs worse than the Modification of Diet in Renal Disease (MDRD) equation in estimating glomerular filtration rate in type 2 diabetic chronic kidney disease. *Diabet Med.* 2011;28:1279. doi:10.1111/j.1464-5491.2011.03281.x
14. Mathews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia.* 1985;28:412–419.
15. Thompson FE, Subar AF. Dietary assessment methodology. In: Coulston AM, Boushey CJ, editors. *Nutrition in the Prevention and Treatment of Disease.* 2nd ed. Philadelphia, PA: Academic Press; 2008:3–39.
16. Salim I, Al Suwaidi J, Ghadban W, Alkilani H, Salam AM. Impact of religious Ramadan fasting on cardiovascular disease: a systematic review of the literature. *Curr Med Res Opin.* 2013;29:343–354. doi:10.1185/03007995.2013.840568
17. Bahijri S, Borai A, Ajabnoor G, et al. Correction: relative metabolic stability, but disrupted circadian cortisol secretion during the fasting month of Ramadan. *PLoS One.* 2013. doi:10.1371/journal.pone.0060917
18. Benaji B, Mounib N, Roky R, et al. Diabetes and Ramadan: review of the literature. *Diabetes Res Clin Pract.* 2006;73:117–125. doi:10.1016/j.diabres.2005.10.028
19. Celik A, Saricicek E, Saricicek V, et al. Effect of Ramadan fasting on serum concentration of apelin-13 and new obesity indices in healthy adult men. *Med Sci Monit.* 2014;20:337–342. doi:10.12659/MSM.890139
20. Kul S, Savaş E, Oztürk ZA, Karadağ G. Does Ramadan fasting alter body weight and blood lipids and fasting blood glucose in a healthy population? A meta-analysis. *J Relig Health.* 2014;53:929–942. doi:10.1007/s10943-013-9687-0
21. Trepanowski JF, Bloomer RJ. The impact of religious fasting on human health. *Nutr J.* 2010;9:57. doi:10.1186/1475-2891-9-57
22. Kareem I, Asegaonkar S, Bavikar J, Pagdhune A, Aghade S, Thorat A. Effect of Ramadan fasting on metabolic profile in healthy adults of Aurangabad, India. *Ijsr.* 2013;2:451–452. doi:10.15373/22778179/NOV2013/149

23. Gnanou JV, Caszo BA, Khalil KM, Abdullah SL, Knight VF, Bidin MZ. Effects of Ramadan fasting on glucose homeostasis and adiponectin levels in healthy adult males. *J Diabetes Metab Disord.* 2015;14:55. doi:10.1186/s40200-015-0183-9
24. Aziz KM. Effect of fasting Ramadan in diabetes control status – application of extensive diabetes education, serum creatinine with HbA1c statistical ANOVA and regression models to prevent hypoglycemia. *Recent Pat EndocrMetab Immune Drug Discov.* 2013;7:233–251. doi:10.2174/18715303113139990010
25. Karatoprak C, Yolbas S, Cakirca M, et al. The effect of long term fasting in Ramadan on glucose regulation in type 2 diabetes mellitus. *Eur Rev Med Pharmacol Sci.* 2013;17:2512–2516.
26. Ouhdouch F, Adarmouch L, Errajaji A, Amine M, El Ansari N. Absence d'effets délétères du jeûne du Ramadan sur l'équilibre glycémique chez des patients diabétiques: rôle des consultations de préparation au jeûne. *Médecine Des Maladies Métaboliques.* 2011;5:448–452. doi:10.1016/S1957-2557(11)70289-9
27. Maislos M, Abou-Rabiah Y, Zuili I. Improved diabetes control after prolonged fasting: the Ramadan model. *Practical Diabetes Int.* 2001;18:149–151. doi:10.1002/pdi.166
28. Al-Shafei AI. Ramadan fasting ameliorates oxidative stress and improves glycemic control and lipid profile in diabetic patients. *Eur J Nutr.* 2014;53:1475–1481. doi:10.1007/s00394-014-0650-y
29. Ibrahim WH, Habib HM, Jarrar AH, Al Baz SA. Effect of Ramadan fasting on markers of oxidative stress and serum biochemical markers of cellular damage in healthy subjects. *Ann Nutr Metab.* 2008;53:175–181. doi:10.1159/000172979
30. Aksungar FB, Eren A, Ure S, Teskin O, Ates G. Effects of intermittent fasting on serum lipid levels, coagulation status and plasma homocysteine levels. *Ann Nutr Metab.* 2005;49:77–82. doi:10.1159/000084739
31. Trabelsi K, Stannard SR, Ghilissi Z, et al. Effect of fed-versus fasted state resistance training during Ramadan on body composition and selected metabolic parameters in bodybuilders. *J Int Soc Sports Nutr.* 2013;10:23. doi:10.1186/1550-2783-10-29
32. Amer H, Eid Y, Aboromia M. Ramadan in Egyptian people with type 2 diabetes: effect on cardiometabolic parameters. *British Journal of Diabetes.* 2022; 29;22(1):14-9.
33. Oueslati I, Kardi A, Boukhatia F, Hammami B, Cheikh M, Romdhane NB, Feki M, Yazidi M, Chihaoui M. Impact of Ramadan intermittent fasting on metabolic and inflammatory profiles in type 2 diabetic patients. *Journal of Diabetes & Metabolic Disorders.* 2022 ;21(1):751-8.

34. Hassanein, M. M., Hanif, W., Malek, R., & Jabbar, A. (2021). Changes in fasting patterns during Ramadan, and associated clinical outcomes in adults with type 2 diabetes: A narrative review of epidemiological studies over the last 20 years. *Diabetes research and clinical practice*, 172, 108584.
35. Abdelrahim D, Faris ME, Hassanein M, Shakir AZ, Yusuf AM, Almeneessier AS, BaHammam AS. Impact of Ramadan diurnal intermittent fasting on hypoglycemic events in patients with type 2 diabetes: a systematic review of randomized controlled trials and observational studies. *Frontiers in Endocrinology*. 2021; 8;12:624423.
36. Akhtar AM, Ghouri N, Chahal CA, Patel R, Ricci F, Sattar N, Waqar S, Khanji MY. Ramadan fasting: recommendations for patients with cardiovascular disease. *Heart*. 2022; 1;108(4):258-65.
37. Hassanein M, Afandi B, Ahmedani MY, Alamoudi RM, Alawadi F, Bajaj HS, Basit A, Bennakhi A, El Sayed AA, Hamdy O, Hanif W. Diabetes and Ramadan: practical guidelines 2021. *Diabetes research and clinical practice*. 2022; 8:109185.

UNDER PEER REVIEW