

# INFLUENCE OF SEED PRIMING AND FOLIAR SPRAY OF NUTRIENTS ON GROWTH AND YIELD OF *RABI* SORGHUM (*Sorghum bicolor* L.)

## Abstract

The experimental trial was conducted at Agricultural Research Station, Tandur, Professor Jayashankar Telangana State Agricultural University (PJ TSAU), for three consecutive years *viz.* 2015-16, 2016-17 and 2017-18 to study the effect of seed priming and foliar spray of nutrients on growth and yield of *Rabi* Sorghum during the winter season. The treatment combinations were laid out in split plot design with two factors Factor 1: Seed priming (5) 1. Priming for 12 Hours in Water 2. Priming with ZnSO<sub>4</sub> (2% solution) 3. Priming with K<sub>2</sub>HPO<sub>4</sub> (2% solution). 4. Priming with KNO<sub>3</sub> (2% solution) and 5. Control (No Priming). Factor 2: Foliar Spray (4) 1. KNO<sub>3</sub> @ 2%, 2. DAP @ 2%, 3. Urea @ 2% and 5. Control (Simple water spray) with three replications. The pooled results of the trial indicated that seed treatment with KNO<sub>3</sub> @ 0.5% and seed priming with KH<sub>2</sub>PO<sub>4</sub> @ 0.5% recorded significantly on par and highest among all other seed priming treatments with respective to the parameters on. Days to 50 % flowering (71 & 72 days), 100 seed weight (3.6 and 3.57 g), grain yield 2.73 & 2.70 t/ha, dry fodder yield (5.68 & 5.70 t/ha) respectively. The 100 seed weight and grain yield recorded significantly on par by foliar spraying of KNO<sub>3</sub> @ 2 % and DAP @ 2 %. (3.60 & 3.62 g) and (2.70 & 2.75 t/ha) respectively over the control.

**Key words:** seed priming, foliar spray, *Rabi* Sorghum and Yield

## INTRODUCTION

Sorghum is an important cereal crop of poor, small and marginal farmers in semi-arid regions of the world. Yield and quality of the sorghum crop often suffers due to presence of insufficient soil moisture during its growth period. The productivity of *rabi* sorghum is dependent on quantity of rains during pre-season monsoon and water holding capacity of soil, use of moisture conservation practices, use of high yielding cultivars on basis of soil types and available production technologies. *Rabi* sorghum is valued mainly for direct human food consumption and fodder for livestock.

In India, *Rabi* Sorghum is cultivated in an area of 30.69 L ha with a production of 30.75 L t and productivity of 1002 kg ha<sup>-1</sup>. In Telangana, it occupies an area of 0.31 L ha with a production 0.56 L t and an average yield of 1818 kg ha<sup>-1</sup> (www.Indiastat.com, 2019-20). It is an important winter season crop grown in Telangana, Andhra Pradesh, Maharashtra and Karnataka. In Telangana it is confined to Ranga Reddy, Mahabubnagar, Adilabad, Nizamabad and Khammam districts. Among different dryland crops, the crops like *Rabi* Sorghum which are grown in post rainy season under receding soil moisture conditions frequently suffer due terminal drought conditions. Among various drought mitigation options, seed priming and foliar nutrition known to impart tolerance to drought in several other crops. Timely and uniform

germination are two essential prerequisites to increase yield and quality in crops. Homogeneity and enhanced seedling emergence of direct-seeded crops have great impact on yield and quality (Gupta et al. 2008). Seed priming is a pre-sowing seed treatment that involves the controlled imbibition of seeds, sufficient to allow pre-germinate metabolic events to take place, but radical emergence does not occur (Heydecker, 1975). Priming is an enhancement method that increases germination and emergence (Khan, 1992; Taylor, 1998). Foliar application is widely used to mitigate the nutrient stress in many crops. Soluble salts are generally effective in foliar sprays. Deficiency symptoms are usually corrected within the few days and after there the entire field could be sprayed with the appropriate micronutrients source. Keeping this in view, the present investigation was under taken to study the effect of seed priming and foliar spray on yield of rabi sorghum during rabi seasons of 2015-16, 2016-17 and 2017-18.

## **MATERIAL AND METHODS**

A field experiment was conducted at Agriculture Research Station, Tandur, Professor Jayashankar Telangana State Agricultural University for three consecutive years 2015-16, 2016-17 and 2017-18. The soil of the experimental field was clay in texture, low in organic carbon (0.22 %), low in available nitrogen ( $148 \text{ kg ha}^{-1}$ ), high in available phosphorus ( $34 \text{ kg ha}^{-1}$ ), high in available potassium ( $418 \text{ kg ha}^{-1}$ ) and slightly alkaline in reaction (pH 7.35). The treatment combinations were laid out Split Plot design with two factors Factor 1: Seed priming (5) 1.Priming for 12 Hours in Water 2.Priming with  $\text{ZnSO}_4$  (2% solution) 3.Priming with  $\text{K}_2\text{HPO}_4$  (2% solution). 4. Priming with  $\text{KNO}_3$  (2% solution) and 5.Control (No Priming). Factor 2: Foliar Spray (4) 1. $\text{KNO}_3$  @ 2%, 2.DAP @ 2%, 3.Urea @ 2% and 5.Control (Simple water spray) with three replications. After priming seeds were dried in shade for about one hour and then used for sowing. The foliar spraying was done at 55 days after emergence (at boot leaf stage). The crop was sown with the spacing of  $45 \times 20$  cm. The fertilizer application was done as per recommended dose of  $60 \text{ kg N}$  and  $40 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$  by placement method. Nitrogen was applied through urea (46% N), phosphorus in the form of single super phosphate (16%  $\text{P}_2\text{O}_5$ ) and which was applied along the marked lines (i.e. line placement) 5 cm below the soil. Net plot yields were used for calculating yield per hectare. The results were analyzed using standard statistical procedure given by Panse and Sukhatme (1967).

## **RESULTS AND DISCUSSION**

### **SEED PRIMING**

The pooled analysis of results revealed that (Table.1), there is a significant difference between the growth and yield parameters of *Rabi Sorghum*. Among the treatments of seed priming with  $\text{KNO}_3$  @ 0.5% for 12 hours produced significantly higher plant height (172 cm), while seed priming with  $\text{KNO}_3$  @ 0.5% and seed priming with  $\text{KH}_2\text{PO}_4$  @ 0.5% recorded significantly on par and highest among all other seed priming treatments with respect to the parameters of days to 50% flowering (71 & 72 days), 100 seed weight (3.6 and 3.57 g), grain yield 2.73 & 2.70 t/ha, dry fodder yield (5.68 & 5.70 t/ha) respectively. Harvest index did not exhibit any significant variation among the different seed priming treatment. This might be due to the seeds priming resulted in earlier emergence of seedlings by 1-3 days and significantly increased plant stand and initial growth vigour. Similar results were obtained by Musa et al. (1999). Srivastava and Bose (2012) reported the beneficial effect of priming treatments [ $\text{Mg}(\text{NO}_3)_2$ ] and  $\text{KNO}_3$ ) which was clearly exhibited in plant height, leaf area and number of leaf and yield attribute characteristics in rice. Ahmadvand et al., (2012) reported that seed priming with  $\text{KNO}_3$  caused a significant increase in germination and emergence percentage, radicle and plumule length, seedling dry weight, plant height, plant leaf area and plant dry weight in soybean.

### **FOLIAR SPRAY**

The plant height (cm) did not exhibit any significant difference by the foliar spray treatment. The days to 50 % percent flowering was significantly higher and on par with the foliar spray of  $\text{KNO}_3$  @ 2 %, DAP @ 2 % and urea @ 2% spray over the control. It recorded 70.93, 70.47 and 70.47 respectively. While the 100 seed weight and grain yield recorded significantly on par by foliar spraying of  $\text{KNO}_3$  @ 2 % and DAP @ 2 %. (3.60 & 3.62 g) and (2.70 & 2.75 t/ha) respectively over the control. While, the dry fodder yield recorded higher yield of 5.69 t/ha with urea @ 2% spray over the other foliar sprayings. Harvest index did not have any significant change by the foliar spray of the nutrients. This might be due to prolonged vegetative growth which increased the plant height (Sivamurugan *et al*, 2018). This result is in conformity with the Alston (1979). Poornima and Koti (2019) indicated that higher grain yield was recorded in 500 ppm Nano ZnO Foliar Spray as compared to 1000 ppm Bulk  $\text{ZnSO}_4$  Foliar Spray. The grain yield of ragi was increased by 31.49 % due to the application of urea spray followed by 19:19:19 spray which has recorded 30.37 % increase in yields (Gokul and Senthilkumar). The findings of

Dudhade and Gadakh (2020) reported that foliar spray of  $\text{KNO}_3$  increased the grain yield of sorghum to 13.08 per cent over the control.

The combination of seed priming and foliar spray of the nutrients (Table.2) resulted in significant yield increase over the control. The combination of seed priming with  $\text{ZNSO}_4$  @ 0.5% and foliar spraying of  $\text{KNO}_3$  and DAP @ 2% spray resulted in higher grain yield of 2.98 and 2.91 t/ha respectively, while the combination of seed priming with  $\text{KNO}_3$  and foliar spray of  $\text{KNO}_3$  and DAP @ 2 % and Urea @ 2% resulted in highest and on par grain yield of rabi sorghum(3.09, 2.99 and 2.77 t/ha) respectively. While that of the combination of seed priming with  $\text{KH}_2\text{PO}_4$  and  $\text{KNO}_3$  and DAP @ 2% spray resulted in highest and on par grain yield of 2.87 and 2.98 t/ha respectively over the control. Bhadane et al., (2020) reported that seed priming with 2%  $\text{CaCl}_2$  followed by 1% foliar spraying at 30 DAS significantly improved most of morpho-physiological parameters and yield contributing traits in Green gram. Seed treatment with bulk  $\text{ZnSO}_4$  has produced higher total dry matter and grain yield but grain zinc content was highest in nano  $\text{ZnO}$ . (Poornima and Koti, 2019). While Dudhade and Gadakh (2020) reported by combined effect of seed priming  $\text{KNO}_3$  @ 0.5% and foliar spray of  $\text{KNO}_3$  @ 2 % produced higher grain yield of 3.0 t/ha and fodder yield 7.6 t/ha respectively.

## **ECONOMICS**

Gross returns of Rupees 98280  $\text{ha}^{-1}$ , Net returns of rupees 75780  $\text{ha}^{-1}$  and benefit cost ratio were recorded highest with the seed priming of  $\text{KNO}_3$  @ 0.5% for 12 hours. While Gross returns of Rupees 99000  $\text{ha}^{-1}$ , Net returns of rupees 76500  $\text{ha}^{-1}$  and benefit cost ratio of 3.40 accrued with the foliar spray of DAP @ 2% were recorded highest. This might be due to higher yield of crop with these treatments. Similar results were earlier reported by (Poornima and Koti, 2019) Dudhade and Gadakh (2020) in Rabi Sorghum.

## **CONCLUSION**

From the experimental results, it can be concluded that seed priming with  $\text{KNO}_3$  @ 0.5% or  $\text{ZNSO}_4$  for 10-12 hrs and foliar spray of  $\text{KNO}_3$  and DAP @ 2% at 55 DAS is equally effective as compared to other seed priming and foliar sprays. Seed priming has been used to improve germination, reduce seedling emergence time, improve stand establishment and yield. Due to foliar spray increase in seed yield observed was due to more absorption of micronutrients through leaves and flowers.  $\text{KNO}_3$  and DAP foliar spray mitigates the water stress during critical stages of flowering and grain formation.

## REFERENCES

- Ahmadvand, G., Soleimani, F., Saadatian, B. and Pouya, M. 2012. Effect of seed priming on germination and emergence traits of two soybean cultivars under salinity stress. *Int Res J Appl Basic Sci.* 3, 234-41.
- Alston, A.M. 1979. Effects of soil water content and foliar fertilization with nitrogen and phosphorus in late season on the yield and composition of wheat. *Aus. J. Exper. Agric. Animal Husbandry.* 30(4), 577-585.
- Bhadane R.S., Prajapati K.R., Ombase K.C. and Patel D.B. 2020. Effect of Seed Priming and Foliar Spraying of PGRs on Morpho-Physiology, Growth and Yield in Green gram (*Vigna radiata* L.) *Legume research*, DOI: 10.18805/LR-4434
- Dhudande D.D and Gadakh S.S. 2020 Effect of seed priming and foliar spray on grain and fodder yield of Rabi Sorghum. *Journal of Progressive Agriculture.* 11(2):69-73.
- Gokul G. and Senthilkumar N. 2019. Effect of foliar feeding technique on growth and yield of ragi. *Plant Archives*, 19(2), 2889- 2892.
- Gupta, A., Kumar, A., Roy, M.B., Naseem, M., Chaudhary, V.K., Maiti, R.K., 2008. Seed Priming: Aftermath. *International Journal of Agriculture, Environment and Biotechnology.* 1, 219-212.
- Heydecker, W., Higgins, J. and Turner, Y.J. 1975. Invigoration of seeds. *Seed Science and Technology.* 3, 881-888.
- Indiastat, 2020. [https://www.indiastat.com/table/agriculture-data/2/jowar-great\\_millet/17197/1131129/data.aspx](https://www.indiastat.com/table/agriculture-data/2/jowar-great_millet/17197/1131129/data.aspx)
- Khan, A.A. 1992. Preplant physiological and seed conditioning. In: Janik, J. (Ed.), *Horticultural Reviews.* John Wiley and Sons, New York. pp.131-181.
- Musa, A. M., Johansen, C., Kumar, J. and Harris, D. 1999. Response of chickpea to seed priming in the High Barind Tract of Bangladesh. *Inter Chickpea and Pigeonpea Newsletter.* 6, 20-22.
- Panase, V. G and Sukhatme, P. V.1967. *Statistical methods for agricultural workers.* ICAR, New Delhi, 1967, 199-200.
- Poornima R. and Koti R.V. 2019. Effect of nano zinc oxide on growth, yield and grain zinc content of sorghum (*Sorghum bicolor*). *Journal of Pharmacognosy and Phytochemistry.* 8(4), 727-731

Srivastava, A. K. and Bose, B. 2012. Effect of nitrate seed priming on phenology, growth rate and yield attributes in rice (*Oryza sativa* L.). *Vegetos*, 25, 174-81.

Taylor, A.P., Allen, P.S., Bennet, M.A., Bradford, K.J., Burns, J.S., Misra, M.K., 1998. Seed characteristics. *Seed Science Research*. 8: 245-256.

UNDER PEER REVIEW

**Table 1. Effect of seed priming and foliar spray on plant height, yield attributes, yield and economics of rabi sorghum. (Pooled means 2014-15, 2015-16 and 2016-17)**

<b>Treatment</b>	Plant height (cm)	Days to 50% flowering	100 Seed weight (g)	Grain yield (t/ha)	Dry fodder yield (t/ha)	Harvest index (%)	Gross returns (Rs/ha)	Net returns (Rs/ha)	B:C Ratio
<b>Seed priming for 12 hours</b>									
P 1 Control (No priming)	153	67	3.18	2.11	4.23	33.28	75960	53585	2.39
P 2 Seed priming with water	159	69	3.31	2.40	5.44	30.61	86400	64025	2.86
P 3 Seed priming with ZNSO <sub>4</sub> (0.5%)	161	69	3.51	2.58	5.36	32.49	92880	70380	3.13
P 4 Seed priming with KNO <sub>3</sub> (0.5%)	172	72	3.60	2.73	5.68	32.46	98280	75780	3.37
P 5 Seed priming with KH <sub>2</sub> PO <sub>4</sub> (0.5%)	159	71	3.57	2.70	5.70	32.14	97200	74700	3.32
SE m	4.52	0.64	0.09	0.05	0.20	0.96			
C.D. (5%)	9.21	1.32	0.18	0.13	0.56	NS			
<b>Foliar spray</b>									
Control (water spray)	156	66.33	3.31	2.17	4.84	30.96	78120	55820	2.50
KNO <sub>3</sub> @2%	163	70.93	3.60	2.70	5.20	34.01	97200	74700	3.32
DAP @2%	159	70.47	3.62	2.75	5.40	33.74	99000	76500	3.40
Urea @2%	160	70.27	3.27	2.40	5.69	29.67	86400	63900	2.84
SE m	3.56	1.17	0.07	0.04	0.17	0.90			
C.D. (5%)	NS	2.29	0.15	0.12	0.50	1.70			
C.V. (%)	7.00	6.81	5.94	6.77	7.48	7.15			
<b>Interaction PXF</b>									
SE m	7.2	0.7	0.12	0.13	0.31	1.58			
C.D. (5%)	NS	NS	NS	0.27	NS	NS			

**Table 2. Pooled means of Interaction effect of seed priming and foliar spray on grain yield t/ha (2014-15, 2015-16 and 2016-17).**

<b>Seed priming with foliar spray</b>	<b>(water spray)</b>	<b>KNO<sub>3</sub> @ 2%</b>	<b>DAP @ 2%</b>	<b>Urea @ 2%</b>	<b>Mean</b>
<b>No priming)</b>	1.99	2.15	2.11	2.21	2.11
<b>Seed priming with water</b>	2.11	2.41	2.78	2.29	2.40
<b>Seed priming with ZNSo<sub>4</sub> (0.5%)</b>	2.33	2.98	2.91	2.08	2.58
<b>Seed priming with KNO<sub>3</sub></b>	2.08	3.09	2.99	2.77	2.73
<b>Seed priming with KH<sub>2</sub>PO<sub>4</sub> (0.5%)</b>	2.34	2.87	2.98	2.63	2.70
<b>Mean</b>	2.17	2.70	2.75	2.40	

Seed priming	SE (m) 0.05	CD 0.13
Foliar spray	SE (m) 0.04	CD 0.12
Seed priming with foliar spray	SE (m) 0.09	CD 0.27