

## Original Research Article

Effective drought index based analysis of drought characteristics and its propagation in the Vindhyan region

### **Abstract**

Drought is a disastrous natural phenomenon that significantly impacts socio-economic, agricultural, and environmental spheres and is considered as the primary factor of uncertainty that continues to haunt Indian agriculture and its economy. In this context, the drought scenario in the Mirzapur and Sonbhadra districts of the Vindhyan region motivates the investigation of drought characteristics through an index called Effective Drought Index (EDI). The analysis revealed that the region faced drought during 1989, 1992, 1998, 2005, 2009, 2014, and 2018. Maximum extreme drought events occurred in Citynagar block (21 events) followed by Kon block (20 events) in Mirzapur district, while blocks located in Sonbhadra district of the study area face 1 to 6 extreme drought events. Haliya block (28 events) in Mirzapur district and Myorpur block (26 events) in Sonbhadra district faces maximum severe drought events. Chhanvey block in Mirzapur district faces 25 severe drought events, while the rest of the block of the study area faces severe drought conditions that vary between 6 to 22 events. The severity of extreme drought is maximum in Citynagar block (-47.46) followed by Kon block (-44.87) in Mirzapur district. Maximum drought severity based on EDI computation used to prevails in Chopan block (-170.48) in Sonbhadra block followed by Haliya block (-171.12) in Mirzapur district. Maximum drought events of varying severity classes used to prevail in Robertsganj block (172 events) in Sonbhadra district. The analysis unveils the characteristics of drought that prevail in the region and through a thorough inspection suitable strategy can be synthesis that renders the impact of drought.

**Keywords:** Drought, Effective Drought Index (EDI), Vindhyan region, Drought characteristics

## Introduction

Drought is the imbalance in hydro-meteorological parameters due to the appreciable deficit in rainfall and combined episodes of dry spells (Mishra and Singh 2010). Lack of sufficient precipitation causes meteorological drought, which is subsequently converted to agricultural and hydrological drought if the rainfall is inadequate to replenish the amount of moisture depleted due to evapotranspiration and absorption by plants (Spinoni et al., 2019; Wilhite and Glantz, 1985). Since drought is the consequence of natural and manmade activities and the circumstances created induce to classify drought based on its impacts, drought is classified as meteorological, agricultural, hydrological, and socioeconomic (Paton et al., 2000).

Meteorological drought is defined as the deficit in the rainfall amount. It is the leading elucidation for other categories of drought events. When meteorological drought causes a significant reduction in the environmental flows which helps to thrive river and stream then resulting effect converted to hydrological drought. Inadequate soil moisture and perseverance of meteorological drought over a continuous period of time result to cause agricultural drought. Drought with varying degrees of severity and intensity globally causes a significant loss which is equivalent to more than 6–8 billion US\$ annually (Pandey et al. 2008).

Studies and extensive research highlighted the difficulty in detecting and investigating drought and its complexes hence developing different drought indices that are area-specific and region-oriented. Most of the drought indices have their own advantages and disadvantages. Thus, indices requiring less cumbersome data that are readily available and less complicating are pretty extensively used. One such index is the Effective Drought Index (EDI), which has been used in the present study to detect drought on a monthly basis. The proposed index defined drought as the objective measure of daily water accumulation based on the weight of the current and antecedent rainfall (Byun and Wilhite 1999).

The EDI concept was considered in following recent researches on droughts (Oh et al. 2014, 2015; Deo et al., 2015; Kumar et al., 2018; Sharma et al., 2019). The main strength of this drought indicator is its ability to detect the onset and end of the drought and drought conditions before the other indices (Jain et al., 2015; Zarei et al., 2017). Most of the current indices use normalized statistical metrics to analyze the deficit periods (Mishra and Singh, 2010, 2011), while their multi-scale drought characterization ranges from the month (smallest

time scale) to longer periods (Deo et al., 2017). These indices are not considered the daily consecutive or accumulated stress of drought (Byun and Wilhite, 1999). Drought severity is mostly calculated from the climatological mean of water deficiency for some predefined duration without considering the diminishing water resources over time.

Studies have confirmed that EDI is more efficient than the SPI in assessing both short term (e.g. Daily, weekly and monthly) and long-term (e.g. Seasonal and annual) droughts (Byun et al., 2010; Dogan et al., 2012). Byun and Kim (2010) compared the EDI with the SPI, and their analysis indicates that EDI detects short term/long term drought that cannot be detected by the long term/short term SPIs, respectively. EDI considers any rainfall amount and rainfall day, one single value for the day or period, long memory of rainfall (365 days) and therefore the consecutive weight of rainfall; confirming that EDI is superior to SPI in measuring the drought severity (Byun and Kim, 2010). Moreover, Morid et al. (2006), cited in Deo et al. (2017), indicates that EDI is best for detecting the beginning and end of drought than the percent of normal, SPI, China Z index and therefore the Z score. EDI are often used due to its self-defined time step, which is free from fixing time step problem (Byun and Wilhite, 1999; Deo et al., 2017; Jain et al., 2015).

Kar et al., (2016) studied the drought severity using EDI for the Bundelkhand region and found a gradually increasing severity in the region. Kumar et al., (2018) characterized meteorological drought using effective drought index for Banswara district, Rajasthan. As the area under consideration is completely rain-fed, variation in rainfall amount during the monsoon period severely affects crop growth. Due to the wide application of EDI for drought investigation and drought monitoring, it has been used in this research paper to assess the meteorological drought characteristics in the Vindhyan region with the aim the results obtained help to formulate efficient mitigation planning during stress conditions.

## **Methodology**

### **Study area**

The Vindhyan region in Mirzapur and Sonbhadra district of Uttar Pradesh lies between 82° to 83°23' E longitude and 22°45' to 24°34' N latitude. The region has an area of 11,309 sq km with an average elevation of 298 m. The region has a relatively subtropical climate with high variation in temperatures during summer (avg. 32°C- 42°C) and winter season (avg. 2°C-15°C). The average annual rainfall in the area is 935 mm, about 90% of which occurs between June and September (monsoon period). Surplus runoff water flows

through ephemeral creeks and streams during the monsoon season, ultimately meeting to river Ganges. Due to its hilly topographic features, the surface runoff is mostly not harvested in the study area. Winds are generally strong in the area, increasing force during summer and southwest monsoon season. The mean wind velocity is 4 kmph and average potential evapotranspiration rate is 1456.7 mm. In the region, primarily alluvial soil is found. Districts in the Vindhyan region comprise Archaean to Neoproterozoic rocks and Quaternary flood plain/ alluvial deposits. The location map of the study area is given in Figure 1.




Figure1: Location map of the study area

### **Data availability**

Gridded daily rainfall data has been used to investigate drought and its characteristics, which was obtained from Indian Meteorological Department (IMD), Pune. The data consists of 40 years period from 1980 to 2019. ArcGIS 10.4 software has been utilized for delineating drainage map and is given in Figure 2a. Land use and Land cover map of the region was obtained from Decadal map of India, which was available online and LULC map of

Vindhyan region is shown in Figure 2b. In the Vindhyan region, 8 blocks lie in Sonbhadra district and 12 blocks are lying in Mirzapur district, whose locations are given in Table 1.

Table 1: Location of blocks in Mirzapur and Sonbhadra district of Vindhyan region

District Name	Block	Lat	Long	Avg. Rainfall (mm)
Mirzapur	Chhanvey	25.17	82.37	900.09
	Citynagar	24.91	82.56	916.57
	Haliya	24.8	82.3	1003.26
	Jamalpur	25.16	83.08	863.34
	Kon	25.2	82.53	900.78
	Lalganj	24.99	82.38	923.56
	Madihaon	24.92	82.57	941.89
	Majhawan	25.22	82.7	973.13
	Narainpur	25.13	82.95	871.46
	Pahari	25.07	82.68	907.45
	Rajgarh	24.98	82.86	893.09
	Sikhdi	25.15	82.79	959.42
Sonbhadra	Babhani	23.95	83.1	962.31
	Chatra	24.66	83.24	974.89
	Chopan	24.45	82.99	968.17
	Dudhi	24.22	83.23	943.02
	Ghorawal	24.75	82.81	976.82
	Myorpur	24.14	82.88	901.63
	Nagwa	24.62	83.4	943.93
	Robertsganj	24.69	83.06	970.36




Figure 2a): Drainage Map; 2b): Land use and Land cover map

### Effective Drought Index (EDI)

Byun and Wilhite (1999) proposed Effective Drought Index (EDI) and stated that it can deal with all of the limitations of SPI. EDI provides more information regarding the duration and severity of rainfall deficit and the start and end of the drought period. Effective precipitation helps to represent daily depletion of water (Morid *et al.*, 2006). The original form of EDI is computed with a daily time step using daily rainfall data. Smakhtin and Hughes (2007) modified his algorithm to be tested with monthly data.

Byun and Wilhite (1999) suggested the following equation for EP:

$$EP_i = \sum_{n=1}^i \frac{\sum_{m=1}^n P_m}{n} \quad \text{.....Eq. (1)}$$

Where,  $EP_i$  = valid effective precipitation,  $P_m$  = daily precipitation,  $m$  = number of days before a specific day,  $i$  = the number of days of the time window,  $n$  = running from 1 till  $i$  (Chhajer *et al.* 2015).

$$DEP = EP / MEP \quad \text{.....Eq. (2)}$$

Where, EP = effective precipitation for 365 days counting from a specific day, MEP = the mean of effective precipitation, DEP = deviation of EP from MEP.

EDI is calculated as,

$$EDI = DEP / SD (EP) \quad \text{.....Eq. (3)}$$

SD (EP) = the standard deviation of each day's EP.

Ranks of EDI reflect drought conditions (Byun and Wilhite 1999] indicate: extreme drought ( $EDI \leq -2.0$ ), severe drought ( $-1.99 \leq EDI \leq -1.5$ ), moderate drought ( $-1.49 \leq EDI \leq -1.0$ ) and near normal conditions ( $-0.99 \leq EDI \leq 0.99$ ).

## Result and Discussion

### Evaluation of drought characteristics based on Effective Drought Index (EDI)

The drought over the Vindhyan region was analysed using the EDI method using monthly rainfall time-series data (1980–2019). While evaluating drought based on EDI, the drought severity classification criterion was considered for the evaluation of the results. Temporal variations of drought characteristics are of great significance for understanding the drought pattern in the region. Figure 3 to 6 depicts the temporal variation of drought severity for different blocks of Mirzapur and Sonbhadra district of the Vindhyan region.




Figure 3: Temporal variation of EDI in Jmalpur block in Mirzapur district




Figure 4: Temporal variation of EDI in Narainpur block in Mirzapur district




Figure 5: Temporal variation of EDI in Robertsganj block in Sonbhadra district




Figure 6: Temporal variation of EDI in Ghorawal block in Sonbhadra district

Temporal variation of EDI indicates that the region faced severe drought conditions during 1989, 1992, 1998, 2005, 2009, 2014, and 2018. Table 2 represents drought events evaluated based on EDI. A maximum number of extreme drought conditions occurred in Citynagar block (21 events) followed by Kon block (20 events) in Mirzapur district. Block namely Lalganj, Chhanvey, Madihaon in Mirzapur district faces 15, 14, and 10 extreme drought events, respectively. Most of the blocks located in Sonbhadra district of the study area face 1 to 6 extreme drought events over the span of 40 years. Haliya block in Mirzapur district faces maximum severe drought events (28 events) followed by Lalganj block in Mirzapur district and Myorpur block (26 events) in Sonbhadra district. Chhanvey block in Mirzapur district faces 25 severe drought events while rest of block of the study area faces severe drought condition that varies between 6 to 22 events. Maximum moderate drought events occurred in Robertsganj block (69 events) in Sonbhadra district followed by Narainpur and Rajgarh block (64 events) in Sonbhadra district of the study area. Occurrence of mild drought events in the study area are frequent i.e., once in every two years.

Table 2: Drought events evaluated based on EDI

District Name	Block	Extreme	Severe	Moderate	Mild	Total Drought Events
Mizrapur	Chhanvey	14	25	39	54	132
	Citynagar	21	15	39	62	137
	Haliya	4	28	57	59	148

	Jamalpur	0	8	60	94	162
	Kon	20	17	37	63	137
	Lalganj	15	26	37	56	134
	Madihaon	10	19	49	68	146
	Majhawan	9	13	55	70	147
	Narainpur	1	11	64	92	168
	Pahari	9	6	58	88	161
	Rajgarh	5	8	64	81	158
	Sikhdi	6	12	57	81	156
Sonbhadra	Robertsganj	2	9	69	92	172
	Ghorawal	1	10	60	97	168
	Chatra	1	16	55	85	157
	Nagwa	0	9	63	83	155
	Chopan	1	22	58	84	165
	Babhani	6	22	43	67	138
	Myorpur	3	26	58	60	147
	Dudhi	1	20	59	65	145

Table 3 showed drought characteristics based on EDI. Severity of extreme drought is maximum in Citynagar block (-47.46) followed by Kon block (-44.87) in Mirzapur district whereas in Sonbhadra district Babhani block (-13.04) has maximum extreme drought severity. Maximum severe drought severity prevails in Haliya and Lalganj block (approx. -46) in Mirzapur district. Moderate drought severity is maximum in Nagwa block (-78.96) in Sonbhadra district followed by Rajgarh block (-78.36) in Mirzapur district. Severity of mild drought events was maximum in Ghorawal block (-73.33) in Sonbhadra district followed by Jmalpur block (-70.36) in Mirzapur district. Maximum drought severity based on EDI computation used to prevails in Chopan block (-170.48) in Sonbhadra block followed by Haliya block (-171.12) in Mirzapur district. Maximum drought events of varying severity class used to prevails in Robertsganj block (172 events) in Sonbhadra district. Drought intensity, which was computed as the ratio of total drought severity to the study period, suggested that Chhanvey Kon and Lalganj block have the highest drought intensity compared to other blocks in the study area.

Table 3: Drought characteristics evaluated based on EDI

District Name	Block	Drought severity				Total drought severity	Drought intensity
		Extreme	Severe	Moderate	Mild		
Mizrapur	Chhanvey	-32.1	-44.73	-47.11	-39.52	-163.46	-1.24

	Citynagar	-47.46	-25.46	-48.28	-45.19	-166.39	-1.21
	Haliya	-8.72	-46.38	-71.79	-44.23	-171.12	-1.16
	Jamalpur	0	-14.04	-72.72	-70.36	-157.12	-0.97
	Kon	-44.87	-29.12	-46.3	-47.9	-168.19	-1.23
	Lalganj	-35.13	-46.01	-43.5	-39.47	-164.11	-1.22
	Madihaon	-22.61	-33.38	-61.7	-49.45	-167.14	-1.14
	Majhawan	-19.94	-22.21	-66.15	-52.88	-161.18	-1.1
	Narainpur	-2.09	-18.29	-77.12	-68.36	-165.86	-0.99
	Pahari	-19.77	-9.97	-69.8	-66.6	-166.14	-1.03
	Rajgarh	-11.01	-14.41	-78.36	-64.14	-167.92	-1.06
	Sikhdi	-13.37	-21.76	-70.15	-60.11	-165.39	-1.06
Sonbhadra	Robertsganj	-4.25	-15.11	-83.5	-68.01	-170.87	-0.99
	Ghorawal	-2.02	-16.85	-73.63	-73.33	-165.83	-0.99
	Chatra	-2.01	-26.36	-68.68	-63.5	-160.55	-1.02
	Nagwa	0	-14.16	-78.96	-63.95	-157.07	-1.01
	Chopan	-2.21	-37.05	-72.66	-60.56	-172.48	-1.05
	Babhani	-13.04	-37.92	-51.94	-48.05	-150.95	-1.09
	Myorpur	-6.77	-43.92	-71.83	-45.61	-168.13	-1.14
	Dudhi	-2.35	-33.01	-74.34	-46.19	-155.89	-1.08

## Conclusion

Changes in the climatic phenomenon that occurred due to human-induced activities over a period of time exacerbated the consequences especially concerning drought. The situation has been extensively devastating for India, which has an agrarian economy. The Vindhyan region where agriculture flourishes has been severely affected due to frequent drought occurrence of various severity classes. The present study highlight that in Mirzapur district the Citynagar block face the maximum number of extreme drought events whereas Babhani block in Sonbhadra district. Maximum drought events occurred in the Robertsganj block of Sonbhadra district and minimum drought events occurred in the Chhanvey district of Mirzapur district. The overall occurrence of drought events in Sonbhadra district (155 events) of Vindhyan region is more as compared to Mirzapur district (149 events) whereas average drought severity in Mirzapur district (-166) is more as compared to Sonbhadra district (-162.5). Therefore, it can be concluded that both the district lying in the Vindhyan region is critical to drought, thus suggesting adaptation of suitable mitigation strategies that can subsume the impact of drought.

## References

- Byun HR, Kim DW. Comparing the effective drought index and the standardized precipitation index. *Economics of drought and drought preparedness in a climate change context*. CIHEAM: Zaragoza, Spain. 2010; 95: 85-89.
- Byun HR, Wilhite DA. Objective quantification of drought severity and duration. *Journal of Climate*. 1999; 12(9): 2747-2756.
- Chhajjer V, Prabhakar S, Rama CPP. Development of index to assess drought conditions using geospatial data a case study of Jaisalmer district, Rajasthan, India. *Geoinformatica Polonica*. 2015; 14(1): 29-40.
- Deo RC, Byun HR, Adamowski JF, Begum K. Application of effective drought index for quantification of meteorological drought events: a case study in Australia. *Theoretical and Applied Climatology*. 2017; 128(1): 359-379.
- Deo RC, Byun HR, Adamowski JF, Kim DW. A real-time flood monitoring index based on daily effective precipitation and its application to Brisbane and Lockyer Valley flood events. *Water Resources Management*. 2015; 29(11): 4075-4093.
- Dogan S, Berktaç A, Singh VP. Comparison of multi-monthly rainfall-based drought severity indices, with application to semi-arid Konya closed basin, Turkey. *Journal of Hydrology*. 2012; 470-471: 255-268.
- Jain VK, Pandey RP, Jain MK, Byun HR. Comparison of drought indices for appraisal of drought characteristics in the Ken River Basin. *Weather and Climate Extremes*. 2015; 8: 1-11.
- Kar SK, Thomas T, Singh RM. Assessment of drought characteristics for Dhasan basin in Bundelkhand region. *International Journal of Agriculture, Environment and Biotechnology*. 2016; 9(5): 897-902.
- Kumar CK, Chandola VK, Kumar R. Meteorological Drought Characterization Using Effective Drought Index (EDI) for Banswara District (Rajasthan), India. *International Journal of Current Microbiology and Applied Sciences*. 2018; 7(9): 3407-3416.

- Mishra AK, Singh VP. A review of drought concepts. *Journal of Hydrology*. 2010; 391: 202–216.
- Mishra AK, Singh VP. Drought modeling - A review. *Journal of Hydrology*. 2011; 403:157–175.
- Morid S, Smakhtin V, Moghaddasi M. Comparison of seven meteorological indices for drought monitoring in Iran. *International Journal of Climatology: A Journal of the Royal Meteorological Society*. 2006; 26(7): 971-985.
- Oh K, Yu I, Kim H, Kim S, Kim LH, Jeong S. Drought assessment based on real-time drought index. *Desalination and Water Treatment*. 2015; 53(11): 3111-3117.
- Oh SB, Byun HR, Kim DW. Spatiotemporal characteristics of regional drought occurrence in East Asia. *Theoretical and applied climatology*. 2014; 117(1): 89-101.
- Pandey RP, Dash BB, Mishra SK, Singh R. Study of indices for drought characterization in KBK districts in Orissa (India). *Hydrological Processes: An International Journal*. 2008; 22(12): 1895-1907.
- Paton D, Smith L, Violanti J. Disaster response: risk, vulnerability and resilience. *International Journal of Disaster Prevention and Management*. 2000; 9 (3): 173-180.
- Sharma G, Kumar CK, Singh BP. Assessment of Meteorological Drought for Ujjain District of Madhya Pradesh, India using Effective Drought Index and ArcGIS. *International Journal of Current Microbiology and Applied Sciences*. 2019; 8(5): 604-612.
- Smakhtin VU, Hughes DA. Automated estimation and analyses of meteorological drought characteristics from monthly rainfall data. *Environmental Modelling and Software*. 2007; 22(6): 880-890.
- Spinoni J, Barbosa P, De Jager A, McCormick N, Naumann G, Vogt JV, Magni D, Masante D, Mazzeschi M. A new global database of meteorological drought events from 1951 to 2016. *Journal of Hydrology: Regional Studies*. 2019; 22: 100593.
- Wilhite DA, Glantz MH. Understanding the drought phenomenon: the role of definitions. *Water international*. 1985; 10(3): 111-120.

Zarei A, Asadi E, Ebrahimi A, Jafary M, Malekian A, Tahmoures M, Alizadeh E. Comparison of Meteorological Indices for Spatio-Temporal Analysis of Drought In Chahrmahal-Bakhtiyari Province In Iran. *Hrvatski Meteorološki Časopis, Croatian Meteorological Journal*. 2017; 52, 13-26.

UNDER PEER REVIEW