

**Yield of Mustard as Influenced by Date of Sowing and Varieties in Western Rajasthan**

---

**ABSTRACT**

An experiment was conducted at agricultural research station, SKRAU, Bikaner during *rabi* season of 2017-18 to evaluate yield of mustard as influenced by date of sowing and varieties in western Rajasthan. The treatment consisted of three dates of sowings viz., 10<sup>th</sup> October, 25<sup>th</sup> October and 9<sup>th</sup> November were kept in main plot and five varieties viz. RH-119, NRCHB-101, RGN-48, RH-749 and Laxmi was kept as sub plot replicated thrice in split plot design. Crop sown on 25<sup>th</sup> October recorded significantly higher seed yield as compared to 9<sup>th</sup> November sowing. In case of stover yield of mustard maximum was recorded under 25<sup>th</sup> October as compared to 10<sup>th</sup> October and 9<sup>th</sup> November. Days taken for emergence were no significant difference in varieties. RH-749 taken maximum days for 50 per cent flowering as compared to all varieties and for Siliqua appearance maximum days also taken by RH-749 as compared to NRCHB-101 and it statistically at par with RH-119, RGN-48 and Laxmi. RH-119 days to maturity and no. of branches/plant were minimum as compared to all varieties and these were remained statistically at par with each other. Maximum number of siliqua /plants was recorded under RGN-48 which was superior over rest of varieties and statistically at par with NRCHB-101. Highest grain/siliqua was recorded under NRCHB-101 over RH-119, RGN-48 and RH-749.

**Key words:** Mustard, Date of Sowing, Varieties and Yield

**1. INTRODUCTION**

India is one among the leading oil seed producing countries in the world. Oilseeds form the second largest agricultural commodity after cereals. In India mustard is the second important edible oil seed crop after groundnut. It plays an important role in the oil seed economy of the country. In India, during 2018-19, the rapeseed and mustard crop had production of about 9.34 million tonnes from an area of 6.13 mha with an average productivity of 1499 kg ha<sup>-1</sup>. (GOI, 2018-19). However, in Rajasthan it is grown in 2.79 mha with production of 4.30 million tonnes. The average productivity of Rajasthan is 1586 kg ha<sup>-1</sup>. (GOR, 2019-20). Weather variability is considered one of the

major factors of inter-annual variation of crop growth and yield in all environments and it is more important in rainfed situation. The adverse agro-meteorological events like extreme hot and cold temperatures, the lesser brighter sunny days and irregular and unequal distributions of rains are the major factors for the decreasing growth and yields of any field crops especially under rainfed environment. Rainfalls being vital, the growth phases of any variety of crops are determined basically with growing season in which the atmospheric ambient temperature and solar radiation are considered the major governing factors (Sastry *et al.*, 2000).

## **2. MATERIALS AND METHODS**

A field experiment was conducted at College of Agriculture, Bikaner (28.01 °N latitude and 73.22 °E longitude at an altitude of 234.7 M above mean sea level). The soil was loamy sand, low in organic carbon (0.08%) and available N (78 kg/ha) and medium in available phosphorus (22 kg P/ha) and available K (210 kg/ha) with pH 8.3. The experiment was laid-out in split plot design with three replications having fifteen treatment combinations of three dates of sowings viz., 10<sup>th</sup> October, 25<sup>th</sup> October and 9<sup>th</sup> November (in main plot) and five varieties viz. RH-119, NRCHB-101, RGN-48, RH-749 and Laxmi (in sub plot) were used to evaluate their effect on productivity of different varieties. The data related to growth, yield attributes and yield of mustard were recorded and statistically analyzed as per guidelines of Fischer (1950)

## **3. RESULTS AND DISCUSSION**

### ***Effect of irrigation scheduling***

Thermo and photosensitivity of oil seed crop limits the sowing period. Very early sowing causes the mortality of tender seedling due to high temperature. However delayed sowing also reduces the production due to poor performance of the crop as the result of reduced growing period, has tens maturity and ultimately causes the reduction in yield. Besides, westerly hot winds accelerate the maturity and thereby reduce the yield. Therefore, to harvest a good crop, sowing at appropriate time which provides congenial environment is utmost important. Initial plant population of the crop was maintained properly by thinning. Since soil moisture in the seedling zone was not a limiting factor for germination of the seeds of the crop pre-sowing irrigation was applied in all the three dates of sowing. The dates of sowing brought significant variation on different agronomical parameters recorded at successive stages of Indian mustard, viz.

Day to emergence, Plant height, days taken to 50 % flowering, Days to Siliqua appearance, Number of branches/plant, and Days to maturity. Early sown crop (10 October) taken minimum days to emergence and had significant variation to normal sowing (25th October) and late sowing (09 November). Dates of sowing had significant influence on 50 per cent flowering of mustard. Due to higher number of branches/plant recorded at harvest as shown in (Table 2). This could be ascribed due to prolonged vegetative growth period because of congenial environmental conditions, especially atmospheric temperature which formed a basis for rapid cell division in the meristematic tissues of the experimental crop which led to better growth attributes under normal sowing (25th October). The early sown crop experienced sub-optimal temperature regime, especially first fortnight of October which retarded their growth as compared to those sown either normal date 25th October or 09 November. Shorter plant under delayed sowing of mustard crop in India have also been reported by Singh and Singh (2002) and Kumar *et al.* (2002), Yield attributes like number of siliqua /plant, number of seeds/Siliqua and 1000 seed weight were successively decreased with early sowing of the crop (Table 2). However, the difference between in number of seeds/Siliqua between on 25th October and 15th October sowing were non-significant. Early and late sowing restricted the crop growth duration and also induced early flowering. It is also reduced pod initiation and seed setting to a great extent as compared to 25th October. This might be due to the fact that low temperature during the month of September not only restricted the crop growth but also induced lower rate of pollination and flower abortion, shedding in early sown mustard crop. Similar results were also observed by, Singh *et al.*, 2001, Singh and Singh 2002, and Panda *et al.*, 2004). Seed and Stover yield of mustard were significantly affected due to different dates of sowing (Table 3). Crop sown on 25th October recorded significantly higher seed yield as compared to 9th November sowing mainly due to better translocation of photosynthates from source to sink. In case of stover yield of mustard maximum yield were recorded under 25 October as compared to 10 October and 9th November; this might be due to poor growth and better translocation of photosynthates from source to sink. All the growth and yield attributes which determined the seed and stover yield of mustard crop, were adversely influenced when the sowing was done on early dates. Significant reduction in seed and stover yield of mustard in early have also been reported by several other workers Singh *et al.*, 2001, Sihag *et al.*, 2003, Panda *et al.*, 2004.

### ***Effect of varieties***

Days taken for emergence were no significant difference in varieties. RH-749 taken maximum days for 50 per cent flowering as compared to all varieties and for Siliqua appearance maximum days also taken by RH-749 as compared to NRCHB-101 and it statistically at par with RH-119, RGN-48 and Laxmi. The probable reason may be attributed to genetic characters of Coral-437 which has higher capacity to utilized the photosynthates more efficiently through maximum leaf area index, number of branches/plant and ultimately the dry matter production, the similar findings have been reported by Kumar *et al.*, 2000, Shukla *et al.*, 2001 and Chaplot *et al.*, 2012. RH-119 taken minimum days to maturity and number of branches/plant as compared to all varieties and these remained varieties were statistically on par with each other. Maximum number of siliqua/plant was recorded with RGN-48 which was superior over rest of varieties and statistically at par with NRCHB-101. Highest grain/siliqua was recorded under NRCHB-101 superior to RH-119, RGN-48 and RH-749. It was attributed due to genetic characters which truly indicated of total photosynthates production, have been reported by several other workers: Kumar *et al.*, 2000, Shukla *et al.*, 2001 and Chaplot *et al.*, 2012. NRCHB-101 recorded highest grain yield over RH-119, RGN-48 and RH-749 and statistically at par with Laxmi. NRCHB-101 given 13.28 per cent higher seed yield over RH-119. Stover yield and harvest index were not influenced by varieties. Varieties have significantly variation in test weight and maximum recorded under Laxmi varieties of mustard which statistically at par with RH-119 and NRCHB-101. The similar resulted also recorded by Shukla *et al.*, 2001.

**Table 1: Effect of date of sowing and varieties on phenophases of mustard**

Treatments	Day to emergence	Days to 50% Flowering	Days to Siliqua appearance	Days to maturity	No. of Branches/plant
<b>Date of sowing</b>					
10 October	5.2	50.1	66.7	139.7	11.1
25 October	6.1	55.4	74.4	134.9	11.6
09 November	6.4	62.9	82.0	126.5	8.8
S. Em.±	0.1	0.4	0.4	0.5	0.3
CD (p=0.05)	0.4	1.1	1.0	1.4	0.9
<b>Varieties</b>					
RH-119	5.9	56.2	74.2	130.8	8.8
NRCHB-101	6.1	52.9	71.9	134.4	11.2
RGN-48	5.8	56.6	75.6	134.9	11.1
RH-749	5.7	59.2	76.0	134.3	10.1
LAXMI	6.0	55.7	74.2	134.1	11.4
S. Em.±	0.2	0.5	0.5	0.6	0.4
CD (p=0.05)	NS	1.5	1.3	1.8	1.2

**Table 2 : Effect of date of sowing and varieties on yield attributes, yield and harvest index of mustard**

Treatments	No. of siliqua /plant	No. of grain/siliqua	Grain Yield (Kg/ha)	Straw Yield (Kg/ha)	HI (%)	TW (g)
<b>Date of sowing</b>						
10 October	211.6	15.7	4018.9	16073.3	25.2	6.3
25 October	236.1	15.9	4106.7	18182.6	22.7	6.3
09 November	194.9	16.8	2693.7	10226.3	26.8	6.4
S. Em.±	2.8	0.2	113.7	633.6	0.5	0.0
CD (p=0.05)	8.1	0.6	331.9	1849.5	1.4	0.1
<b>Varieties</b>						
RH-119	209.7	15.9	3267.3	13226.5	25.2	6.5
NRCHB-101	213.5	17.8	3767.9	15102.3	25.7	6.4
RGN-48	224.1	14.2	3669.8	15365.6	24.3	6.1
RH-749	207.6	16.0	3614.8	14939.2	24.7	6.0
LAXMI	216.0	16.7	3712.3	15503.3	24.5	6.6
S. Em.±	3.6	0.3	146.8	818.0	0.6	0.1
CD (p=0.05)	10.4	0.8	428.5	2387.7	1.8	0.2

## CONCLUSION

Based on the findings of the present study, it may be concluded that the sowing of wheat cultivar Raj 4079 on 15<sup>th</sup> November with irrigation scheduling at 1.2 ETc was found better with respect to yield attributes and yield in relation to economics as compared to other cultivars. This remained at par with irrigation scheduling at 1.0 ETc under 15<sup>th</sup> November sown crop. However, under late sowing (15<sup>th</sup> December) the cultivar Raj 4238 produced higher grain yield as compared to all other cultivars. This study helps to the farmers in gaining an accurate knowledge of the optimum sowing window and efficient irrigation scheduling approach to achieve crops productive potential and higher yield.

## References

- Chaplot, P.C. Vandeeep, Amita and Kumar Rakesh. Effect of balanced fertilization and Agrochemicals on growth, yield attributes and yield of mustard varieties. 3rd *International Agronomy Congress Nov.*, 2012, 26-30, New Delhi, 1110-1111.
- Fisher RA. *Statistical Methods for Research Workers*, Oliver and Boyd, Edinburg, London, 1950, pp.57-63
- Kumar, R., Singh, D. and Singh, H. Growth and yield of *Brassica* species as influenced by sulphur application and sowing dates. *Indian Journal of Agronomy*, 2000, **47** (3):417-421.
- Panda, B. B., Shivayy, Y. S. and Bandyopadhyay, S. K. Growth and development of Indian mustard (*Brassica juncea* L.) under different varieties of irrigation and date of sowing. *Indian Journal of Plant Physiology*, 2004, **9** (4): 419-425.
- Shukla, D.K. Singh, R.P. and Shukla, A. Response of yellow sarson varieties to different nitrogen levels in Tarai region of U.P. *Annals of Agricultural Research*, 2001, **22** (4): 586-588
- Sihag, J. S. Manohar S.S. and Chaudhary T. Combined effect of sulphur and time of sowing on yield attributes, yield and quality of mustard (*Brassica juncea* L.). *Journal of Eco physiology*, 2003, **6** (1-2): 65-68.
- Singh, R., Patidar, M. and Singh, B. Response of Indian mustard cultivars to different sowing time *Indian Journal of Agronomy*, 2001, **46** (2): 292-295.
- Singh, S. K. and Singh, G. Response of Indian mustard (*Brassica juncea*) varieties nitrogen under varying sowing dates in eastern Uttar Pradesh. *Indian Journal of Agronomy*, 2002, **47** (2):242-248.