

Effects of Potassium on yield of Summer rice (*Oryza sativa*)

ABSTRACT

The effects of potassium on growth and yield parameters of summer rice was assessed through an experiment at the agricultural farm of Uttar Banga Krishi Viswavidyalaya, Pundibari, Coochbehar during the year 2018 and 2019. The important agronomic parameters were significantly influenced with potassium fertilization. Increasing doses of potassium enhanced economic produce of rice to the tune of 6.34 t ha⁻¹ with 150% potassium fertilization (T₅) which was statistically at par with T₄, viz., 125% of the Recommended Dose (RD) of K and T₈ (Nutrient Expert based potassium recommendation) treatment. The Straw yield (11.77 t ha⁻¹) was also enhanced with increasing K levels @150% of the RD of potassium fertilization under same levels of nitrogen and phosphorus. It was concluded that, the current dose of potassium for rice has to be enhanced for desired yield and to keep balance of K⁺ in soil.

Key words: Fertilizer, Nutrient Expert Software, Potassium, Rice, Yield

1. INTRODUCTION

Potassium (K) has been the “forgotten” nutrient in terms of quality of soil-environment during the last few decades, receiving less attention than nitrogen and phosphorus [1]. For food security, a reducing effect of potassium balance is tremendously difficult in terms of food security in global scales [2]. Potassium depletion in soil leads to reduction in yield and use efficiency of nutrients under high density cropping systems [3]. Although, the application of nitrogen in balanced proportion with phosphate and potash in soil is essential [4]. The crop establishment is restricted by insufficient amount of K [5]. Potassium is essential in enzymatic activities, energy metabolism, synthesis of protein and solute transport. The cell turgor, especially in rapidly

expanding cells is regulated by K-ions [6]. The K-nutrition in optimum level enhances the yield attributes of rice and improves strength to the plants which facilitate stand firm against strong winds and reduce lodging. Overall quality of the cereal crops is greatly affected by the application of potassic fertilizer [7].

Being one of the important cereal crops in the world, rice takes on about 90% of the global rice area and production is represented by Asian countries [8] where productivity in India is 3632.9 kg ha⁻¹ [9]. In India, rice production with 22.45 million tonnes in West Bengal, followed by Punjab (20.07 million tonnes) and Uttar Pradesh (19.91 million tonnes) have been reported [10].

Potassium in rice plants are absorbed for proper function of various activities [11]. Farmers often ignore applying K-fertilizer, particularly in Asian subcontinent compared to N and P containing fertilizer [12]. Yield of rice is enhanced significantly by K fertilization ranging from 78 to 93 kg ha⁻¹ [13]. Organic matter application would be an inevitable practice for an alternative supply of nutrients to the crops especially under constraints in resources [14]; although, may not be supplemented by the sole application of organic matter at the present situation. Hence, an integrated nutrient management approach is effective way to enhance productivity of crop [15]. Hence, rice plants become susceptible to biotic and abiotic stresses due to K deficiency *vis-à-vis* applied nutrients [16], although, negative K-balances was observed [17]. The lesser PBC^K (K buffering capacity) of soils of Pundibari (Coochbehar) reflecting that K-fertilization is needed more frequently for higher production of crop. Without the use of K-fertilizer, the lower K status of subsurface soils may not sustain agricultural production system in future [18].

Based on the above perspectives, the present experiments were conducted to assess the effect of potassium on production potential of rice.

2. MATERIALS AND METHODS

The experiment in the field was carried out on rice in a randomized block design (RBD) with three replications at the farm of Uttar Banga Krishi Viswavidyalaya, Coochbehar during two consecutive years, viz., 2018 and 2019 during Summer season. It consisted of eight treatment combinations which are as follows: T₁- Farmer's Practice (120:60:60); T₂- RD (Recommended dose) of NPK (140:70:70); T₃- NP (RD) + K₀ (control); T₄- NP (RD) +125% K; T₅- NP (RD) +150% K; T₆- NP (RD) + vermicompost (5 t ha⁻¹); T₇- NP (RD) + crop residue (rice straw) (5 t ha⁻¹); T₈- NP (RD) + Nutrient expert software based Potassium recommendation (80 kg ha⁻¹).

Gotra Bidhan -1, a popular high yielding variety of rice was used in the experiment with a spacing of 20 cm × 15 cm with 2-3 seedlings hill⁻¹ in 5 m × 4 m plot size. Each plot received a recommended dose of nitrogen and phosphorus as per the state recommendation except the plot where farmer's practice was employed. Split application of urea in three equal splits, viz., basal, first top dressing [21 Days after Transplanting (DAT)] and second top dressing (42 DAT) was applied. The Phosphorus as single super phosphate (SSP) was applied after puddling. Muriate of Potash (MOP) was applied as a source of potassic fertilizer in two splits, 3/4th as basal and rest 1/4th in 42 DAT. All recommended agronomic practices including plant protection measures for rice were followed during the length of growing period of the crop. Plant samples were collected randomly from each net plot area to determine growth and yield parameters at harvest. Partial factor productivity and Agronomic potassium use efficiency were calculated as follows:

Partial Factor Productivity of potassium (PFP_K) = Grain yield in kg/Amount of K applied in kg -----1

Agronomic Potassium Use Efficiency (AKUE) = (Grain yield in fertilized plot - Grain yield in control plot)/Amount of K applied -----2

The statistical analysis was done by employing SPSS software (version 26). The data were analyzed by using standard analysis of variance procedures.

3. RESULTS AND DISCUSSION

It was observed that, in general potassium application had no significant effect on the plant height of rice (Table 1). The plant height (Table 1) was recorded maximum (105.53 cm) in T₄ (125% of the RD of K) and lowest (93.97 cm) in T₃ qualifying the findings of Islam *et al.* (2016) [13].

Different doses of potassium fertilization had influenced on number of tillers (Table 1) and panicle per m² (Table 1) at maturity. Number of tillers was significantly influenced by various K-levels. Average number of tillers m⁻² varied from 216 to 311, the highest being recorded under T₅ (150% of the RD of K). The agronomic parameters were enhanced by higher rate of potassium fertilization. Islam *et al.*, (2015) [19] suggested that number of tillers and number of panicles were increased with application of K beyond 80 kg ha⁻¹. Bagheri *et al.*, (2011) [20] also mentioned that, effective tillers per plant were affected by rate of potassium. Average number of panicle m⁻² (Table 2) varied from 190 (Control) to 262 (150% RD of K)

. Table 1: Effect of treatments on growth and yield contributing parameters of rice

Treatment	Plant height (cm)			No. tillers m ⁻²			No. of panicle m ⁻²			Panicle length (cm)			No. of grains panicle ⁻¹		
	1 st	2 nd	Mean	1 st	2 nd	Mean	1 st	2 nd	Mean	1 st	2 nd	Mean	1 st	2 nd	Mean
	year	year		year	year		year	year		year	year		year	year	
T ₁	102.00 ^a	100.27 ^{ab}	101.13 ^a	249 ^{cde}	253 ^{cd}	251 ^d	207 ^{de}	220 ^{bc}	214 ^{de}	17.80 ^{cd}	19.03 ^c	18.41 ^{cd}	108 ^{cd}	116 ^{cd}	112 ^{bc}
T ₂	103.67 ^a	102.05 ^{ab}	102.86 ^a	256 ^{bcd}	272 ^{bc}	264 ^{cd}	221 ^{cd}	236 ^{abc}	229 ^{cd}	19.50 ^{bcd}	20.11 ^{bc}	19.81 ^{bc}	125 ^{bc}	122 ^{bc}	123 ^b
T ₃	96.67 ^a	91.27 ^b	93.97 ^b	215 ^f	217 ^e	216 ^e	193 ^e	187 ^d	190 ^f	17.27 ^d	16.37 ^e	16.82 ^d	94 ^d	94 ^e	94 ^d
T ₄	105.33 ^a	105.73 ^a	105.53 ^a	278 ^a	305 ^a	291 ^b	245 ^{ab}	256 ^a	250 ^{ab}	22.57 ^{ab}	20.88 ^{ab}	21.73 ^{ab}	138 ^{ab}	140 ^{ab}	139 ^a
T ₅	106.00 ^a	104.75 ^a	105.37 ^a	309 ^a	312 ^a	311 ^a	263 ^a	260 ^a	262 ^a	24.03 ^a	21.42 ^a	22.73 ^a	144 ^a	149 ^a	146 ^a
T ₆	101.33 ^a	97.06 ^{ab}	99.20 ^{ab}	226 ^{ef}	236 ^{de}	231 ^e	196 ^e	209 ^{cd}	203 ^{ef}	18.93 ^{bcd}	17.82 ^d	18.38 ^{cd}	103 ^d	101 ^{de}	102 ^{cd}
T ₇	103.67 ^a	101.20 ^{ab}	102.43 ^a	239 ^{def}	259 ^{bcd}	249 ^d	217 ^{cd}	226 ^{bc}	222 ^{cd}	17.90 ^{cd}	19.39 ^c	18.64 ^{cd}	113 ^{cd}	134 ^{ab}	124 ^b
T ₈	105.00 ^a	105.01 ^a	105.00 ^a	266 ^{bc}	286 ^{ab}	276 ^{bc}	229 ^{bc}	242 ^{ab}	236 ^{bc}	21.42 ^{abc}	19.89 ^{bc}	20.65 ^b	133 ^{ab}	141 ^a	137 ^a
SEM(±)	2.783	3.321	2.166	7.634	8.947	5.880	6.337	8.282	5.214	1.215	0.359	0.634	5.741	5.746	4.061
LSD	NS	NS	6.275	23.155	27.138	17.034	19.221	25.121	15.104	3.685	1.089	1.837	17.414	17.429	11.764

($p=0.05$)

Note: NS: Non significant

Values with small letters indicate differences at 5% level of significance

UNDER PEER REVIEW

The grains per panicle (Table 2) was influenced by potassium fertilization which varied from 94 in control plot to 146 in T₅ (150% of the RD of K) plot and no significant effect between T₅ (150% of the RD of K), T₄ (125% of the RD of K) and T₈ (Nutrient expert software based Potassium recommendation which was 80 kg ha⁻¹) was observed. Where, T₂ and T₇ were also statistically at par. Bahmaniar *et al.*, (2007) [21] found that, application of potassium in field has significantly affected the number of grains per panicle. Zaman *et al.*, (2015) [22] also documented that K fertilization in addition to nitrogenous fertilizer increased yield of rice due to improved grain quality and enhanced the grains per panicle.

Length of panicle was improved by the application of K and their pooled data reflected that the panicle length ranged from 16.82 cm (T₃ - control) to 22.73 cm (T₅ - 150% of the RD of K). Similar result was obtained by Fageria (2015) [23]. The test weight (Table 1) was recorded maximum (24.04 g) in T₅ (125% of the RD of K) and lowest (23.48 g) in T₃ (control). Addition of potassium did not affect the test weight significantly but was certainly improved with increasing rate of potassium to some extent. Similar result was also observed by Islam *et al.* (2015) [19]. The sufficient amount of mineral nutrition, including K fertilization improved number of panicle, length of panicle and 1000-grain weight [24].

Table 2: Effect of treatments on economic produces of rice


Treatment	Grain yield (t ha ⁻¹)			Straw yield (t ha ⁻¹)			Harvest Index (%)			1000-grain weight (g)		
	1 st year	2 nd year	Mean	1 st year	2 nd year	Mean	1 st year	2 nd year	Mean	1 st year	2 nd year	Mean
T1	5.35	5.23	5.29	10.0	10.1	10.0	34.7	34.00	34.4	23.6	23.7	23.7
	bc	c	c	2 ^c	6 ^{cd}	9 ^d	9 ^a	abc	0 ^{ab}	9 ^a	4 ^a	1 ^{ab}

T2	5.60	5.81	5.70	10.8	10.5	10.6	34.1	35.47	34.8	23.8	23.8	23.8
	b	b	b	1 ^{bc}	7 ^{bc}	9 ^c	4 ^a	a	1 ^{ab}	5 ^a	7 ^a	6 ^{ab}
T3	4.09	4.28	4.19	7.79 ^e	8.91 ^e	8.35 ^g	34.4	32.46	33.4	23.4	23.5	23.4
	e	e	e				4 ^a	c	5 ^b	2 ^a	3 ^a	8 ^b
T4	6.13	6.12	6.13	11.3	11.2	11.2	35.1	35.32	35.2	24.0	23.9	24.0
	a	ab	a	4 ^{ab}	1 ^{ab}	7 ^{ab}	1 ^a	a	2 ^a	9 ^a	5 ^a	2 ^{ab}
T5	6.31	6.37	6.34	11.7	11.8	11.7	35.0	34.99	35.0	23.9	24.1	24.0
	a	a	a	1 ^a	4 ^a	7 ^a	4 ^a	ab	2 ^{ab}	6 ^a	2 ^a	4 ^a
T6	4.59	4.78	4.68	8.24 ^d	9.67 ^d	8.96 ^f	35.7	33.06	34.4	23.5	23.5	23.5
	d	d	d	e			5 ^a	bc	1 ^{ab}	5 ^a	7 ^a	6 ^{ab}
T7	5.13	5.41	5.27	9.05 ^d	9.99 ^c	9.52 ^e	36.2	35.11	35.6	23.6	23.7	23.7
	c	c	c		d		2 ^a	a	7 ^a	6 ^a	9 ^a	3 ^{ab}
T8	6.04	6.24	6.14	11.0	11.1	11.0	35.3	35.95	35.6	23.9	23.9	23.9
	a	a	a	5 ^{ab}	1 ^{ab}	8 ^{bc}	5 ^a	a	5 ^a	4 ^a	8 ^a	6 ^{ab}
SEM(±	0.14	0.10	0.08	0.27	0.23	0.17	0.86		0.53	0.21	0.20	0.14
)	3	7	9	0	1	8	6	0.616	1	9	0	8
LSD												
(p=0.05	0.43	0.32	0.25	0.81	0.70	0.51						
)	4	5	8	9	1	6	NS	1.868	NS	NS	NS	NS

Note: NS: Non significant

Values with small letters indicate differences at 5% level of significance

Figure 1: Effects of treatments on grain yield of rice. The error bar indicates the standard deviation at 5% level of significance



The grain yield of rice was observed (Figure 1) where, variation in yield corresponding to each treatment combination was noted. The grain yield (Table 3) of rice varied from 4.19 to 6.34 t ha⁻¹ in T₅ (150% of the RD of K). Maximum grain yield (6.34 t ha⁻¹) obtained under application of 150% of the RD of K (T₅) in soil was statistically at par with T₄ (125% of the RD of K) and T₈ (Nutrient expert software based potassium recommendation which was 141: 44: 80 kg ha⁻¹). The yield attributes of rice are heavily influenced with higher doses of K results in higher yield of rice crop [19]. Nath and Purkayastha (1988) [25] observed that increasing doses of potassium when applied to soils, the grain yield could increase significantly on application of 80 kg K₂O ha⁻¹, which was at par with that of 120 kg K₂O ha⁻¹. The treatments T₁ (Farmer's practice- 120: 60: 60 kg ha⁻¹) and T₇ [110% of the RD of K replaced by crop residue (rice straw)] for grain yield of rice were also statistically at par. Application of rice straw could build up a soil conditions favorable for improving establishment of rice [26]. Singh *et al.*, (2001) [27] also reported that incorporation of straw could increase the soil fertility and enhance crop yield.

The yield of straw was found to be increased with increasing rate of potassium. Straw yield (Table 3) was recorded highest (11.77 t ha⁻¹) in T₅ (150% of the RD of K). Enhanced potassium rate facilitates production of starch and also translocates the photosynthets efficiently to the spikelets [28]. Hence, K induced the grain as well as the straw of rice to obtain greater volume and weight [19]. In recent times, rice residues have been also emerged as an alternative source of energy which reduces production of CO₂ [29]. Increase in Harvest Index results in enhancing grain yield which indicates more partitioning of assimilates to grain and/or total biomass production [30]. It was noted that, the harvest index had negligible influences on the added doses of K (Table 2). Highest harvest index (35.61%) was obtained at T₈ and lowest (31.77%) was at control. It was observed that harvest index was decreased in highest potassium receiving plot and tends to increase with decreasing rate of potassium .This might be due to production of more straw than grain with application of higher doses of K. Bagheri *et al.*, (2011) [20] observed that, higher doses of potassium application reduced the harvest index and enhanced the biological yield. Similar result was obtained by Islam *et al.*, (2015) [19].

Table 3: Effect of treatments on Partial factor productivity and Agronomic potassium use efficiency of rice

Treatment	Partial factor productivity (kg kg ⁻¹)		Agronomic potassium use efficiency (kg kg ⁻¹)	
	1 st year	2 nd year	1 st year	2 nd year
T1	89.11	87.22	20.89	15.89
T2	80.00	82.95	21.52	21.81
T4	70.10	69.94	23.31	21.03
T5	60.06	60.67	21.08	19.90

T6	119.13	124.07	12.81	12.90
T7	66.67	70.22	13.51	14.63
T8	75.54	77.96	24.38	24.46

Partial Factor Productivity of K (PFP_K) [total grain yield (kg) per kg applied K] varied from 59 (T_6) to 89 (T_1) in 2018 and 60 (T_5) to 87 (T_1) in 2019 (Table 3). With increasing rate of potassium, there was gradual decline in partial factor productivity of K except T_7 treatment where crop residue was applied. Crop residue (especially rice straw) contain higher amount of potassium which causes low PFP_K value. On another side, marginal increase in potassium application did not result in much yield increment and hence, the PFP_K value of T_1 was greater than even T_2 . Though in T_6 treatment PFP_K was higher but due to low potassium content in vermicompost, appreciable yield was not achieved. The highest Agronomic Potassium Use Efficiency (AKUE) was observed in both years with T_8 treatment (24.38 in 2018 and 24.46 in 2019) (Table 3) followed by T_4 treatment as yield increment over control in these treatments were much higher; the lowest was observed in T_6 treatment (12.81 in 2018 and 12.90 in 2019) (Table 3). On treatment T_6 and T_7 , the yield increment over control was very less and resulting the lower Agronomic Potassium Use Efficiency (AKUE). Nutrient Expert-based potassium recommendation which was based on SSNM (Site Specific Nutrient Management) performed best. It was observed from this experiment, that, the crop did not respond well beyond 80 kg K_2O ha^{-1} .

4. Conclusion

On the basis of the 2-years experiment, it can be concluded that, application of potassium obviously increased the yield of summer rice. The results indicated that 125% of the RD of K

gave a better yield of rice, being at par with 150% of the RD of K. The application of potassium through Nutrient Expert gave best result with respect to growth and yield which was at par with 150% of the RD of K and 125% of the RD of K. Rice straw incorporation in soil at harvest might be an alternative source of potassium build-up in soil.

REFERENCES

1. Romheld V, Kirkby EA. Research on potassium in agriculture: needs and prospects. *Plant Soil*. 2010;335(1):155–180.
2. Cakmak I. Potassium for better crop production and quality. *Plant Soil*. 2010;335:1–2.
3. Regmi A, Ladha J, Pasuquin E, Pathak H, Hobbs P, Shrestha, L, Gharti D, Duveiller E. The role of potassium in sustaining yields in a long-term rice-wheat experiment in the Indo-Gangetic Plains of Nepal. *Biol. Fertil. Soils*. 2002;36(3):240–247.
4. Roy RN, Braun H. (1985). Systems approach to nutrient management with special emphasis to nitrogen. *Nitrogen management in farming systems in humid and subhumid tropics*. 1985;339-348.
5. Rengel Z, Damon PM. Crops and genotypes differ in efficiency of potassium uptake and use. *Physiol. Plant*. 2008;133(4):624-636.
6. White PJ, Karley AJ. 2010. Potassium. In *Cell biology of metals and nutrients*. Springer, Berlin, Heidelberg. pp. 199-224.
7. Pavithira E, Sirisena DN, Herath HMS. Effect of potassium fertilizer split applications together with straw on optimum level in leaf and stem of Rice. . *J. Agric. Sci*. 2017;12(1):24- 33.
8. FAO STAT online database. January 2013.

9. FAO. FAO Statistical data, 2017. [http: stat. www.faostat.org.in](http://stat. www.faostat.org.in).
10. Pathak H, Tripathi R, Jambhulkar NN, Bisen JP, Panda BB. (2020). Eco-regional Rice Farming for Enhancing Productivity, Profitability and Sustainability. NRRI Research Bulletin No. 22, ICAR-National Rice Research Institute, Cuttack 753006, Odisha, India. pp 28.
11. Zhang F, Niu J, Zhang W, Chen X, Li C, Yuan, L. and Xie, J. Potassium nutrition of crops under varied regimes of nitrogen supply. *Plant soil*. 2010;335(1): 21-34.
12. Li J, Lu J, Li X, Ren T, Cong R, Zhou L. Dynamics of potassium release and adsorption on rice straw residue. *PLoS One*. 2014;9(2): e90440.
13. Islam A, Muttaleb A. Effect of potassium fertilization on yield and potassium nutrition of Boro rice in a wetland ecosystem of Bangladesh. *Arch. Agron. Soil Sci*. 2016;62(11):1530-1540.
14. Palm CA, Myers RJ, Nandwa SM. Combined use of organic and inorganic nutrient sources for soil fertility maintenance and replenishment. *Replenishing soil fertility in Africa*. 1997;51:193-217.
15. Bandara, W. M. J., Wickramasinghe, W. M. A. D. B., Sirisena, D. N. and Wijewardana, J. D. H. Response of rice to applied potassium in soils of Sri Lanka. Use of phosphorus and potassium fertilizers in SriLankanagriculture.(Kumaragamage, D., Satyanarayana, T., Singh, H. and Majumdar, K. Eds.). International Plant Nutrition Institute (IPNI), Gurgaon, Haryana, India, 89-117;2009.
16. Johnston AE, Poulton PR and Syers JK. Phosphorus, potassium and sulphur cycles in agricultural soils. *Proceedings No. 465*. York (UK): The International Fertiliser Society; p. 44;2001.

17. Miah MM, Saha PK, Islam A, Hasan MN, Nosov V. Potassium fertilization in rice-rice and rice-wheat cropping system in Bangladesh. *Bangladesh J. Agric. and Environ.* 2008;4:51-67.
18. Singh NK, Banik GC, Ghosh A. Spatial Variation of Potassium Quantity/Intensity Relationships and Buffering Capacity of Some Entisols of the Himalayan Floodplain. *J. Indian Soc. Soil Sci.* 2019;67(3):321-328.
19. Islam A, Chandrabiswas J, Sirajul Karim AJM, Salmapervin MST, Saleque MD. Effects of potassium fertilization on growth and yield of wetland rice in grey terrace soils of Bangladesh. *Research on Crop Ecophysiology.* 2015;10(2):64-82.
20. Bagheri R, Mobasser HR, Malidarreh AG, Dastan S. Effect of seedling age and potassium rates on morphological traits related-lodging, yield and yield components of rice (*Oryza sativa L.*) in Iran. *American-Eurasian Journal of Agricultural & Environmental Sciences.* 2011;11(2):261-268.
21. Bahmaniar MA, Ranjbar GA. Response of Rice Cultivars to Rates of Nitrogen and Potassium Application in. *Pak. J. Biol. Sci.* 2007;10(9):1430-1437.
22. Zaman U, Ahmad Z, Farooq M, Saeed S, Ahmad M, Wakeel A. Potassium fertilization may improve stem strength and yield of Basmati rice grown on nitrogen-fertilized soils. *Pak. J. Agric. Sci.* 2015; 52(2):437-443
23. Fageria NK. Potassium requirements of lowland rice. *Commun Soil Sci Plant Anal.* 2015;46(12):1459-1472.
24. Fageria NK, Baligar VC, Jones CA. Growth and mineral nutrition of field crops, 3rd ed. Boca Raton, FL: CRC;2011

25. Nath AK, Purkaystha S. A study on soil test and crop response in respect of potassium in acid alluvial soils of Assam. *J. Indian Soc. Soil Sci.* 1988;36(1):120-124.
26. Sarkar MIU, Islam MN, Jahan A, Islam A, Biswas JC. Rice straw as a source of potassium for wetland rice cultivation. *Geol. ecol. landsc.* 2017;1(3):184-189.
27. Singh M, Singh VP, Reddy KS. Effect of integrated use of fertilizer nitrogen and farmyard manure or green manure on transformation of N, K and S and productivity of rice-wheat system on a Vertisol. *J. Indian Soc. Soil Sci.* 2001;49(3): 430-435.
28. Beringer H. Functions of Potassium in Plant Metabolism with Particular Reference to Yield. In: Sekhon GS (ed). *Potassium in Soils and Crops*, Potash Research Institute of India, 1978. New Delhi, India, pp: 185-202;1978.
29. Matsumura Y, Minowa T, Yamamoto H. Amount, availability, and potential use of rice straw (agricultural residue) biomass as an energy resource in Japan. *Biomass Bioenergy.* 2005;29: 347-354.
30. Richards RA.. Selectable traits to increase crop photosynthesis and yield of grain crops. *J. Exp. Bot.* 2000;51:447-458.