

Evaluating the suitability of System of Rice Intensification practices for enhancing rice and water productivity in semi-arid environment, Tamil Nadu, India

Abstract

In the world of rapid change in climate, irregular rainfall pattern tends to pose serious impact on water availability for agriculture. Rice is one of the important food crops to get affected much by the less water availability because of its higher water requirement. Various techniques were adopted to meet low water availability or increase productivity at the expense of either. System of Rice Intensification (SRI) is being tried by many countries with several modifications based on their priorities with the aim of enhancing productivity besides reducing the water demand for rice cultivation. It is essential to have more insight into the individual and compounding effect of multiple components of SRI on yield, and water productivity of rice for identifying the potential and suitable SRI practices. Investigating the influence of different practices of SRI viz., planting of young and single seedlings per hill in wider spacing, water saving irrigation like Alternate Wetting and Drying (AWD), and weed control using cono-weeders on rice using the data obtained from the field experiment carried out during 2021 in Tamil Nadu Agricultural University, Coimbatore, India. Water productivity of rice plants under SRI were compared with conventional practices. The results revealed that plants grown with complete SRI practices had increased water productivity by 0.25 kg grain/m³ of water which is almost twice that of conventional cultivation system. The yield obtained in SRI treatments was higher about 39 % than conventional treatments. The total water savings was 20 % higher in AWD treatments than continuous flooding treatments.

Keywords: Water productivity, SRI, Alternate Wetting and Drying, Cono-weeding, rice

1. Introduction

India's food requirement is majorly dependent on rice as staple food for the past few decades feeding about half of the country's population. The per capita consumption of milled rice is around 74 kg per year in India (Maclean *et al.*, 2002). Rice being a semi aquatic crop, is cultivated over 43.5 million hectares in India with productivity of 2722 kg/ha during 2020 (Indiastat, 2022). Despite of its importance as food crop, water requirement of rice cultivation is about 80 percent of total irrigation water resources in Asia (Maclean *et al.*, 2002; Tuong *et al.*, 2015). Rice consumes on average of about 1300 to 1500 mm of water (IRRI, 2022) which is very high compared to other major food crops. World is facing water scarcity due to climate change which is expected to worsen in the future, putting water and food security at risk. There is a need for improved crop management practices to increase productivity of rice without depending on more water. System of Rice Intensification (SRI) is one such approach that promises an increase in yields with low external inputs and at the same time reduces water consumption (Uphoff *et al.*, 2002). This system is developed through farmers' participatory experiments conducted by a French missionary named Henri de Laulanie in Madagascar in 1980s-90s (Dobermann, 2004; Nirmala *et al.*, 2021).

SRI principles generally include good nursery management, careful transplanting of young seedlings, single seedlings at a wider spacing in a square pattern, intermittent irrigation to avoid continuous flooding except at the flowering stage, use of mechanical weeding without using herbicides, and improved nutrient management mostly with organic manures. Nevertheless, the SRI practices are not a set of the standardized package of practices but an empirical set that depends on location-specific conditions (Uphoff *et al.*, 2002).

SRI is known to increase productivity, while reducing water use through a practice known as Alternate Wetting and Drying (AWD) irrigation. There are reports of water savings from 20% to 50% savings with a small or no reduction in yield while some reported slight increase in yield under AWD irrigation system (Zhao *et al.*, 2010; Howell *et al.*, 2015; Shaibu *et al.*, 2015; Mondal, 2017). Water productivity is also increased largely through SRI practices. However, there only a few studies available to explain the combined effect of each SRI practices on yield and water productivity of rice compared to conventional practices. Therefore, the current study aims to fathom out the impacts of alternate wetting and drying irrigation and its influence on yield and water productivity of rice when combined with other principles of SRI.

2. Materials and Methods

Experimental site details: The field experiment was undertaken during the summer of 2021 at wetlands of Tamilnadu Agricultural University, Coimbatore, India. The soil in the study area was clay loam with slightly basic pH of 8.2, with 225:54:290 kg/ha of N: P: K and about 0.56 percent of available carbon. The site recorded an average maximum temperature of 34 °C and a minimum of 23.9 °C, with a cumulative precipitation of 121 mm. The average bright sunshine hours were 7.9 hours, average daily evaporation was 6.8 mm d⁻¹ and 14.9 MJ m⁻² solar radiation was observed during cropping period.

Experimental design and treatments: Each practice of SRI was added to conventional cultivation to create different treatment combinations for observing their influence on rice yield and water productivity as given in the Table 1. Randomized block design with four replications was used for this experiment.

Table 1. Treatment details of the experiment

| | |
|----|---|
| T1 | Square planting (25 cm x 25 cm) of 2 leaf stage single seedling |
| T2 | T1 + Cono-weeding four times on 15, 25, 35 and 45 days after planting |
| T3 | T1 + Alternate wetting and Drying (AWD) method of irrigation |
| T4 | T2 + Alternate wetting and Drying (AWD) method of irrigation (complete SRI) |
| T5 | 21 days old seedling @ 2–3 seedlings/hill -spacing 15 cm x 10 cm (conventional) |
| T6 | T5 + Alternate wetting and Drying (AWD) method of irrigation |

Field preparation and crop management: Nursery was raised with 4 kg of seeds in 10 m² area where 1 kg of DAP was applied as basal fertilizer. The main field preparation involved trimming and plastering, followed by puddling of the field. The plots were laid out with buffer zones to prevent water flow across the plots. Pre-emergence herbicide butachlor was applied at a rate of 1.25 kg ha⁻¹ broadcasted after mixing with sand. Recommended dose of fertilizers (150:50:50 N: P: K) was applied to all treatments (Directorate of Agriculture & Tamil Nadu Agricultural University, 2020).

2.1 Irrigation and weeding

For AWD treatments, irrigation was done to fill up to a thin water layer once the water level reaches 10 cm below soil layer. The irrigation level was monitored with the help of field water tube (Cabangon *et al.*, 2009) installed in the plots as shown in the Fig 1 and Fig 2. A practice named “safe-AWD” given by Bouman *et al.* (2007) was adopted where the field was irrigated sufficiently during the week of peak flowering to prevent yield losses due to water stress. Cono weeding was done four times starting from 10 days after transplanting at ten days interval in a criss-cross pattern. Manual weeding also was carried out once in 10 days. The buffer and irrigation channels were weeded manually to keep weed growth low.



Fig 1. Field water tube installed in AWD plots to monitor water depth




Fig 2. Alternate Wetting and Drying irrigation management using field water tube

2.2 Harvesting Index

Harvest index (HI) was calculated as ratio between economical yield (grain weight) to biological yield (grain weight + stover weight) expressed as percentage, as given by the following equation (1)

$$HI = \frac{\text{Economical yield}}{\text{Biological yield}} \times 100 \quad (1)$$

2.3 Water productivity

Water productivity is the ratio of total yield of the crop to the total amount of water supplied to the crop including irrigation and rainfall during the cropping period (Molden *et al.*, 2001). It can be expressed by the equation (2) given below

$$WP = \frac{\text{grain yield}}{\text{total water supplied}} \quad (2)$$

2.4 Statistical analysis

The observed data were statistically analysed for significance using analysis of variance (ANOVA) as per randomized complete block design. The mean dataset was then arranged in descending order and grouped or differentiated using Duncan multiple range test (DMRT) with 5% significance level and the results are presented in Table 2.

3. Results and Discussion

3.1 Yield

Yield variation obtained from the treatments with SRI as well as conventional cultivation practices is presented in the Table 2. Yield was higher in the SRI based cultivation rather than conventional method. Complete SRI cultivation practice (T₄) registered the maximum yield of 6,499 kg/ha which was followed (5,925 kg/ha) by square planting with cono weeding treatment (T₂). Conventional cultivational practices registered the lower yield as compared to SRI methods. Complete conventional practices (T₅) showed the lowest yield of 3,919 kg/ha which was on par (4,136 kg/ha) with conventional method with altered wetting and drying treatment (T₆). There are multiple experimental reports of increased yields in SRI treatments (Zhao *et al.*, 2010; Barison and Uphoff, 2011; Thakur *et al.*, 2020) and also farmers surveys indicating yield gains through SRI (Senthilkumar *et al.*, 2008; Nirmala *et al.*, 2021).

3.2 Harvest Index

Ratio of grain yield to total dry matter produced gives the harvest index (HI) which was considered as reproductive efficiency of the crop. It was influenced by interaction between genotype, atmospheric condition and crop cultivation practices. Complete SRI practices gave the maximum HI of 47.05%. The conventional methods, T₅ and T₆ produced the least HI of 37.27% and 39.81% respectively. Chapagain *et al.* (2011) found that the harvest index varied among cultural practices wherein SRI cultivation produced higher HI than conventional methods.

Table 2. Effect of SRI practices on yield, harvest index and water productivity of Rice

| Treatment | Yield (kg/ha) | Harvest Index (%) | Water productivity (kg grain/cu. m water) |
|-----------|----------------------------|----------------------------|---|
| T1 | 4588 ^d ± 269.88 | 47.16 ^a ± 4.10 | 0.29 ^e ± 0.02 |
| T2 | 5925 ^b ± 278.64 | 44.98 ^{ab} ± 2.39 | 0.37 ^c ± 0.02 |
| T3 | 5525 ^c ± 142.68 | 42.88 ^b ± 0.98 | 0.43 ^b ± 0.01 |
| T4 | 6491 ^a ± 284.34 | 47.05 ^a ± 2.84 | 0.51 ^a ± 0.02 |
| T5 | 3911 ^e ± 192.45 | 37.27 ^c ± 2.05 | 0.25 ^f ± 0.01 |
| T6 | 4136 ^e ± 346.69 | 39.81 ^c ± 3.81 | 0.32 ^d ± 0.03 |

3.3 Water productivity

Water productivity was higher in the SRI based cultivation method compared to conventional cultivation methods. Complete SRI method of cultivation (T₄) produces about 0.51 kg of biomass per cubic meter of water whereas the lowest water productivity was obtained in complete conventional method of cultivation with 0.25 kg per cubic meter of water. The square planting with alternate wetting and drying cropping practice recorded 0.43 kg per cubic meter of water. Thakur *et al.* (2018) found that SRI treatments showed almost twice the increase in water productivity compared to conventional treatments and AWD further enhanced water productivity when combined with SRI.

4. Conclusion

Water productivity of rice has increased twice with alternate wetting and drying than conventional flooding irrigation. It is inferred that drying the soil at a protective level for rice crop without subjecting to moisture stress will not reduce the rice productivity and may even increase yield to some extent. AWD can also increase soil aeration which leads to increased root growth and eventually boosted the productivity of rice crop. AWD in combine with other

practices of SRI, could increase both yield and water productivity in semi-arid regions which experience frequent water shortages leading to reduction in yield or even crop failures.

5. References

- Barison, Joeli, and Norman Uphoff. 2011. "Rice Yield and Its Relation to Root Growth and Nutrient-Use Efficiency under SRI and Conventional Cultivation: An Evaluation in Madagascar." *Paddy and Water Environment* 9(1):65–78.
- Bouman, B. A. M., R. M. Lampayan, and T. P. Tuong. 2007. *Water Management in Irrigated Rice: Coping with Water Scarcity*. Los Banos, Philippines: International Rice Research Institute.
- Cabangon, R., F. G. Corcuera, D. O. Angeles, and R. Lampayan International Rice Research Inst. Los Banos Laguna (Philippines). 2009. "Perched Water Tube: A Simple Tool for Managing Water under Alternate Wetting and Drying (AWD) Irrigation in Rice." P. 86 in *Philippine Journal of Crop Science (Philippines), Philippine Journal of Crop Science (Philippines)*, edited by F. A. O. of the UN.
- Chapagain, Tejendra, Andrew Riseman, and Eiji Yamaji. 2011. "Assessment of System of Rice Intensification (SRI) and Conventional Practices under Organic and Inorganic Management in Japan." *Rice Science* 18(4):311–20.
- Dobermann, A. 2004. "A Critical Assessment of the System of Rice Intensification (SRI)." *Agricultural Systems* 79:261–81.
- Howell, Katharine R., Pitambar Shrestha, and Ian C. Dodd. 2015. "Alternate Wetting and Drying Irrigation Maintained Rice Yields despite Half the Irrigation Volume, but Is Currently Unlikely to Be Adopted by Smallholder Lowland Rice Farmers in Nepal." *Food and Energy Security* 4(2):144–57.
- Indiastat. 2022. "Statewise and Seasonwise Rice Area, Production and Productivity." *Ministry of Agriculture & Farmers Welfare, Govt. of India. (ON2873) & Past Issues*. Retrieved March 3, 2022 (<https://www.indiastat.com/data/agriculture/rice/data-year/2012-2022>).
- IRRI. 2022. "Water Management - IRRI Rice Knowledge Bank." 1. Retrieved April 28, 2022 (<http://www.knowledgebank.irri.org/step-by-step-production/growth/water-management#for-safe-alternate-wetting-and-drying>).
- Maclean, J. L., D. C. Dawe, B. Hardy, and G. P. Hettel. 2002. "Source Book for the Most

Important Economic Activity on Earth W A R D A A D R A O 2002 Third Edition.”

- Molden, D., R. Sakthivadivel, and Z. Habib. 2001. *Basin-Level Use and Productivity of Water: Examples from South Asia*.
- Mondal, Dibyendu. 2017. “Prospects of Alternate Wetting and Drying (AWD) Methodology of Irrigation through System Intensification on Productivity of Summer Transplanted Rice (*Oryza Sativa* L.)” *International Journal of Pure & Applied Bioscience* 5(4):629–34.
- Nirmala, B., M. D. Tuti, R. Mahender Kumar, Amtul Waris, P. Muthuraman, Brajendra Parmar, and T. Vidhan Singh. 2021. “Integrated Assessment of System of Rice Intensification vs. Conventional Method of Transplanting for Economic Benefit, Energy Efficiency and Lower Global Warming Potential in India.” *Agroecology and Sustainable Food Systems* 45(5):745–66.
- Senthilkumar, K., P. S. Bindraban, T. M. Thiyagarajan, N. De Ridder, and K. E. Giller. 2008. “Modified Rice Cultivation in Tamil Nadu, India: Yield Gains and Farmers’ (Lack of) Acceptance.”
- Shaibu, Y. A., H. R. Mloza Banda, C. N. Makwiza, and J. Chidanti Malunga. 2015. “Grain Yield Performance of Upland and Lowland Rice Varieties Under Water Saving Irrigation Through Alternate Wetting and Drying in Sandy Clay Loams of Southern Malawi.” *Experimental Agriculture* 51(2):313–26.
- Thakur, A. K., K. G. Mandal, Rajeeb K. Mohanty, and S. K. Ambast. 2018. “Rice Root Growth, Photosynthesis, Yield and Water Productivity Improvements through Modifying Cultivation Practices and Water Management.” *Agricultural Water Management*.
- Thakur, Amod K., Krishna G. Mandal, and Sachidulal Raychaudhuri. 2020. “Impact of Crop and Nutrient Management on Crop Growth and Yield, Nutrient Uptake and Content in Rice.” *Paddy and Water Environment* 18(1):139–51.
- Tuong, To P., B. A. M. Bouman, and Martin Mortimer. 2015. “More Rice, Less Water—Integrated Approaches for Increasing Water Productivity in Irrigated Rice-Based Systems in Asia.”
[Http://Www.Tandfonline.Com/Action/AuthorSubmission?JournalCode=tpps20&page=i](http://www.tandfonline.com/action/authorSubmission?journalCode=tpps20&page=i)

nstructions 8(3):231–41.

Uphoff, Norman T., Erick C. M. Fernandes, Yuan Longping, Peng Jiming, Sebastien Rafaralahy, and Justin Rabenandrasana. 2002. “Assessments of The System of Rice Intensification (SRI).” Pp. 23–25 in. Sanya, China.

Zhao, Limei, Lianghuan Wu, Yongshan Li, Sarkar Animesh, Defeng Zhu, and Norman Uphoff. 2010. “Comparisons of Yield, Water Use Efficiency, and Soil Microbial Biomass as Affected by the System of Rice Intensification.” *Communications in Soil Science and Plant Analysis* 41(1):1–12.

UNDER PEER REVIEW