

Original Research Article

STUDIES ON CORRELATION AND PATH COEFFICIENT ANALYSIS OF YIELD AND YIELD ATTRIBUTING CHARACTERS IN RICE LANDRACES

(*Oryza sativa* L.)

ABSTRACT

In order to characterise the correlation and path analysis for yield and yield attributing traits for 66 genotypes including 4 checks in Augmented Block Design replicated thrice during Kharif, 2021, the present investigation was conducted at the Agricultural College, Aswaraopet, Bhadradi, Telangana. The most positive and significant correlation between important traits and grain yield per plant was found to be between the number of panicles per plant and the number of filled grains per panicle. The features number of panicles per plant and number of filled grains per panicle had reported the strongest positive direct effect on grain yield per plant, and may be considered as excellent selection criteria for yield improvement, according to path coefficient analysis.

KEY WORDS: *Correlation, Path coefficient analysis, Landrace, Rice.*

INTRODUCTION

India's main food crop is rice (*Oryza sativa* L.). It belongs to the Oryzoideae subfamily of the Gramineae. *Oryza sativa* and *Oryza glaberrima* are two of the 23 species in the genus, according to D. Chatterjee [1], 21 of which are wild. In India, rice is referred to as "Prana," which means "breath of life." 700 million tonnes of rice are produced annually on 158 million hectares of land worldwide. 114 of the 193 countries in the world's

population grow it. But more than 90% of the world's rice is produced and consumed in Asia. More over half of the world's rice-growing acreage and 56% of its rice production are in China and India (Patra) [2]. With an area, production, and productivity of 43.80 million ha, 118.88 million tonnes, and 2650 Kg ha⁻¹, respectively, India is one among the world's top producers of rice. Nearly all of the states in India, including Telangana, Andhra Pradesh, Bihar, Uttar Pradesh, Maharashtra, and West Bengal, cultivate rice. Out of these, West Bengal and Uttar Pradesh generate the most rice. The production and productivity of rice in Telangana State, however, is 60.50 million tonnes and 3550 kg ha⁻¹ on an area of 1.8 million ha. With an increasing area of 55.50 lakh acres in kharif and 54.30 lakh acres in rabi, Telangana is gradually becoming known as the "Rice Bowl of India".

Given that grain yield is a complicated trait, independent variables other than environmental factors are also involved. Characters that are important to breeders tend to be complicated as a result of various factors interacting. Understanding the link between yield and its components is necessary to make the right selection options. By assessing the relative influence of various component attributes on grain yield, the correlation coefficient is utilised to determine which plant is the best. To distinguish between direct and indirect impacts of correlation, path coefficient analysis is utilised (Ekka et al.) [3]. The purpose of the current study was to assess the direct and indirect effects of yield attributing qualities on grain yield and to understand the relationship between yield and yield component traits.

MATERIAL AND METHODS

The material used in the present investigation comprised of 70 genotypes which included 66 rice land races and 04 Standard check varieties (Chittimutyalu, DRR Dhan 45, BPT5204 and Zincorice). The experiment was carried out during *kharif*, 2021, at Agricultural College, Aswaraopet, which is situated at an altitude of 64 m above mean sea

level on 18° 80' N latitude and 7° 55' E longitudes in Northern Zone of Telangana State. The experimental material was planted in a Augmented block design, Standard check varieties replicated into 3 times. Each replication consisted of three rows of 1 m length with a spacing of 20 cm between the rows and 15 cm between the plants. All required precautions were taken to ensure a uniform plant population in each treatment per replication. Five plants were selected at random and observations were recorded from each replication. The characters studied were Days to 50% flowering, Days to maturity, Plant height (cm), Panicle length (cm), Number of panicles per plant, Number of filled grains per panicle, Grain length (mm), Grain width (mm), 1000 grain weight (g), Kernel length (mm), Kernel breadth (mm), Grain iron content (ppm), Grain zinc content (ppm) and Grain yield per plant (g). Relevant data was collected and subjected to statistical analysis by employing the methods Correlation coefficient and Path analysis by Dewey and Lu [4].

RESULTS AND DISCUSSION

Phenotypic selection is influenced by several biotic and abiotic factors, and the selection is always deceptive. Therefore, every crop development programme will benefit from selection based on the set of qualities that significantly correlate with grain yield.

The degree and direction of association between diverse features, which is the key to the selection process to develop high yielding cultivars, will be shown by correlation studies on various yield and yield attributing traits. The fundamental information about the direct and indirect effects of various independent factors on the final dependent variable, grain yield, is provided by path coefficient analysis.

The correlation coefficients between yield and yield components, including plant height, panicle length, number of panicles per plant, number of filled grains per panicle, grain length, grain width, 1000 grain weight, kernel length, kernel breadth, grain iron content, and grain zinc content, were evaluated and the results have been presented in Table 1.

Positive and significant associations were noticed for panicle length, number of panicles per plant, number of filled grains per panicle. Positive but non-significant association was observed with days to maturity, grain length, grain width, 1000 grain weight, kernel length, kernel breadth, grain iron content, grain zinc content and showed negative non significant correlation with days to 50% flowering and plant height (Ratna et al. [5] ; Rahman et al. [6]. The findings revealed that late flowering types may have more chance of vegetative growth, increasing source and sink relationship paving way for more number of grains and yield.

Positive and significant associations were observed for days to maturity with days to 50% flowering, zinc content and non significant positive correlation with the panicle length, number panicles per plant (Devi et al.) [7], number of filled grains per panicle, grain length, 1000 grain weight, kernel length, grain iron content, grain yield per plant. It recorded significant negative association with the plant height, grain width and non significant negative association with the kernel breadth.

Positive and significant associations were observed for days to maturity with days to fifty percent flowering ,iron content and zinc content. While non significant positive correlation with the panicle length, number panicles per plant, number of filled grains per panicle (Immanuel et al.) [8], grain length, 1000 grain weight, kernel length, grain iron content, grain yield per plant. Days to maturity recorded significant negative association

with the plant height, grain width and non significant negative association with the kernel breadth.

Plant height revealed that there was a non significant positive correlation with panicle length (Umarani et al.) [9], number of filled grains per panicle, grain width, 1000 grain weight, kernel breadth. While significant negative association with days to maturity, grain iron content (Gangashetty et al.) [10] and non significant negative association with the plant height, number of panicles per plant, grain length, kernel length, grain zinc content (Dore et al.) [11] and grain yield per plant (Ratna et al. [5] ; Rahman et al.) [6].

Panicle length was significantly and positively correlated with number of panicles per plant (Singh et al.) [12], grain width, kernel length, grain yield per plant (Babaeian and Bagheri) [13] and non significant positive association with the number of filled grains per panicle, grain length, kernel breadth and grain zinc content (Shivani et al.) [14]. The concentrations of days to 50% flowering, 1000 grain weight, grain iron content has a non-significant negative correlation.

Panicle length, Number of filled grains per panicle, grain yield per plant had a strong positive connection with the number of panicles per plant showing that it is one of the important selection attributes for yield increase. The trait recorded that there was a non significant positive correlation with days to 50% flowering, days to maturity, grain width, kernel length, grain iron content and grain zinc content. Plant height, grain length, 1000 grain weight and kernel breadth have negative non-significant association with number of panicles per plant. The results were in conformity with Patel et al. [15], Pandey et al. [16], Seneega et al. [17], Islam et al. [18], Prasannakumari et al. [19], Parimala et al. [20] and Sudeepti et al. [21] for grain yield per plant.

Filled grains per panicle were significantly and positively correlated with number of panicles per plant, grain zinc content, grain yield per plant (Patel et al.) [15]. positive non significant association with days to 50% flowering, days to maturity, plant height, panicle length, grain width, 1000 grain weight, kernel length, kernel breadth and grain iron content and non significant negative correlation with the grain length. Selection based on the number of filled grains per panicle improves yield.

Grain length shows a positive significant correlation with grain width and kernel length and also exhibits a positive non significant association with days to maturity, panicle length, 1000 grain weight, kernel breadth, grain iron content and grain yield per plant. It recorded non significant negative correlation with the days to 50% flowering, plant height, number of panicles per plant, number of filled grains per panicle and grain zinc content. These results are in line with the earlier reports of (Ekka et al. [3]; Islam et al. [18]).

Positive significant association were observed for panicle length, grain length, kernel length, kernel breadth with grain width and positive non significant association with the plant height, number of panicles per plant, number of filled grains per panicle, 1000 grain weight, grain iron content and grain yield per plant. The trait also shows a negative significant association with days to maturity and negative non significant association with the days to 50% flowering and grain zinc content. These results are in line with the earlier reports of Ekka et al. [3] , Mallimar et al. [22] and Islam et al. [18]

1000 grain weight revealed a significant positive correlation with kernel breadth and non significant positive correlation with days to 50% flowering, days to maturity, plant height, number of filled grains per panicle, grain length, grain width, kernel length, grain zinc content, grain iron content and grain yield per plant. This trait had a negative non

significant association with panicle length and number of panicles per plant (Hasan et al. [23]). As a result, simple selection of this attribute could be able to improve yield.

Positive significant correlation were noticed for panicle length, grain length, grain width with kernel length and positive non significant correlation with the days to maturity, number of panicles per plant, number of filled grains per panicle, 1000 grain weight , kernel breadth, grain iron content and grain yield per plant. The trait also reveals a negative non significant correlation with the days to 50% flowering, plant height and grain zinc content. These results are in line with the earlier reports of Ekka et al.[3], Islam et al. [18] and Prasannakumari et al. [19].

Kernel breadth exhibits a positive significant correlation with grain width, 1000 grain weight and positive non significant correlation with plant height, panicle length, number of filled grains per panicle, grain length, kernel length, grain iron content, grain zinc content and grain yield per plant. This trait had a negative non significant correlation with the days to 50% flowering, days to maturity and number of panicles per plant. These results are in line with the earlier reports of Ekka et al. [3], Islam et al. [18] and Prasannakumari et al. [19].

Grain iron concentration has a significant positive correlation ($r=0.248$) with days to 50% flowering and grain zinc content and also exhibits a non-significant positive connection (0.165) with days to maturity, number of panicles per plant, number of filled grains per panicle, grain length, grain width, 1000 grain weight, kernel length, Kernel breadth and grain yield per plant. This trait also shows a negative significant correlation with the plant height and negative non significant correlation with the panicle length. Oliveira et al. [24], Mallimar et al. [22] and Raza et al. [25] all found a positive association between iron and zinc in their studies.

Grain zinc concentration has a significant positive correlation with days to 50% flowering, days to maturity, number of filled grains per panicle and grain iron content and also recorded a non-significant positive connection with panicle length, number of panicles per plant, 1000 grain weight, kernel breadth, grain yield per plant and number of filled grains per panicle. This trait reveals a negative non significant correlation with the plant height, grain length , grain width and kernel length. For iron, the results were consistent with Oliveira et al. [24], Mallimar et al. [22], and Raza et al. [25], and for yield, they were consistent with Singh et al. [12].

PATH COEFFICIENT ANALYSIS

It is generally recognised that simple correlation does not reflect the underlying relationship between qualities and yield, nor does it explain the relationship between causes and effects between the numerous yield parameters and, ultimately, the yield. By separating the correlation coefficients into direct and indirect effects, the path analysis technique gives insight on the true impact of independent factors on yield.

For the yield and yield component traits, path coefficient analysis estimates have been presented in Table 2. From the path analysis it was observed that, number of panicles per plant reported a high direct positive effect on grain yield per plant (0.7353) and the correlation between the two traits is positive and significant (0.7991). It expressed positive indirect effects through days to 50% flowering (0.0465), days to maturity (0.0495), panicle length (0.1762), number of filled grains per panicle (0.4457), grain width (0.1164), kernel length (0.0771), grain iron content (0.1477), grain zinc content (0.1155) and negative

indirect effects through plant height (-0.0783), grain length (-0.0193), 1000 grain weight (-0.0155) and kernel breadth (-0.0066). The results were in agreement with the findings of Edukondalu et al.[26], Archana et al. [27] and Rachana et al. [28].

The trait, panicle length was noticed as the second most important character with a direct positive effect on grain yield per plant (0.1018) and the correlation between the two traits is positive and significant (0.3118). It expressed positive indirect effects through days to maturity (0.0022), plant height (0.0157), number of panicles per plant (0.0244), number of filled grains per panicle (0.0235), grain length (0.0056), grain width (0.0246), kernel length (0.0257), kernel breadth (0.0071), grain zinc content (0.0043) and negative indirect effects through days to 50% flowering (-0.0092), 1000 grain weight (-0.0056) and grain iron content (-0.0065). The results were in agreement with the findings of Islam et al.[18] .

Days to maturity shows a high direct positive effect on grain yield per plant (0.2302) and the correlation between the two traits is positive but non significant (0.1751) due to positive indirect effects through days to 50% flowering (0.1301), panicle length (0.0049), number of panicles per plant (0.0155), number of filled grains per panicle (0.0318), grain length (0.0147), 1000 grain weight (0.0199) kernel length (0.0153), grain iron content (0.0539), grain zinc content (0.0671) and negative indirect effect through plant height (-0.0752), grain width (-0.0555), and kernel breadth (-0.0096). The results were in agreement with the findings of Immanuel et al. [8] and Babu et al. [29].

Grain width reported a direct positive effect on grain yield per plant (0.0659) and the correlation between the two traits is positive and non significant(0.1846) due to positive indirect effects through plant height (0.0076), panicle length (0.0159), number of panicles per plant (0.0104), number of filled grains per panicle (0.0083), grain length (0.0367), 1000 grain weight (0.0050), kernel length (0.0407, kernel breadth (0.0259), grain iron content

(0.0022) and negative indirect effects through days to 50% flowering (-0.0151), days to maturity (-0.0159) and grain zinc content (-0.0017). The results were in agreement with the findings of Ekka et al. [3], Mallimar et al. [22] and Patel et al. [15].

Number of filled grains per panicle reported a direct positive effect on grain yield per plant (0.0613) and the correlation between the two traits is positive and significant (0.5121). It expressed positive indirect effects through days to 50% flowering (0.0133), days to maturity (0.0085), plant height (0.0074), panicle length (0.0141), number of panicles per plant (0.0372), grain width (0.0077), 1000 grain weight (0.0025), kernel length (0.0062), kernel breadth (0.0116), grain iron content (0.0051), grain zinc content (0.0195) and negative indirect effect through grain length (-0.0042). The results were in agreement with the findings of Khare et al. [30], Ratna et al. [5], Bhati et al. [31], Babaeian and Bagheri [13] and Prasannakumari et al. [19].

1000 grain weight reported a direct positive effect on grain yield per plant (0.0480) and the correlation between the two traits is positive and non significant (0.0462) due to positive indirect effects through days to 50% flowering (0.0003), days to maturity (0.0041), plant height (0.0030), number of filled grains per panicle (0.0019), grain length (0.0041), grain width (0.0036), kernel length (0.0053), kernel breadth (0.0146), grain iron content (0.0027), grain zinc content (0.0037) and negative indirect effects through panicle length (-0.0026) and number of panicles per plant (-0.0010). The results were in agreement with the findings of Mallimar et al. [22], Nayak et al. [32], Babaeian and Bagheri [13], Kishore et al. [33], Shivani et al. [14], Islam et al. [18] and Parimala et al. [20].

Kernel breadth reported a direct positive effect on grain yield per plant (0.0220) and the correlation between the two traits is positive and non significant (0.0777) due to positive indirect effects through plant height (0.0011), panicle length (0.0015), number of

filled grains per panicle (0.0042), grain length (0.0021), grain width (0.0086), 1000 grain weight (0.0067), kernel length (0.0040), grain iron content (0.0015), grain zinc content (0.0040) and negative indirect effects through days to 50% flowering (-0.0028), days to maturity (-0.0009) and number of panicles per plant (-0.0002). The results were in agreement with the findings of Ekka et al. [3] and Prasannakumari et al. [19].

Grain zinc content shows a direct positive effect on grain yield per plant (0.0196) and the correlation between the two traits is positive and non significant (0.1410) due to positive indirect effects through days to 50% flowering (0.0046), days to maturity (0.0057), panicle length (0.0008), number of panicles per plant (0.0031), number of filled grains per panicle (0.0062), 1000 grain weight (0.0015), kernel breadth (0.0036), grain iron content (0.0125) and negative indirect effects through plant height (-0.0033), grain length (-0.0013), grain width (-0.0005) and kernel length (-0.0002). The results were in agreement with the findings of Bekele et al. [34] and Shivani et al. [14].

Grain length reported a direct positive effect on grain yield per plant (0.0118) and the correlation between the two traits is positive but non significant (0.0712) due to positive indirect effects through days to maturity (0.0008), panicle length (0.0006), grain width (0.0066), 1000 grain weight (0.0010), kernel length (0.0087), kernel breadth (0.0011), grain iron content (0.0009), and negative indirect effects through days to 50% flowering (-0.0026), plant height (-0.0016), number of panicles per plant (-0.0003), number of filled grains per panicle (-0.0008) and grain zinc content (-0.0008). The results were in agreement with the findings of Ekka et al. [3], Mallimar et al. [22] and Patel et al. [15].

Grain iron content shows a direct negative effect on grain yield per plant (-0.0899) and the correlation between the two traits is positive and non significant (0.0947) due to positive indirect effects through plant height (0.0011), panicle length (0.0015), and negative

indirect effects through days to 50% flowering (-0.0226), days to maturity (-0.0210), number of panicles per plant (-0.0180), number of filled grains per panicle (-0.0075), grain length (-0.0070), grain width (-0.0030), 1000 grain weight (-0.0051), kernel length (-0.0155), kernel breadth (-0.0062) and grain zinc content (-0.0573). The results were in agreement with the findings of Bekele et al. [34] and Singh et al. [12].

Kernel length reported a direct negative effect on grain yield per plant (-0.0434) and the correlation between the two traits is positive and non significant(0.1323) due to positive indirect effects through days to 50% flowering (0.0031), plant height (0.0011), grain zinc content (0.0005) and negative indirect effects through days to maturity (-0.0011), panicle length (-0.0110), number of panicles per plant (-0.0045), number of filled grains per panicle (-0.0044), grain length (-0.0318), grain width (-0.0268), 1000 grain weight (-0.0048), kernel breadth (-0.0078) and grain iron content (-0.0075). The results were in agreement with the findings of Ekka et al. [3] and Prasannakumari et al. [19].

Plant height shows a direct negative effect on grain yield per plant (-0.1005) and the correlation between the two traits is negative and non significant(-0.1755) due to negative indirect effects through panicle length (-0.0155), number of filled grains per panicle (-0.0122), grain width (-0.0115), 1000 grain weight (-0.0062), kernel breadth (-0.0052) and positive indirect effects through days to 50% flowering (0.0177), days to maturity (0.0328), number of panicles per plant (0.0107), grain length (0.0134), kernel length (0.0027), grain iron content (0.0254) and grain zinc content (0.0169). The results were in agreement with the findings of Islam et al. [18].

Days to 50% flowering reported a direct negative effect on grain yield per plant (-0.2121) and the correlation between the two traits is negative and non significant(-0.0550) due to indirect negative effect through days to days to maturity (-0.1198), number of

panicles per plant (-0.0134), number of filled grains per panicle (-0.0460), 1000 grain weight (-0.0012), grain iron content (-0.0534), grain zinc content (-0.0503) and positive indirect effect through plant height (0.0247), panicle length (0.0193), grain width (0.0485), grain length (0.0464), kernel length (0.0153) and kernel breadth (0.0274). The results were in agreement with the findings of Babu et al. [29] and Kalyan et al. [35].

CONCLUSION

The key variables that affect yield, such as the number of panicles per plant, the number of filled grains per panicle, and panicle length, all showed highly significant association values, according to research on correlation and path analysis. Additionally, during Path analysis, the same attributes showed greater values of direct effects on grain yield per plant, demonstrating the significance of these qualities and the need for selection criteria to maximise yield potential.

FUTURE SCOPE

The features that were beneficial for improving yield were determined through investigations on correlation and path coefficient analysis. The relationships between these traits also became quite obvious, and these relationships can be used in rice breeding programmes in the future to increase grain yield without affecting other traits like crop length and plant architecture. In order to improve yield effectively, the data can also be employed as a selection method.

REFERENCES

Chatterjee, D. (1947). Botany of the wild and cultivated rice. *Nature*, **160**: 234-237.

Patra, B.C. (2000). Collection and characterization of rice genetic resources from Keonjhar district of Orissa. *Oryza*, 34: 324-326.

Ekka, R.E., Sarawgi, A.K and Kanwar, R.R. 2011. Correlation and path analysis in traditional rice accessions of Chhattisgarh. *Journal of rice research*, 4(1), pp.11-18.

Dewey, D.R and Lu, K. 1959. A Correlation and Path-Coefficient Analysis of Components of Crested Wheatgrass Seed Production 1. *Agronomy Journal*. 51 (9): 515-518.

Ratna, M., Begum, S., Husna, A., Dey, S.R and Hossain, M.S. 2015. Correlation and path coefficients analyses in basmati rice. *Bangladesh Journal of Agricultural Research*. 40(1): 153-161.

Rahman, M.A., Hossain, M.S., Chowdhury, I.F., Matin, M.A and Mehraj, H. 2014. Variability study of advanced fine rice with correlation, path co-efficient analysis of yield and yield contributing characters. *International Journal of Applied Sciences and Biotechnology*. 2(3): 364-370.

Devi, R.K., Chandra, S.B., Lingaiah, N., Hari, Y and Venkanna, V. 2017. Analysis of Variability, correlation and path coefficient studies for yield and quality traits in rice (*Oryza sativa* L.). *Agricultural Science Digest*. 37(1) :1-9.

Immanuel S. C., Nagarajan, P., Thiyagarajan, K., Bharathi, M and Rabindran, R. 2011. Genetic parameters of variability, correlation and path coefficient studies for grain yield and other yield attributes among rice blast disease resistant genotypes of rice (*Oryza sativa* L.). *African Journal of Biotechnology*. 10(17): 3322-3334.

Umarani, E., Radhika, K., Padma, V and Rao, S.L.V. 2019. Agro-morphological characterization of rice (*Oryza sativa* L.) landraces based on DUS descriptors. *International Journal Pure Applied Biosciences*. 5(4):466-475.

Gangashetty, P. I., Salimath, P. M and Hanamaratti, N.G. 2013. Association analysis in genetically diverse non-basmati local aromatic genotypes of rice (*Oryza sativa* L.). *Molecular Plant Breeding*. 4.

Dore, V., Koti, R.V and Math, K.K. 2018. Response of zinc application on growth, zinc content and grain yield of rice genotypes and correlation between zinc content and yield attributes of rice genotypes. *Indian Journal of Agricultural Research*. 52 (6): 625-630.

Singh, A., Manjri, S.D.G., Kumar, G., Dubey, V., Rampreet, K.N and Dwivedi, D.K. 2018. Path coefficient analysis studies in Iron and Zinc containing rice varieties. *Journal of Pharmacognosy and Phytochemistry*. 7(2): 3729-3732.

Babaeian-Jelodar, N and Bagheri, N. 2018. Correlation and Path Coefficient Analysis in F₂ Generation of Rice Genotypes Derived from Crosses between Tarom-Jelodar and 229R Cultivars. *Journal of Crop Breeding*. 9 (24): 152- 157.

Shivani, D., Cheralu, C., Neeraja, C.N and Shankar, V.G. 2018. Grain Zinc and Iron Association Studies in Swarna X Type 3 RIL Population of Rice. *International Journal of Current Microbiology and Applied Sciences*. 7(5): 708-714.

Patel, J. R., Dixita, K., Patel, K. N., Prajapati, N. V., Soni and Ankit, P. 2017. Correlation and path coefficient analysis in rainfed upland rice (*Oryza sativa* L.). *Environment and Ecology*. 35(2): 789-794.

Pandey, S., Doss, D.D and Shashidhar, H.E. 2018. Correlation and path analysis of yield determinants and micronutrient content in rice (*Oryza sativa* L.). *Journal of*

Pharmacognosy and Phytochemistry. 7(1): 2723-2728.

Seneega, T.V., Gnanamalar, R.P., Parameswari, C., Vellaikumar, S and Priyanka, A.R. 2019. Genetic variability and association studies in F2 generation of rice (*Oryza sativa* L.). *Electronic Journal of Plant Breeding*. 10(2): 512-517.

Islam, M.Z., Mian, M.A.K., Ivy, N.A., Akter, N and Rahman, M.M., 2019. Genetic variability, correlation and path analysis for yield and its component traits in restorer rice. *Bangladesh Journal of Agricultural Research*. 44(2): 291-301.

Prasannakumari, M., Akilan, M., Kalaiselvan, S., Subramanian, A., Janaki, P and Jeyaprakash, P. 2020. Studies on genetic parameters, correlation and path analysis for yield attributes and Iron content in a backcross population of rice [*Oryza sativa* (L.)]. *Electronic Journal of Plant Breeding*. 11(03): 881-886.

Parimala, K., Raju, C.S., Prasad, A.H., Kumar, S.S and Reddy, S.N. 2020. Studies on genetic parameters, correlation and path analysis in rice (*Oryza sativa* L.). *Journal of Pharmacognosy and Phytochemistry*. 9(1): 414-417.

Sudeepthi, K., Srinivas, T., Kumar, B.R., DPB, J and Umar, S.N. 2020. Genetic variability, character association and path analysis for anaerobic germination traits in rice (*Oryza sativa* L.). *Journal of Pharmacognosy and Phytochemistry*. 9(1): 553-556.

Mallimar, M., Surendra, P., Hundekar, R., Jogi, M and Lakkangoudar, M.C.S. 2016. Correlation studies for micronutrients, yield and yield components in F3 population of rice (*Oryza Sativa* L.). *Res. Environ. Life Sci*. 9(9): 1140-1142.

Hasan, M.J., Kulsum, M.U., Akter, A., Masuduzzaman, A.S.M and Ramesha, M.S. 2013. Genetic variability and character association for agronomic traits in hybrid rice (*Oryza sativa* L.). *Bangladesh Journal Plant Breeding and Genetics*. 24 (1): 45-51.

Oliveira, A.L., Tan, L., Fu, Y and Sun, C. 2009. Genetic identification of quantitative trait loci for contents of mineral nutrients in rice grain. *Journal of Integrative Plant Biology*. 51(1): 84-92. *International Journal of Current Microbiology and Applied Sciences*. 7(09): 97-108.

Raza, Q., Saher, H., Shahzadi, F., Riaz, A., Bibi, T and Sabar, M. 2019. Genetic diversity in traditional genotypes for grain iron, zinc and beta-carotene contents reveals potential for breeding micronutrient dense rice. *Journal of Experimental Biology and Agricultural Sciences*. 7(2):194-203.

Edukondalu, B., Reddy, V. R., Rani, T. S., Kumari, C. A and Soundharya, B. 2017. Studies on variability, heritability, correlation and path analysis for yield, yield attributes in rice (*Oryza sativa* L.). *International Journal of Current Microbiology and Applied Sciences*. 6(10): 2369-2376.

Archana, R.S., Rani, M.S., Vardhan, K.V and Fareeda, G. 2018. Genetic diversity studies among rice (*Oryza sativa* L.) genotypes for grain yield, yield components and nutritional traits in rice. *International Journal of Communication Systems*. 6 (6): 134-137.

Rachana, B., Eswari, K. B., Jyothi, B and Raghuvier Rao, P. 2018. Correlation and path analysis for yield and its component traits in NPT core set of rice (*Oryza sativa* L.).

Babu, V.R., Shreya, K., Dangi, K.S., Usharani, G and Nagesh, P. 2012. Genetic variability studies for qualitative and quantitative traits in popular rice (*Oryza sativa* L.) hybrids of India. *International Journal of Scientific and Research Publications*. 2 (6): 1-5.

Khare, R., Singh, A.K., Eram, S and Singh, P.K. 2014. Genetic variability, association and diversity analysis in upland rice (*Oryza sativa* L.). *SAARC Journal of Agriculture*. 12

(2): 40-51.

Bhati, M., Babu, G.S and Rajput, A.S. 2015. Genetic variability, correlation and path coefficient for grain yield and quantitative traits of elite rice (*Oryza sativa* L.) genotypes at Uttar Pradesh. *Electronic Journal of Plant Breeding*. 6 (2): 586- 591.

Kishore, C., Kumar, A., Pal, A.K., Kumar, V., Prasad, B.D and Kumar, A. 2018. Character association and path analysis for yield components in traditional rice (*Oryza sativa* L.) genotypes. *International Journal of Current Microbiology and Applied Sciences*. 7(3): 283-291.

Nayak, R., Singh, V.K., Singh, A.K and Singh, P.K. 2016. Genetic variability, character association and path analysis of rice genotypes. *Annals of Plant and Soil Research*.18(2): 161-164.

Bekele, B.D., Rakhi, S., Naveen, G.K., Kundur, P.J and Shashidhar, H.E. 2013. Estimation of genetic variability and correlation studies for grain zinc concentrations and yield related traits in selected rice (*Oryza sativa* L.) genotypes. *Asian Journal of Experimental Biological Sciences*. 4(3): 345-351.

Kalyan, B., Radha Krishna, K.V and Rao, L.V.S. 2017. Path coefficient analysis for yield and yield contributing traits in rice (*Oryza sativa* L.) genotypes. *International Journal of Current Microbiology and Applied Sciences*. 6(7): 2680-2687.

Table 1. Correlation coefficients of yield and yield attributing traits in rice genotypes

Character	DFE	DM	PH	PL	NPP	NFG	GL	GW	TGW	KL	KB	FC	ZC	GYP
DFE	1.000	0.5649***	-0.1162	-0.1162	0.0632	0.2170	-0.2189	-0.2288	0.0055	-0.0723	-0.1289	0.2519*	0.2372*	-0.0549
DM		1.000	-0.3267**	0.0212	0.0673	0.1383	0.0639	-0.2411*	0.0864	0.0252	-0.0417	0.2340	0.2916*	0.1750
PH			1.000	0.1545	-0.1065	0.1214	-0.1337	0.1146	0.0620	-0.0264	0.0515	-0.2530*	-0.1680	-0.1755
PL				1.000	0.2396 *	0.2306	0.0547	0.2411 *	-0.0546	0.2525 *	0.0692	-0.0633	0.0424	0.3118 **
NPP					1.000	0.6061***	-0.0263	0.1582	-0.0210	0.1047	-0.0089	0.2008	0.1570	0.7991 ***
NFP						1.000	-0.0688	0.1255	0.0403	0.1013	0.1898	0.0830	0.3180**	0.5121 ***
GL							1.000	0.5569 ***	0.0861	0.7323 ***	0.0941	0.0780	-0.0658	0.0712
GW								1.000	0.0756	0.6173 ***	0.3923 ***	0.0334	-0.0257	0.1846
TGW									1.000	0.1109	0.3037*	0.0569	0.0771	0.0461
KL										1.000	0.1802	0.1724	-0.0112	0.1323
KB											1.000	0.0694	0.1815	0.0776
FC												1.000	0.6378 ***	0.0946
ZC													1.000	0.0609

* Significance at p = 0.01 **Significance at p = 0.05

DFE- Days to 50% flowering, **DM**- Days to maturity, **PH**- Plant height, **PL**- Panicle length, **NPP**- Number of panicles per plant, **NFP**- Number of filled grains per panicle, **GL**- Grain length, **GW**- Grain width, **TGW**- 1000 grain weight, **KL**- Kernel length, **KB**- Kernel breadth, **FC**- Fe(iron) content, **ZC**- Zn(zinc) content, **GYP**- Grain yield per plant.

Table. 2 Path coefficients of yield and quality traits in rice

CHARACTER	DFE	DM	PH	PL	NPP	NFG	GL	GW	TGW	KL	KB	FC	ZC	GYE
DFE	-0.2121	-0.1198	0.0247	0.0193	-0.0134	-0.0460	0.0464	0.0485	-0.0012	0.0153	0.0274	-0.0534	-0.0503	-0.0550
DM	0.1301	0.2302	-0.0752	0.0049	0.0155	0.0318	0.0147	-0.0555	0.0199	0.0058	-0.0096	0.0539	0.0671	0.1751
PH	0.0117	0.0328	-0.1005	-0.0155	0.0107	-0.0122	0.0134	-0.0115	-0.0062	0.0027	-0.0052	0.0254	0.0169	-0.1755
PL	-0.0092	0.0022	0.0157	0.1018	0.0244	0.0235	0.0056	0.0246	-0.0056	0.0257	0.0071	-0.0065	0.0043	0.3118
NPP	0.0465	0.0495	-0.0783	0.1762	0.7353	0.4457	-0.0193	0.1164	-0.0155	0.0771	-0.0066	0.1477	0.1155	0.7991
NFE	0.0133	0.0085	0.0074	0.0141	0.0372	0.0613	-0.0042	0.0077	0.0025	0.0062	0.0116	0.0051	0.0195	0.5121
GL	-0.0026	0.0008	-0.0016	0.0006	-0.0003	-0.0008	0.0118	0.0066	0.0010	0.0087	0.0011	0.0009	-0.0008	0.0712
GW	-0.0151	-0.0159	0.0076	0.0159	0.0104	0.0083	0.0367	0.0659	0.0050	0.0407	0.0259	0.0022	-0.0017	0.1846
TGW	0.0003	0.0041	0.0030	-0.0026	-0.0010	0.0019	0.0041	0.0036	0.0480	0.0053	0.0146	0.0027	0.0037	0.0462
KL	0.0031	-0.0011	0.0011	-0.0110	-0.0045	-0.0044	-0.0318	-0.0268	-0.0048	-0.0434	-0.0078	-0.0075	0.0005	0.1323
KB	-0.0028	-0.0009	0.0011	0.0015	-0.0002	0.0042	0.0021	0.0086	0.0067	0.0040	0.0220	0.0015	0.0040	0.0777
FE	-0.0226	-0.0210	0.0227	0.0057	-0.0180	-0.0075	-0.0070	-0.0030	-0.0051	-0.0155	-0.0062	-0.0899	-0.0573	0.0947
ZN	0.0046	0.0057	-0.0033	0.0008	0.0031	0.0062	-0.0013	-0.0005	0.0015	-0.0002	0.0036	0.0125	0.0196	0.1410

Residual effect = 0.5236. **DFE**- Days to 50% flowering, **DM**- Days to maturity, **PH**- Plant height, **PL**- Panicle length, **NPP**- Number of panicles per plant, **NFE**- Number of filled grains per panicle, **GL**- Grain length, **GW**- Grain width, **TGW**- 1000 grain weight, **KL**- Kernel length, **KB**- Kernel breadth, **FE**- Fe(iron) content, **ZC**- Zn(zinc) content, **GYE**- Grain yield per plant.

Figure 1. Phenotypic correlation coefficients among different traits in rice

