

## **COMPARATIVE STUDY OF BIOFFERTILIZER AND PLANT EXTRACTS ON GROWTH, YIELD AND YIELD ATTRIBUTING TRAITS OF FIELD PEA (*Pisum sativum*).**

### **ABSTRACT**

The field experiment entitled “Comparative Study of Biofertilizer and Plant Extracts on Growth, Yield and Yield Attributing Traits of Field pea (*Pisum Sativum* L.)” was conducted during Rabi 2021 at field experimental centre, Department of Genetics and Plant Breeding, SHUATS, Prayagraj, (U.P). The soil of experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.36 %), available N (171.48 kg/ha), available P (15.2 kg/ha) and available K (232.5 kg/ha). The experiment was laid out in Randomized Block Design with thirteen treatments including control which were replicated thrice. The treatments are as follows, TO-Control, T1, T2, T3, - Rhizobium @ 5%, 8%, 10%, T4,T5,T6 – Vermi wash @ 5%, 8%, 10%, T7,T8,T9 – Neem leaf extract @ 5%, 10%, 15% and T10, T11, T12 – Tulsi leaf extract @ 5%, 10%, 15% respectively. The experiment results revealed that seeds treated with Rhizobium @ 8% gave better than other treatments viz, Field emergence (87.47 %), Plant height (95.71 cm), Days to 50% flowering (40.60), number of branches per plant (8.87), Number of pods per plant (13.53), Number of seeds per pod (8.47), seed yield per plant (24.88 g), seed yield per plot (746.50 g), biological yield (33.73 kg/ha), Harvest index (73.48 %) were recorded significantly higher compared to other treatments.

**Key words: Rhizobium, Vermiwash, Neem leaf extract, Tulsi leaf extract, Seed Priming.**

## INTRODUCTION:

The Field pea is one of the oldest domesticated crops, cultivated for at least 7,000 years. Field peas are now grown in many countries for both human consumption and stock feed. It is a climbing annual legume with weak, viny, and relatively succulent stems. Vines often are 4 to 5 feet (120 to 150 cm) long, but when grown alone, field pea's weak stems prevent it from growing more than 1.5 to 2 feet (45 to 60 cm) tall. Leaves have two leaflets and a tendril. Flowers are white, pink, or purple. Pods carry seeds that are large (4,000 seeds/lb), nearly spherical, and white, gray, green, or brown. The root system is relatively shallow and small, but well nodulated. The field pea is a cool-season legume crop that is grown on over 25 million acres worldwide. It has been an important grain legume crop for millennia, seeds showing domesticated characteristics dating from at least 7000 years ago have been found in archaeological sites around what is now Turkey. Field peas or "dry peas" are marketed as a dry, shelled product for either human or livestock food, unlike the garden pea, which is marketed as a fresh or canned vegetable. Peas, like many legumes, contain symbiotic bacteria called Rhizobia within root nodules of their Root system. These bacteria have the special ability to fix N<sub>2</sub> from atmospheric, molecular nitrogen (N<sub>2</sub>) into Ammonia (NH<sub>3</sub>).

Rhizobium starvation is common in micronutrient deficient soils. Seeds treated with Rhizobium shows maximum germinability and vigour index. Now a days seeds treated with Biofertilizers shows best results producing high yield and enhance plant growth (Maurya *et al.*2021). It also acts as a plant growth promoter by mechanisms of tolerance of abiotic stresses. Vermiwash is a liquid extract produced from vermicompost in a medium where earthworms are richly populated. It comprises a massive decomposer bacteria count, mucus, vitamins, different bioavailable minerals, hormones, enzymes, different antimicrobial peptides, etc. Seeds primed with Vermiwash results in highest field emergence and better yield in various field crops (Manjunadh *et al.* 2021). It acts as a plant growth promoter. Rhizobium biofertilizer contains symbiotic Rhizobium bacteria which is the most important nitrogen-fixing organism. These organisms have the ability to drive atmospheric Nitrogen and provide it to plants. It is recommended for crops such as Groundnut, Soybean, Red-gram, Greengram, Black-gram, Lentil, Cowpea, Field pea, Bengal-gram and Fodder legumes, etc. They are the most efficient biofertilizer as per the quantity of Atmospheric Nitrogen fixed. Vermiwash is nutrient rich liquid manure, extracted from vermicomposts riches with a greater number of earthworms feeding on organic waste material and plant residues. It is non-toxic and eco-friendly, which arrests bacterial growth and forms as a protective layer for their survival and growth. Vermiwash contains N, P, K, Ca & hormones such as auxin, cytokinin, some other secretions and many useful microbes like heterotrophic bacteria, fungi etc.). Neem leaf extract is a powerful insecticide which contains an pivotal insecticidal ingredient called Azadirachtin which repelling them, by inhibiting feeding, and by disrupting their growth, metamorphosis and reproduction. Azadirachtin affects insect physiology by mimicking a natural hormone. It has been shown to affect egg production and hatching rates. In larvae, azadirachtin can inhibit molting, preventing them from developing into pupae. Tulsi leaf extract is an anti fungal agent and an insecticide but it shows least inhibition over control against fungi such as *Aspergillus niger*, *Alternaria alternata*, *Penicillium rubrum*, and *Fusarium moniliformae*. But neem leaf extract was more effective than tulsi leaf extract..

## **MATERIALS AND METHODS:**

The present research on **EFFECT OF VARIOUS PRESOWING SEED TREATMENTS ON GROWTH AND YIELD PARAMETERS OF FIELD PEA (*Pisum sativum.L*)** was made to identify the effect of seed priming of different kinds on seed quality parameters of mustard and to find out suitable seed priming method for mustard. For this purpose, 13 priming treatments including control on Field pea seeds were used to study under field conditions during Rabi, 2021-22. The treatments were T0–Control, T1-Rhizobium (5%), T2- Rhizobium (8%), T3- Rhizobium (10%), T4- Vermiwash (5%), T5- Vermiwash (8%), T6- Vermiwash (10%), T7-Neem leaf extract (5%), T8-Neem leaf extract (10%), T9-Neem leaf extract (15%), T10-Tulasi leaf extract-(5%), T11-Tulasi leaf extract-(10%), T12-Tulasi leaf extract-(15%). The mustard seeds were primed with above different priming agents in above different concentrations and intensities for a given duration. After priming seeds were dried to initial moisture content at room temperature. After that the primed seeds were used to grow under field conditions. Field experiment was laid out in Randomized Block Design (RBD) with three replications during Rabi 2021-22. Data were recorded for 10 characters i.e.

**Pre-harvest characters viz.,** Field emergence percentage, Plant height (60 DAS), number of primary branches (at harvest), 50% of flowering and Days to maturity.

**Post-harvest characters viz.,** number of pods/plant, number of seeds/pod, seed yield/plant (gm), seed yield/plot (gm), biological yield (gm), harvest index (%). Analysis of variance (Table-1) for the data revealed that significance mean sum of squares due to seed priming treatments were observed for all the characters under study viz., Field emergence percentage, Plant height (60, at harvest), number of primary branches (at harvest), 50% of flowering and Days to maturity, number of pods/plant, number of seeds/pod, seed yield/plant (gm), seed yield/plot (gm), biological yield (gm), harvest index (%). which were highly significant at 5% level of significance indicating presence of good amount of variability among the treatments for these characters.

## **RESULTS AND DISCUSSION:**

### **GROWTH ATTRIBUTES:**

Seed treatment is a technique to decrease the emergence time, produce uniform emergence time, better allometric (changes in growth of plant parts over time) attributes and provide requisite stand in many horticultural and field crops. Various pre-hydration or priming treatments have been employed to increase the speed and synchrony of seed germination. The treatments showed significant effect of pre-sowing seed treatment on field emergence. The higher mean performance of field emergence was observed in the treatment using rhizobium with the concentration of 8% (2309), followed by treatment of seeds with neem leaf extract with 10% concentration (2272), and the plant height was observed highest in the treatment using Rhizobium @ 8% (95.71cm) followed by treatment with neem leaf extract @ 10% (92.54). In the performance with the parameter 50% flowering, treatment with Rhizobium @ 8% showed significant value (40.6) which is followed by the seed treatment using neem leaf extract @ 10% (41.4) and days to pod maturity was observed highest in the seed treatment using Rhizobium @ 8% (101.43), followed by the treatment using neem leaf extract @ 10% (102.83) and in the consideration of parameter, number of primary branches seed treatment by using Rhizobium @ 8% shown effective results with the mean value range of 8.87 which is followed by the treatment of seeds with neem leaf extract @ 10% (7.8). Based on the above observations seed treatment by using Rhizobium @ 8% shown considerable results in the growth parameters on comparison with the other treatments.

Table 1A: Mean performance of different treatments for pre harvest characters in Field Pea (*Pisum sativum*) .

TREATMENTS	CONCENTRATIONS(%)	Rate of field emergence	Plant Height(cm)	Days to 50% flowering	Days to Maturity	Number of Branches
Control	-	1850	74.85	44.6	107.56	5.53
Rhizobium	5%	2053	89.66	44.07	106.48	6.47
Rhizobium	8%	2309	95.71	40.6	101.43	8.87
Rhizobium	10%	2114	82.78	43.1	105.66	6.73
Vermiwash	5%	2083	86.01	43.27	107.1	6.13
Vermiwash	8%	2136	85.68	43.23	106.12	6.67
Vermiwash	10%	2237	87.27	42.93	106.92	6.13
Neem leaf extract	5%	2201	83.38	45.2	104.31	7.13
Neem leaf extract	10%	2272	92.54	41.4	102.83	7.8
Neem leaf extract	15%	1945	88.62	43.2	107.43	5.73
Tulsi leaf extract	5%	2020	85.63	45	106.76	6.33
Tulsi leaf extract	10%	1898	84.73	44.27	103.79	7.27
Tulsi leaf extract	15%	1991	82.37	43	105.02	6.93
<b>F Test</b>			S	S	S	S
<b>S.EM (±)</b>			0.54	0.41	1.21	0.14
<b>CD (p=0.05)</b>			1.59	1.2	3.54	0.42

Table 1B: Mean performance of different treatments for postharvest characters in Field Pea (*Pisum sativum*).

TREATMENTS	CONCENTRATIONS	Numbers of pods per plant	Number of seeds per pod	Seed yield per plant (gm)	Seed yield per plot (gm)	Biological yield	Harvest Index
Control	-	10.8	5.87	13.25	397.4	24.28	55.1
Rhizobium	5%	11.4	7	17.09	512.57	25.86	66.47
Rhizobium	8%	13.53	8.47	24.88	746.5	33.73	73.48
Rhizobium	10%	13	7.2	20.9	627.09	30.37	64.62
Vermiwash	5%	11.47	6.47	19.95	598.37	29.49	68.42
Vermiwash	8%	11.67	7.6	18.12	543.56	28.84	63.03
Vermiwash	10%	12.6	7.2	19.41	582.29	29.36	66.46
Neem leaf extract	5%	11.47	7.6	16.96	508.78	26.66	68.84
Neem leaf extract	10%	13.2	8	22.09	662.83	32.33	68.99
Neem leaf extract	15%	13.17	6.07	16.07	482.07	27.43	58.16
Tulsi leaf extract	5%	12.2	7.33	16.24	487.14	26.74	61.12
Tulsi leaf extract	10%	12.87	7.87	21.86	655.93	30.88	70.77
Tulsi leaf extract	15%	12.87	6.87	18.21	546.27	28.44	64.15
F Test		S	S	S	S	S	S
S.EM (±)		0.16	0.16	0.57	16.97	1.01	2.22
CD (p=0.05)		0.45	0.46	1.65	49.55	2.95	6.49

## **YIELD ATTRIBUTES:**

The mean performance of number of pods/plant was recorded highest in the seed treatment using Rhizobium @ 8% concentration with the mean value range of (13.53 pods) which is followed by treating of seeds with neem leaf extract @ 10% concentration (13.2) and in the case of number of seeds per pod the treatment of seeds with the treatment using Rhizobium @ 8% shown the highest mean performance (8.47) which is followed by seed treatment using Neem leaf extract @ 10% (8). The parameter of seed yield per plant when taken under consideration, the seed treatment using Rhizobium @ 8% recorded the higher mean performance (24.88), followed by treating of seeds with Neem leaf extract @ 10% (22.09) and seed yield per plot was recorded higher in the treatment of seeds with Rhizobium @ 8% ( 746.5) which is followed by treating of seeds with Neem leaf extract @ 10% (662.83).In the parameter of biological yield higher values were recorded in the treatment of Rhizobium @ 8% (33.73), followed by seed treatment using Neem leaf extract @ 10 % concentration (32.33). On considering the harvest index the treatment using Rhizobium @ 8% recorded the higher mean value (73.48) and it was followed by treatment using Neem leaf extract @ 10% (68.99).These recorded mean performances are comparatively higher than the control used and the treatment of seeds with the Rhizobium @ 8% recorded significant results in the yield attributes of the plant.

## **CONCLUSION:**

It is concluded that seeds treated with Rhizobium @ 8% was found to be more desirable for producing significantly higher seed yield per plant (24.88 g), seed yield per plot(746.50 g). Findings are based on research done in one season in Prayagraj (Allahabad) U.P..These recommendations are based on Three months experimentation and therefore further investigation is needed to arrive at valid recommendation.

## REFERENCE:

- Abdul Wajid, G., Kalneni Jahnavi, Arun Kumar Chaurasia, Bharath Reddy, N. and Prudvi Raj Naidu, B. (2021). Effect of Different Organic and Inorganic Seed Priming Method on Growth, Yield and Quality Parameters of Field Pea (*Pisum sativum* L.). *Int.J.Curr.Microbiol.App.Sci.* **10**(01): 280-286.
- Abebe Megersa (2016). Botanicals extracts for control of pea aphid (*Acyrtosiphon pisum*; Harris). *Journal of Entomology and Zoology Studies.* **4**(1): 623-627.
- Abhishek, S and Sobita Simon (2017). Eco-friendly management of powdery mildew and rust of garden pea (*Pisum sativum* L.). *Journal of Pharmacognosy and Phytochemistry.* **6**(5) : 90-93.
- Akhilesh Patel., Mohd. Danish., Mukesh Kumar Prajapati., Anoop Kumar and Abhilasha A Lal. (2021). In vitro efficacy of plant extracts against *Alternaria lini* (leaf blight of linseed). *The Pharma Innovation Journal.* **10**(4): 675-679.
- Alam, M.S., Solaiman, A.R.M., Rahman, G.K.M.M., Rahman, M.M. and Islam, M.M, (2015). Evaluation of Rhizobium Isolates in Terms of Nodulation, Growth and Yield Of Chickpea. *Bangladesh J. Soil Sci.* **37** (1): 35-45.
- Anjeneth S. Francisco., Njeneth S. Francisco., Felix E. Arcilla Jr. (2019). Agronomic Performance of Mungbean (*Vigna Radiata*) using Vermiwash and Seed Inoculant. *AMURE International Journal of Ecology and Conservation.* **29** : 17-42.
- Archana W Shingne, Dr. Giri, GK., and Ashwini R Bagade. (2020). In vitro evaluation of fungicides, botanicals and bio-agents against *Alternaria alternata* causing leaf spot disease of niger. *International Journal of Chemical Studies.* **8**(4): 3360-3364.
- Bhumi Reddy Divyavani, Ganesh, V., Dhanuka, D. (2020). Effect of integrated nutrient management on growth and yield in black gram (*Vigna mungo* L. *Hepper*) under doon valley condition. *Journal of Pharmacognosy and Phytochemistry.* **9**(5): 2928-2932.
- Choudhary, C.S., Jain,S.C., Ritesh Kumar and Jaipal Singh Choudhary. (2013). Efficacy of different fungicides, biocides and botanical extract seed treatment for controlling seed-borne colletotrichum sp. in chilli (*Capsicum annuum* L.). *The Bioscan.* **8**(1) : 1-4.
- Deeksha Gupta., Shilpa Sivadas., Keerthi Vikram., Suneetha V (2016). To assess the role of natural pesticides made from Tulsi *ocimum tenuiflorum*, Turmeric *curcuma longa* and *Neem* *azadirachta indica* on Culinary crops and its maturing soil. *International Journal of AppliedPharmaceutics.* **8**(2) : 13-15.
- Deepak Chand Bhatishwar., Deepti Prabha., Deepak Jangid and Mohammad Salman. (2020). Effect of

Seed Priming with Botanicals on Plant Growth and Seed Yield of Lentil (*Lens culinaris* M.).

*Int.J.Curr.Microbiol.App.Sci.* **9**(7): 3484-3499.

- Dheeba, B., Niranjana, R., Sampathkumar, P. (2015). Efficacy of Neem (*Azadirachta Indica*) and Tulsi (*Ocimum Sanctum*) Leaf Extracts Against Early Blight of Tomato. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci.* **85**, 327–336.
- Kanaga, A., Muniyammal, P., and Manonmani, R. (2012). Suppressive Effects of Aqueous Extracts of Neem (*Melia Azadirachta* L.) on Some Initial Growth Parameters of Cow Pea & Horse Gram. *Journal of Applied Pharmaceutical Science.* **2**(5): 185-187.
- Kasarkar, A.R., and Barge, A.N. (2016). Effect of aqueous extract of neem (*Azadirachta indica* A. Juss) leaves on germination and growth of some agricultural crops. *Journal of Medicinal Plants Studies.* **4**(5): 11-13.
- Keya Prajapati., Patel, P.H and Rabari, S.R. (2022). Effect of Foliar Spray of Vermiwash and Panchgavya on Growth, Yield and Economics of Field Pea. *Biological Forum – An International Journal.* **14**(2): 117-120.
- Maheswari, V.N., Srikumaran, M.P., Rekha, G.S., Elumalai, D., and Kaleena, P.K. (2016). Growth promoting effects of vermiwash and panchagavya on *Dolichus lablab* under field experimental conditions. *Int J Appl Sci Biotechnol.* **4**(4): 513-518.
- Molla, M.N., Solaiman, A.R.M., Jahiruddin, M., Mridha, M.A.U., Khanam, D. (2011). Influence of different doses of phosphorus in the presence of arbuscular mycorrhiza and Rhizobium on the growth and yield of mungbean. *Bull. Inst. Trop. Agr., Kyushu Univ.* **34**: 49-67.
- Moomal Bharadwaj, Lakhawat, S.S., Upadhaya, B., Shalini Pilia, Devendra Jain and Bunker, R.N. (2021). Effect of organic liquid manures on vegetative growth and yield of pea (*Pisum sativum* L.). *The Pharma Innovation Journal.* **10**(9): 1360-1364.
- Murat Erman., Bunyamin Yildirim., Necat Togay and Fatih Cig (2010). Effect of Phosphorus Application and Rhizobium Inoculation on the Yield, Nodulation and Nutrient Uptake in Field Pea (*Pisum sativum* sp. *arvense* L.). *Journal of Animal and Veterinary Advances.* **8**(2): 301-304.
- Nangial Khan., Farooq Nawaz., Abdullah Khan., Naveed Ul Haq., Saif Ullah, Anis Ur Rehman Khalil., Sohail., Adeel Liaqat., Murad Ali., Junaid Ali Shah., Israr Ali and Mehran Ali (2017). Effect of farmyard manure and rhizobium inoculation on growth of chickpea (*Cicer arietinum* L.) variety “karak-03”. *Pure Appl. Biol.* **6**(1): 378-384.
- Nelakurthi Venkata Praveen, Prashant Kumar Rai, Rupesh Kumar and Ruksana. (2020). Effect of Priming on the Growth, Seedling, Yield and its Attributing Characters of Desi Chickpea

(*Cicer arietinum* L.). *Int.J.Curr.Microbiol.App.Sci.* **9**(09): 3162-3171.

- Ngeno Jonah., Chemining wa George, N., Muthomi James, W., and Shibairo Solomon, I. (2012). Effect of Rhizobium inoculation and Nitrogen fertilizer application on growth, nodulation and yield of two garden pea genotypes. *Journal of Animal & Plant Sciences.* **15**(2): 2147-2156.
- Nwankwo, E.N., Onuseleogu, D.C., Ogbonna Confidence U., Okorochoa, A.O.E (2016). Effect of neem leaf extracts (*Azadirachta indica*) and synthetic pesticide (Carbofuran) on the root-knot nematode (*Meloidogyne* spp) of cowpea (*Vigna unguiculata* L. Walp). *International Journal of Entomology Research.* **1**(3): 01-06.
- Putta Rajesh., Arun Kumar Chaurasia and Jagadam Reddy Kiran (2021). Effect of different invigoration treatments on growth and yield parameters in Cowpea (*Vigna unguiculata* L.). *The Pharma Innovation Journal.* **10**(5): 574-576.
- Rahman, M.M., Solaiman, A.R.M., Khanam, D., Sirajul Karim, A.J.M and Karim, M.A. (2010). Effects of Inoculation with Rhizobium and Arbuscular Mycorrhiza and Phosphorus on Growth, Yield and Nutrient Uptake by Pea Grown in Soil. *Bangladesh J Microbiol.* **27**(1): 22-27.
- Rajan, M.R., Murugesan, P. (2012). Influence of Vermiwash on Germination and Growth of Cow Pea *Vigna Unguiculata* and Rice *Oryza Sativa*. *IOSR Journal of Pharmacy.* **2**(6) : 31-34.
- Saad Abou-El-Hassan and Heba S. Elbatran (2020). Production of Pea without Chemical Fertilizers via Integrating Biofertilizers with Vermiwash. *Plant Archives.* **20**(2): 4319-4325.
- Sakpal, V.M., Jagtap,D.N., Upadhyay, L., Pinjari, S.S., More, S.S., Dhekale, J.S., Jadhav, M.S.,Rajemahadik, V.A and Bodake, P.S. (2021). Response of Cowpea (*Vigna Unguiculata* L.) to Foliar Application of Different Organic Sources and Levels of Fertilizer. *Chemical Science Review and Letters.* **10**(38), 269-273.
- Saurabh Singh., Seweta Srivastava., Jyotiranjana Mishra., Richa Raaj and Asha Sinha. (2014). Evaluation of Some Plant Extract against Predominant Seed Mycoflora of Mungbean (*Vigna Radiata* L.) Wilczek Seed. *Life Sciences Leaflets.* **51** : 83-89.
- Simran Kour, Bairwa,H.L., Lakhawat, S.S., Sharma, S.K., Shalini Pliania B Upadhyay and Dangi, N.L.(2021). Effect of organic manures and vermiwash on growth, yield and quality of cowpea [*Vigna unguiculata* (L.) Walp.] cv. Pusa Komal. *The Pharma Innovation Journal.* **10**(9): 543-548.
- Solaiman, A.R.M., Rabbani, M. G., Hossain, D., Hossain, G.M.A., and Alam, M. S. (2012). Influence of Phosphorus and Inoculation with Rhizobium and AM Fungi on Growth and Dry Matter Yield of Chickpea. *Bangladesh J. Sci. Res.* **25**(1): 23-32.

- Srikanth, K. and Dr. Chaurasia, A.K. (2021). Influence of botanicals, coconut water and PGPR treatments on plant growth, nodulation, yield and seed quality parameters of chick pea (*Cicer arietinum* L.). *The Pharma Innovation Journal*. **10**(12) : 2799-2802.
- Talwinder Singh., Harish Chandra Raturi., Dilip Singh Kachwaya., Sandeep Kumar Singh and Arinderpal Kaur (2019). Effect of biofertilizer and mulch on yield and quality of pea (*Pisum sativum* L.). *Journal of Pharmacognosy and Phytochemistry*. SPI : 205-208.
- Tarun Sharma , Janardan Singh , Navneet Kaur and Shilpa (2020). Effect of organic sources of nutrients on growth and growth indices of cowpea (*Vigna unguiculata*) under mid hill conditions of Himachal Pradesh. *Himachal Journal of Agricultural Research*. **46**(2) : 201-204.
- Tauseef A. Bhat., Meenakshi Gupta., Sheraz Mahdi, S., Manzoor A. Ganai., Hilal A. Bhat., Bhat, J.A., Iqbal A. Wani and Hussain Dar, M. (2013). Growth, Yield and Economics of Field pea (*Pisum sativum* L.) as Influenced by Phosphorus and Bio-fertilizers under Subtropical Conditions of Jammu. *Journal of Pure and Applied Microbiology*. **7**(1): 645-652.
- Tharuna Sree., Abhinav Dayal and Prashanth Kumar Rai (2021). Comparative Effect of Different Treatments in Field Pea (*Pisum sativum* L.). *International Journal of Environment and Climate Change*. **11**(10): 221-225.
- Vivek Mishra., Abhilasha A Lal and Sobita Simon (2017). Efficacy of botanicals and bio-agents against powdery mildew disease of garden pea (*Pisum sativum* L.). *Journal of Pharmacognosy and Phytochemistry*. **6**(4) : 1125-1126.
- Vivek Mishra., Abhilasha A Lal and Sobita Simon (2017). Efficacy of botanicals and bio-agents against rust disease of garden pea (*Pisum sativum* L.). *Journal of Pharmacognosy and Phytochemistry*. **6**(4) : 1125-1126.

UNDER PEER REVIEW