

Effect of different varieties, nitrogen levels and cutting management on growth and yield of fodder bajra (*Pennisetum glaucum* L.)

ABSTRACT

Aims: To identify an ideal pearl millet variety under ideal nitrogen level and cutting management for high green fodder yield and dry fodder yield.

Study design: Randomized block design with factorial concept.

Place and duration of study: AICRP on Forage Crops and Utilization, Agricultural Research Institute, between July 2021 and Nov 2021.

Methodology: Field experiment was conducted at AICRP on Forage Crops and Utilization, ARI, Rajendranagar, Hyderabad during *kharif* 2021. The treatments consisted of three varieties (TSFB 15-4, TSFB 15-8 and Moti bajra), three nitrogen levels (80, 100 and 120 kg N ha⁻¹) and two cutting management practices (C₁: Two cuts: 1st at 60 DAS, 2nd cut at 50% flowering) (C₂: Three cuts: 1st at 50 DAS, 2nd cut at 35 days after 1st cut and 3rd cut at 50% flowering) laid out in randomized block design with factorial concept, with Factor (A) as varieties, Factor (B) as nitrogen levels and Factor (C) as cutting management with three replications. The soil was sandy loam in texture with pH of 7.0 low in available nitrogen, medium in available phosphorus and available potassium.

Results: Variety TSFB 15-8 recorded significantly higher plant height (67.2 cm) and leaf-stem ratio (0.4) green forage yield (821.1 q ha⁻¹), dry fodder yield (198.4 q ha⁻¹) respectively, but the number of tillers m⁻² was recorded highest in Moti bajra (4.5). Application of nitrogen 120 kg ha⁻¹ significantly recorded higher plant height (72.0 cm), number of tillers m⁻² (4.7) and leaf-stem ratio (0.4), green forage yield (817.7 q ha⁻¹), dry fodder yield (201.0 q ha⁻¹). Three cuts for green fodder recorded significantly higher plant height (65.9 cm), number of tillers (5.1) and leaf-stem ratio (0.5) green fodder yield (910.0 q ha⁻¹), dry fodder yield (216.4 q ha⁻¹) compared to two cuts.

Conclusion: Variety TSFB 15-8 with nitrogen level of 120 kg N ha⁻¹ at C₂ found suitable and economical for cultivation in southern Telangana zone.

Keywords: Varieties, nitrogen levels and cutting management.

1. INTRODUCTION

Pearl millet (*Pennisetum glaucum*. L) is an important minor millet being cultivated for high dietary fiber and nutrient source for human beings and also a good fodder crop for livestock and requires less irrigation water as compared to other cereal fodders; hence, it can be

cultivated in command areas under rice fallows, where water is very scarce. It is an ideal crop with high tillering ability, high dry matter production, high protein content (10-12%) with excellent growth habit, high palatability and better nutritive value. The dual-purpose nature of the pearl millet has been recently identified due to its profuse tillering, withstanding capacity for repeated harvesting and absence of anti-nutritional factors like prussic acid, better performance under marginal and low fertile soils. Pearl millet is quick growing cereal as compared to maize and sorghum and it produces green fodder in short relation or time. Being tall vigorous with exceptional fodder yielding potential, it is an indispensable fodder for the animal inhabitants in arid and semi-arid regions of the world. In India, the requirement of green fodder was 611.99 Mt against the availability of only 224.08 Mt (Anonymous. 2006). It reflects a wide gap between demand and supply because the regional deficit are more important than the national deficit especially for forages which account only 3.4 m ha and 9.3 m ha area of total cultivable land is utilized for forage production in 1999 in Rajasthan and India, respectively (GOI, 2004). Therefore, the development of quality fodder cultivars and management to meet out the fodder requirements forever increasing livestock population is imperative, as the quality of fodder is very important issue with respect to livestock health status as well as to maximize the animal production.

2. MATERIAL AND METHODS

Field experiment was conducted during *kharif* season of 2021 at AICRP on Forage Crops and Utilization, ARI, PJTSAU, Rajendranagar, Hyderabad. The present research work is framed with an objective to identify an ideal pearl millet variety under ideal nitrogen level and cutting management for high green and dry fodder yield. The experiment was laid out in randomized block design with factorial concept. The soil was sandy loam in texture with pH of 7.0 low in available nitrogen, medium in available phosphorus and available potassium. The experiment consisted of 18 treatment combinations viz., three varieties (TSFB 15-4, TSFB 15-8 and Moti bajra), three nitrogen levels (80, 100 and 120 kg N ha⁻¹) and two cutting management practices (C₁: Two cuts: 1st at 60 DAS, 2nd cut at 50% flowering) (C₂: Three cuts: 1st at 50 DAS, 2nd cut at 35 days after 1st cut and 3rd cut at 50% flowering) with three replications, phosphorus (40 kg P₂O₅ ha⁻¹) and potassium (20 kg K₂O ha⁻¹) were applied through Di-ammonium phosphate and Muriate of potash respectively. Nitrogen was given in split-doses, half as basal and other dose was given after 30 DAS and after each cut as top dressing. Crop was sown at a row distance of 30 cm. The crop was sown during 2nd week of July and harvested when crop attained 50% flowering. Five plants were randomly select in each net plot area for taking observations on growth and yield attributing parameters. The samples were first dried under shade and then in electric oven at temperature of 60°C till attaining constant weight, on the basis of weight of these samples, the green fodder yield was converted into dry matter yield (q ha⁻¹). Data of growth attributes that is plant height (cm), number of tillers m⁻¹ row length, leaf-stem ratio and forage yield as green and dry (q ha⁻¹). Data obtained were statistically analyzed as mentioned by Gomez and Gomez, 1984.

3. RESULTS AND DISCUSSION

3.1 Growth parameters

3.1.1 Varieties

The growth and fodder yield of pearl millet significantly varied among varieties (Table 1). The pearl millet variety TSFB 15-8 recorded significantly higher plant height (67.2 cm) and

leaf-stem ratio (0.4) as over other varieties but the maximum number of tillers was found in Moti bajra (4.5) and was at par with TSFB 15-8. The TSFB 15-8 recorded highest green fodder yield of 821.2 q ha⁻¹ which was followed by TSFB 15-4 (757.6 q ha⁻¹) and the same variety recorded higher dry matter yield of 198.4 q ha⁻¹ followed by variety Moti bajra (181.7 q ha⁻¹). This was due to the superiority of the genotype to produce more values of growth characters like plant height, leaf-stem ratio, number of tillers.

3.1.2 Nitrogen levels

Application of nitrogen 120 kg N ha⁻¹ recorded significantly higher green fodder yield (817.7 q ha⁻¹) and was at par with 100 kg N ha⁻¹ (787.9 q ha⁻¹) and dry matter yield (201.0 q ha⁻¹) over other levels. Increase in green bio-mass yield was due to improved growth parameters viz., plant height (72.0 cm), number of tillers m⁻² (4.7) and leaf-stem ratio (0.4). Nitrogen is directly involved in cell division, cell elongation, formation of nucleotides and co-enzymes that increased meristematic activity, since nitrogen is an integral part of chlorophyll, plays an important role in photosynthesis and produce more of photosynthates, which helped in accumulation more dry matter yield. This is in conformity with the findings of Golada et al. [6] and Damame et al. [4].

3.1.3 Cutting management

Results with cutting management indicated that the plant height (65.9 cm), number of tillers m⁻² (5.9) were recorded with three cuts for green fodder and recorded significantly higher green fodder yield (910.0 q ha⁻¹) and dry matter yield (216.4 q ha⁻¹) as compared to two cuts of green fodder yield (641.1 q ha⁻¹) and dry matter yield (151.4 q ha⁻¹). This might be due to high production potential of all the varieties and higher number of tillers m⁻² as evident from data (Table 1).

3.1.4 Interaction effect

The interaction between different varieties and nitrogen levels was found to be significant and revealed that variety TSFB 15-8 recorded maximum green fodder yield at 100 kg N ha⁻¹ (860.1 q ha⁻¹) but significantly on par with Moti bajra at 80 kg N ha⁻¹ and 120 kg N ha⁻¹. Contrary to this, at 100 kg N ha⁻¹ TSFB 15-8 was significantly on par with TSFB 15-4 (Table 2).

The interaction between different varieties and nitrogen levels on dry fodder yield was found to be significant and it was observed that TSFB 15-8 recorded maximum dry fodder yield at 100 kg N ha⁻¹ (202.9 q ha⁻¹) and 120 kg N ha⁻¹ (216.2 q ha⁻¹) but significantly on par with TSFB 15-4 at 100 kg N ha⁻¹ and with Moti bajra at 120 kg N ha⁻¹. Contrary to this, at 80 kg N ha⁻¹ Moti bajra (176.4 q ha⁻¹) was found to be superior and was on par with TSFB 15-8 (Table 3).

Table 1: Effect of different varieties, nitrogen levels and cutting management on growth parameters and fodder yield of pearl millet.

Treatments	Plant height (cm)	No. of tillers m ⁻²	L:S ratio	GFY (q ha ⁻¹)	DFY (q ha ⁻¹)
Varieties					
TSFB 15-4	63.6	3.7	0.3	757.6	171.6
TSFB 15-8	67.2	4.4	0.4	821.2	198.4
Moti bajra	60.2	4.5	0.4	747.7	181.7
S Em±	0.9	0.09	0.01	14.9	4.4

CD (P=0.05)	2.7	0.2	0.04	42.7	12.6
Nitrogen levels (kg ha⁻¹)					
80	56.6	3.9	0.4	720.9	167.5
100	62.4	3.9	0.3	787.9	183.2
120	72.0	4.7	0.4	817.7	201.0
S Em±	0.9	0.09	0.01	14.9	4.4
CD (P=0.05)	2.7	0.2	0.04	42.7	12.6
Cutting management					
C ₁ (Two cuts)	61.5	3.3	0.3	641.1	151.4
C ₂ (Three cuts)	65.9	5.1	0.5	910.0	216.4
S Em±	0.8	0.07	0.01	12.1	3.6
CD (P=0.05)	2.2	0.2	0.03	34.9	10.3

NOTE: L:S- Leaf-stem ratio, GFY- Green fodder yield, DFY- Dry fodder yield.

Table 2. Interaction effect of different varieties and nitrogen levels on green fodder yield of fodder pearl millet.

Treatments	Nitrogen levels (kg ha ⁻¹)		
	80	100	120
Varieties			
TSFB 15-4	674.2	807.6	791.1
TSFB 15-8	767.9	860.1	835.8
Moti Bajra	720.7	696.1	826.2

Table 3. Interaction effect of different varieties and nitrogen levels on dry fodder yield of fodder pearl millet

Treatments	Nitrogen levels (kg ha ⁻¹)		
	80	100	120
Varieties			
TSFB 15-4	150.2	181.6	183.1
TSFB 15-8	176.0	202.9	216.2
Moti Bajra	176.4	165.1	203.7

Table 4. Effect of different varieties, nitrogen levels and cutting management on economics of fodder pearl millet

Treatments	Gross returns (Rs. ha ⁻¹)	Net returns (Rs. ha ⁻¹)	B-C ratio
Varieties			
TSFB 15-4	132584	92640	2.3
TSFB 15-8	143713	103768	2.5
Moti bajra	130842	90897	2.2

Nitrogen levels (kg ha ⁻¹)			
80	126162	90897	2.1
100	137882	97938	2.4
120	143095	103024	2.5
Cutting management			
C1 (Two cuts)	112186	73574	1.9
C2 (Three cuts)	159240	117963	2.8




Fig.1. Variation in GFY and DFY (q ha⁻¹) among varieties, nitrogen levels and cutting management.

NOTE: Varieties (TSFB 15-4, TSFB 15-8, Moti bajra)

Nitrogen levels (80, 100, 120 kg ha⁻¹)

Cutting management (C₁ and C₂)

GFY- Green fodder yield, DFY- Dry fodder yield.

Among varieties tested TSFB 15-8 registered higher net returns (103768 Rs. ha⁻¹) and benefit-cost ratio (2.5). Application of nitrogen 120 kg N ha⁻¹ recorded higher net returns (103024 Rs. ha⁻¹) and benefit-cost ratio (2.5) and among cutting management practices C₂ (Three cuts) recorded higher net returns (Rs. 117963 ha⁻¹) and benefit-cost ratio (2.8). This might be due to better growth attributes which resulted higher green forage yield with higher level of nitrogen.

4. CONCLUSION

Based on the research results it can be inferred that forage pearl millet variety TSFB 15-8 with nitrogen level of 120 kg N ha⁻¹ at C₂ (Three cuts) found suitable and economical for cultivation in Southern Telangana Zone. These results are in conformity with findings of Shekara et al. [16] and Shekara et al. [17].

REFERENCES

1. Aboelgoud SA, Ragab MN. Performance of Two Fodder Pearl Millet Varieties under Different Seeding and Nitrogen Fertilizer Rates. *Journal of Plant Production*. 2021;12(10):1069-1076.
2. Anonymous, 2006. Hand book of Agriculture, ICAR, New Delhi. pp. 1353-1355.
3. Ayub M, Nadeem MA, Tahir M, Ibrahim M, Aslam MN. Effect of nitrogen application and harvesting intervals on forage yield and quality of pearl millet (*Pennisetum americanum* L.). *Pak. J. Life Soc. Sci.* 2009;7(2):185-189.
4. Damame SV, Bhingarde RN, Pathan SH. Effect of different nitrogen levels on nutritional quality and nitrate nitrogen accumulation in forage pearl millet genotypes grown under rainfed conditions. *Forage Research*. 2013;39(2): 93-95.
5. GOI, 2004. Livestock Census Report. Directorate of Economics and Statistics, Ministry of Agriculture, Government of India, New Delhi.
6. Golada SL, Patel BJ, Singh P, Buldak LR. Effect of integrated nutrient management on productivity of forage pearl millet (*Pennisetum glaucum* L.). *Forage research*. 2010;36(3): 185-187.
7. Joshi MP, Pankhaniya RM, Mohammadi NK. Response of pearl millet (*Pennisetum glaucum* L.) to levels and scheduling of nitrogen under south Gujarat condition. *International Journal of Chemical Studies*. 2018; 6(1):32-35.
8. Kadam SB, Pawar SB, Jakkawad SR. Response of pearl millet (*Pennisetum glaucum* L.) to levels and scheduling of nitrogen under Maharashtra condition. *J. Pharm. Phytochem.* 2019;8: 2922-2925.
9. Khairwal IS, Yadav SK, Rai KN, Upadhyaya HD, Kachhawa D, Nirwan B, Bhattacharjee R, Rajpurohit BS, Dangaria CJ. Evaluation and identification of promising pearl millet germplasm for grain and fodder traits. *Journal of SAT Agricultural Research*. 2007;5(1):1-6.
10. Kumar ARUN, Kumar ANIL, Midha LK, Duhan BS. Effect of different production factors on productivity, NPK uptake and quality of pearl millet [*Pennisetum glaucum* (L.) R. BR.]. *Forage Research*. 2015;41: 46-49.
11. Kumar DS. Green fodder yield and quality of dual-purpose pearl millet (*Pennisetum glaucum* L.) varieties as influenced by cutting and nitrogen management. *An International Quarterly Journal of Life Sciences*. 2016;11(4): 2311-2315.

12. Makarana G, Yadav RK, Kumar R, Kumar A, Soni PG, Kar S, Rajvaidya SK. Fodder and grain quality of Pearl millet (*Pennisetum glaucum* L.) under cutting management in saline irrigation water. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(3):1251-1257.
13. Midha LK, Arya S, Kumari P, Joshi UN. Performance of forage pearl millet genotypes under different nitrogen levels. *Forage Research*. 2015;41(2):137-138.
14. Noor MA, Fiaz S, Nawaz A, Nawaz MM. The effects of cutting interval on agro-qualitative traits of different millet (*Pennisetum americanum* L.) cultivars. *Journal of the Saudi Society of Agricultural Sciences*. 2018;17(3):317-322.
15. Obeng E, Cebert E, Singh BP, Ward R, Nyochembeng L.M, Mays DA. Growth and grain yield of pearl millet (*Pennisetum glaucum*) genotypes at different levels of nitrogen fertilization in the southeastern United States. *Journal of Agricultural Science*. 2012;4(12):155.
16. Shekara BG, Lohithaswa HC. Fodder and seed yield of forage pearl millet genotypes as influenced by different levels of nitrogen. *Forage Research*. 2009;35(1): 45-47.
17. Shekara BG, Mahadevu P, Chikkarugi NM, Manasa N. Response of pearl millet (*Pennisetum glaucum* L.) varieties to nitrogen levels for higher green forage yield and quality in southern dry zone of Karnataka. *Forage Research*. 2019;45(3): 232-234.

UNDER PEER REVIEW