

Original Research Article

INFLUENCE OF PHOSPHOROUS AND FOLIAR SPRAY OF ZINC ON GROWTH AND YIELD OF LENTIL (*Lens culinaris* M.)

ABSTRACT

A field experiment was conducted during *Rabi* season of 2021-22, at crop research farm of Department of Agronomy at Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj in North Eastern plains of Eastern Uttar Pradesh with the objective to study the Influence of Phosphorus and Foliar Spray of Zinc on Growth and yield of lentil (*Lens culinaris* M.) Var. KSL-0903 was conducted on Randomized block design comprising of 9 treatments of which treatments (T₁-T₉) with different combination of Phosphorus and Foliar Spray of Zinc on which are replicated thrice. The experimental results revealed that [Phosphorus 50kg/ha+ Zinc 0.75%] was recorded maximum plant height (54.53cm), Number of nodules/plant (8.66), Plant dry weight (12.73g), number of pods/plant (163.3), number of seeds/pod (2.00), test weight (25.13g), Seed yield (1.908t/ha), Stover yield (3.534 t/ha). The treatment 9 [Phosphorus 50kg/ha+ Zinc 0.75%] recorded highest net return (66348.30 INR /ha), highest gross return (97308.30 INR/ha) and benefit: cost ratio (2.14).

Key words - Lentil, Phosphorus, Zinc, Growth parameters, Yield attributes and Economics,.

INTRODUCTION

India is largest pulse-growing country which accounts for nearly one-third of the total world area under pulses and one-fourth of the total world production. Lentil (*Lens culinaris* Medikus) is an important *Rabi* season legume crop in India, lentils are drought tolerant crop. It belongs to the sub-family Papilionaceae under the family Fabaceae (Leguminosae). Lentil (*Lens culinaris* M.) Medik a name given by the German botanist medikus in 1778 (**Cubero, 1981**). It is locally known as Masoor. Lentil requires cold climate it is very hardy and can tolerate frost and severe winter to a great extent. It requires cold temperature at the time of maturity. The optimum temperature for growth is 18-30^o C. It grows on a wide range of soils ranging from light loam sand to heavy clay in northern parts, and in moderately deep, light black soil in Madhya Pradesh and Maharashtra. It is one of the important pulse crops of India which can adapt well to a wide range of climate and soil condition. It can tolerate frost & severe winter. India ranked first in area and second in the production with 39.79% and 22.79% of world area and production respectively. Canada rank first in production (41.16%) due to very high level of productivity (1633 kg/ha) as compared to India (611 kg/ha). (**FAO STAT 2014**).

Phosphorous is one of the most important elements significantly effecting the plant growth and metabolism. In legumes, the high requirement for P is consistent with the involvement of P in high rates of energy transfer that take place in the nodule. Phosphorous has also an enhancing impact on growth and biological yield through its importance as energy storage and transfer necessary for metabolic processes (**Singh et**

al., 2014). Phosphorous addition increases the efficiency of plants to photosynthesis, enhances the activity of rhizobia. Phosphorus contributes substantially to increased yield of legumes by enhancing the physiological functions of the crop plants, root development, and nodulation. Phosphorus application not only increases the dry matter and seed yield of lentil but also enhances the N and P content of the seed by increasing nodulation and root development.

Zinc is an important component of various enzymes that are responsible for driving many metabolic reactions in all crops. It contributes to a number of important plant functions such as growth hormone production, plant resistance against disease, photosynthesis. Zinc is very important in reproductive phase like fertilization and pollen grain formation as pollen grains contains a high amount of zinc, most of the zinc is translocated to seeds during fertilization (**Abid Ali et al., 2017**). Plants enzymes activated by Zn are involved in carbohydrate metabolism, and regulation of auxin synthesis and pollen formation. Zn seems to affect the capacity for water uptake and transport in plants and also reduce the adverse effects of short periods of heat and salt stress. As Zn is required for the synthesis of tryptophan which is a precursor of IAA, it also has an active role in the production of an essential growth hormone auxin (**Alloway et al., 2008**).

MATERIALS AND METHODS

The experiment was carried out during winter season of 2021-2022 at the Crop Research Farm SHUATS, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh. The crop Research Farm is situated at 25°24' 42" N latitude, 81° 50' 56" E longitude and at an altitude of 98m above mean sea level (MSL). To assess the influence of phosphorus and Foliar Spray of zinc on growth and yield of Lentil (*Lens culinaris* M.). The experiment was laid out in Randomized Block Design comprising of 9 treatments which are replicated thrice. Each treatment net plot size is 3m × 3m. The treatment are categorized as with recommended dose of nitrogen through Urea and potash through Muriate of Potash, in addition with Phosphorus through Single super phosphate and Foliar Spray of zinc when applied in combinations as follows; T1- Phosphorus 30 kg/ha+ Zinc 0.25%; T2- Phosphorus 30 kg/ha + Zinc 0.5%; T3- Phosphorus 30 kg/ha + Zinc 0.75%; T4- Phosphorus 40 kg/ha + Zinc 0.25%; T5- Phosphorus 40 kg/ha + Zinc 0.5%; T6- Phosphorus 40 kg/ha + Zinc 0.75%; T7- Phosphorus 50 kg/ha + Zinc 0.25%; T8- Phosphorus 50 kg/ha + Zinc 0.5%; T9- Phosphorus 50 kg/ha + Zinc 0.75%. The Lentil crop was harvested treatment wise at harvesting maturity stage. Growth parameters viz. plant height (cm), dry matter accumulation(g /plant) were recorded manually on five randomly selected representative plants from each plot of each replication separately and after harvesting, seeds were separated from each net plot and were dried under sun for three days. Later winnowed, cleaned and grain yield per ha was computed and expressed in tonnes per hectare. After complete drying under sun for 10 days stover yield from each net plot was recorded and expressed in kg/hectare. The data was computed and analyzed by following statistical method of Gomez and Gomez (1976).

RESULTS AND DISCUSSION

Growth parameters:

Plant height (cm)

It is evident from Table 1. that plant height measured increased with advancement in crop growth. At harvest the treatment T₉ application of (Phosphorus 50kg/ha+ Zinc 0.75%) recorded maximum height of 54.43cm. At harvest, significantly higher plant height (54.43 cm) was recorded in treatment-9 (Phosphorus 50 kg/ha+ Zinc 0.75%). However, treatment- 8(Phosphorus 50 kg/ha + Zinc 0.5%) was statistically at par with treatment-9(Phosphorus 50 kg/ha+ Zinc 0.75%). The increase in plant height may be owing to phosphorus application in the soil might be due to increase availability and uptake of soil nutrients by the crop contributed by phosphorus fertilization. The higher availability of nutrients might have increased the photosynthetic ability and translocation of the metabolites to different parts which ultimately increased the root and shoot development of the crop. These findings corroborate the results of **Tophia Yumnam et al., (2018)**, **Singh and Singh (2014)** in lentil.

Number of Nodules/plants

At harvest, significantly higher number of nodules per plant (8.66) was recorded in treatment-9 (Phosphorus 50 kg/ha+ Zinc 0.75%). However, treatment- 8(Phosphorus 50 kg/ha + Zinc 0.5%) was found to be statistically at par with treatment-9(Phosphorus 50 kg/ha+ Zinc 0.75%). Application of phosphorus help in efficient utilization of nutrients, which resulted in attaining better crop canopy and further increased absorption and utilization of radiant energy resulting in higher effective and total number of nodules per plant. These results are confirmed with the findings of other researcher **Togay et al., (2008)**, **Patel et al., (2017)** and **Ashish Masih et al., (2020)** who reported that application of phosphorus increased the number of nodules per plant.

Plant dry weight

At Harvest highest plant dry weight was found in treatment Phosphorus 50 kg/ha+ Zinc 0.75% (12.73g). However, treatment- 8 (Phosphorus 50 kg/ha + Zinc 0.5%) was found to be statistically at par with treatment-9(Phosphorus 50 kg/ha+ Zinc 0.75%). Increase in dry weight due to increase in photosynthetic ability and translocation of metabolites to different parts which ultimately increased the shoot development of the crop. Similar results were found in **Tophia et al., (2018)**.

Table 1. Influence of Phosphorous and Foliar spray of zinc on growth parameters of lentil.

Treatments	No of Pods/ plant	No of Seeds/pod	Test weight (g)
Phosphorus 30 kg/ha + Zinc 0.25%	139.8	1.00	21.82
Phosphorus 30 kg/ha + Zinc 0.5%	142.1	1.30	22.10
Phosphorus 30 kg/ha + Zinc 0.75%	143.1	1.60	22.90
Phosphorus 40 kg/ha + Zinc 0.25%	149.2	1.40	22.63
Phosphorus 40 kg/ha+ Zinc 0.5%	150.8	1.70	22.93
Phosphorus 40 kg/ha + Zinc 0.75%	155.1	1.70	23.73
Phosphorus 50 kg/ha + Zinc 0.25%	158.4	1.60	22.77
Phosphorus 50 kg/ha + Zinc 0.5%	161.0	1.80	23.87
Phosphorus 50 kg/ha+ Zinc 0.75%	163.3	2.00	25.13
F-Test	S	S	S
SEm±	3.63	0.07	0.60
CD (P=0.05)	2.59	0.20	1.80

Yield Attributes:

Number of Pods/plant

The statistically higher number of pods/plant (163.3) was observed in treatment-9 (Phosphorus 50 kg/ha+ Zinc 0.75%), which was significantly higher over rest of the treatments and treatment-8 (Phosphorus 50 kg/ha+ Zinc 0.5%), was statistically at par with treatment-9(Phosphorus 50 kg/ha+ Zinc 0.75%). It might be the reason of moderate plant nutrients availability due to which the plant produces more pods plant⁻¹ as compare to other treatments and also phosphorus strongly increases the reproduction of the plants i.e., flowering and fruiting. These results were similar with that of **Abid et al., (2017)** and **Choubey et al., (2013)**.

Number of Seeds/Pod

Significant effect was observed by the statistical analysis of Number of seeds/pod. Treatment Phosphorus 50 kg/ha+ Zinc 0.75% recorded significant and highest number of seeds/pod (2.00). However, statistically at par with treatment-8 (Phosphorus 50 kg/ha+ Zinc 0.5%).

Test weight (g)

The highest test weight (25.13 g) was observed in treatment-9 with (Phosphorus 50 kg/ha+ Zinc 0.75%), which was found to be significantly higher over rest of the treatments statistically at par with treatment-8 (Phosphorus 50 kg/ha+ Zinc 0.5%). Phosphorus fertilization was effective in the generative period of the plant and that such fertilization led to a much better granulation, enhancement and improvement occurred in test weight with the increasing doses of phosphorus. Similar findings were reported by **Dogan et al., (2014)**.

Seed yield (kg/ha)

The highest seed yield (1908 kg/ha) was observed in treatment-9 with (Phosphorus 50 kg/ha+ Zinc 0.75%), which was found to be significantly higher over rest of the treatments statistically at par with treatment-8 (Phosphorus 50 kg/ha+ Zinc 0.5%). Application of phosphatic fertilizer therefore provided balance nutrition to the crop which resulted in higher seed yield of lentil. Phosphorus also increased the photosynthesis and translocation of assimilates to different plant parts for enhanced growth and yield attributing characters of the crop as observed in number of pods per plant and number of seeds per pod. In the later stage, the excess assimilates stored in the leaves was translocated towards sink development which ultimately contributed to higher seed yield. These findings were supported by **Choubey et al., (2013)**, **Dharvendra Singh (2017)** and **Tophia Yumnam et al., (2018)** in lentil.

Stover yield (kg/ha)

The highest stover yield (3534 kg/ha) was observed in treatment-9 with (Phosphorus 50 kg/ha+ Zinc 0.75%), which was significantly higher over rest of the treatments statistically at par with treatment-8 (Phosphorus 50 kg/ha+ Zinc 0.5%). Each increment in P stimulates the lentil to increase

the straw yield. Similar findings were reported by **Choubey et al (2013)**.

Economics:

Gross returns (INR/ha)

Gross returns (97308.30 INR/ha) were found to be highest in treatment-9 (Phosphorus 50 kg/ha+ Zinc 0.75%) and the lowest gross returns (68680.00 INR/ha) was found to be in treatment-1 (Phosphorus 30 kg/ha + Zinc 0.25%) as compared to other treatments.

Net returns (INR/ha)

Net (66348.30 INR/ha) were found to be highest in treatment-9 (Phosphorus 50 kg/ha+ Zinc 0.75%) and the lowest net returns (39255.00 INR/ha) was found to be in treatment-1 (Phosphorus 30 kg/ha + Zinc 0.25%) as compared to other treatments.

The statistically higher Net returns was with the application of treatment -9 (Phosphorous 50kg/ha+ Zinc 0.75%). With increasing levels of phosphorus, the grain and straw yield increased this attributed to a higher net return. These results are in conformity with those observed by **Mitra et al., (2006)**.

Benefit Cost ratio (B: C)

Benefit Cost ratio (2.14) was found to be highest in treatment-9 (Phosphorus 50 kg/ha+ Zinc 0.75%) and the lowest benefit cost ratio (1.33) was found to be in in treatment-1 (Phosphorus 30 kg/ha + Zinc 0.25%) as compared to other treatments.

The statistically higher Benefit cost ratio was with the application of treatment-9 (Phosphorus 50 kg/ha+ Zinc 0.75%), due to the phosphorus and zinc provides conclusive condition to the soil with the synergistic influence of Phosphorus and Foliar spray of zinc resulting better benefit cost ratio. These results are supported by the findings of **Mitra et al., (2006)** in lentil

Treatments	No of Pods/ plant	No of Seeds/pod	Test weight (g)	Seed yield (t/ha)	Stover yield (t/ha)	Harvest index (%)
Phosphorus 30 kg/ha + Zinc 0.25%	139.8	1.00	21.82	1.346	2.070	39.38
Phosphorus 30 kg/ha + Zinc 0.5%	142.1	1.30	22.10	1.395	2.274	38.02

Table 2. Influence of Phosphorous and foliar spray of zinc on yield attributes of lentil.

Phosphorus 30 kg/ha + Zinc 0.75%	143.1	1.60	22.90	1.523	2.486	38.03
Phosphorus 40 kg/ha + Zinc 0.25%	149.2	1.40	22.63	1.446	2.324	38.36
Phosphorus 40 kg/ha+ Zinc 0.5%	150.8	1.70	22.93	1.564	2.806	35.77
Phosphorus 40 kg/ha + Zinc 0.75%	155.1	1.70	23.73	1.698	2.961	36.68
Phosphorus 50 kg/ha + Zinc 0.25%	158.4	1.60	22.77	1.468	2.471	37.22
Phosphorus 50 kg/ha + Zinc 0.5%	161.0	1.80	23.87	1.814	3.369	37.82
Phosphorus 50 kg/ha+ Zinc 0.75%	163.3	2.00	25.13	1.908	3.534	35.04
F-Test	S	S	S	S	S	NS
SEm±	3.63	0.07	0.60	45.03	116.70	1.78
CD (P=0.05)	2.59	0.20	1.80	135.00	349.87	-----

Table 3. . Influence of Phosphorus and Foliar spray of zinc levels on Economics of lentil.

Treatments	Total cost of cultivation (INR/ha)	Gross returns (INR/ha)	Net returns (INR/ha)	B:C ratio
Phosphorus 30 kg/ha + Zinc 0.25%	29425.00	68680.17	39255.17	1.33
Phosphorus 30 kg/ha + Zinc 0.5%	29675.00	71145.00	41470.00	1.40
Phosphorus 30 kg/ha + Zinc 0.75%	29925.00	77673.00	47748.00	1.60
Phosphorus 40 kg/ha + Zinc 0.25%	29942.00	73762.83	43820.83	1.46
Phosphorus 40 kg/ha+ Zinc 0.5%	30192.00	79764.00	49572.00	1.64
Phosphorus 40 kg/ha + Zinc 0.75%	30442.00	86598.00	56156.00	1.84
Phosphorus 50 kg/ha + Zinc 0.25%	30460.00	74868.00	44408.00	1.46
Phosphorus 50 kg/ha + Zinc 0.5%	30710.00	92683.83	61973.83	2.02
Phosphorus 50 kg/ha+ Zinc 0.75%	30960.00	97308.00	66348.00	2.14

REFERENCES

- Abid, Ali., (2017) Effect of phosphorous and zinc on yield of lentil, *Pure Applied Biology*, **6**(4):1397-1402.
- Anil, Kumar, Singh and Bhatt, B.P (2015) Late-Sown lentil performance in response to foliar application of zinc, *Bangladesh Journal of Botony*. **44**(1): 125-128.
- Biswas, P. K (2015) Conjoint application of bio fertilizer and phosphorous levels on growth, nodulation, nutrient uptake and productivity of lentil (*Lens culinaris* Medikus) in red and lateritic soils of west Bengal, *Journal Crop and Weed*, **11**(1):29-32
- Cubero, J.I (1981) Origin, taxonomy and domestication In: Lentils (eds C.Webb & G.C. Hawtin). *Commonwealth Agricultural Bureau*, Slough, England,15-38
- Choubey, S.K., Dwivedi, V.P and Srivastava, N.K (2013) Effect of different levels of phosphorus and sulphur on growth, yield and quality of lentil (*Lens culinaris* M). *Indian Journal. Science Research*. **4**: 149- 150.
- Datta, S.K., (2013) Effect of variety and level of phosphorous on the yield and yield components of lentil, *International Journal of Research Innovation & Technology* **3**(1): 78-82.
- Deo, Karan (2014) Effect of zinc and boron application on yield of lentil and nutrient balance in the soil under Indo-Gangetic plain zones, *Journal of Agriculture research* **1**(4): 206-209.
- Mukesh, Kumar., Pandey (2016) Effect of integrated phosphorous management on growth, yield and quality of lentil (*Lens culinaris*), *Indian Journal Research*, **50** (3): 238-243.
- Patel, M.M (2011) Effect of zinc and iron on yield and yield attributes of rainfed cowpea (*Vigna unguiculata* L.Walp), *Annals of Arid Zone* **50**(1):17-19.
- Shahid, Rasool (2016) Effect of Bio fertilizers and phosphorous on growth and yield of lentil, *International Journal of Advances in Agricultural Science and Technology*, **3**(7), pg 35-42.
- Sonet, R.A (2020) Influence of phosphorous levels on growth and yield of four lentil varieties, *Bangladesh Agronomy Journal*, **23**(1): 29-36.
- Tophia, Yummam (2018) Influence of phosphorous on growth and yield of promising varieties of lentil (*Lens Culinaris* .M), *International Journal of Current Microbiology and Applied Sciences*.**7**(08):162-170
- Verma, C.B (2017) Enhancing growth, yield and quality of lentil through foliar spray of zinc, urea and

thiourea under rainfed condition, *Agri ways* 5(2): 123-127.

Yesim, Togay., Necat, Togay and Yusuf, Dogan (2008) Research on the effect of phosphorus and molybdenum applications on the yield and yield parameters in lentil ISSN 1684–5315 *Academic Journals*

Zeidan, M.S (2007) Effect of organic manure and phosphorous fertilizers on growth, yield and quality of lentil in sandy soil. *Research Journal of Agriculture and Biological Sciences*, 3(6): 748-752.

UNDER PEER REVIEW