

# Exploring the genetic variability for yield attributing traits among the Indigenous and Exotic collection of wheat in Cis-Himalayan region of West Bengal

## Abstract

Field experiment was conducted in Instructional farm, Uttar Banga Krishi Viswavidyalaya, during 2018-19 rabi season to study the genetic components and genetic variability based on eleven number of quantitative traits in 254 bread wheat lines, which is collected from NBPGR, New Delhi. Analysis of variance for treatment showed high significance for all the traits except grain yield  $\text{polt}^{-1}$ . Estimated GCV and PCV were high for awn length, peduncle length, plant height and AUDPC. Plant height (97.83), awn length (95.49), peduncle length (94.88), test weight (90.98), Days to 50% flowering (87.66), number of grains  $\text{spike}^{-1}$  (70.33) and Spike length (62.3) were showed high heritability along with high degree of genetic advance. Finally, 254 wheat accession were grouped into three cluster based on the dendrogram analysis using ward method.

**Key word:** Wheat, Genetic components, Genetic variability and Ward method.

## Introduction

Wheat is the most widely grown crop and staple food across the world, Wheat plays important role in human nutrition. It provides 20% of the total calories (Goel *et al*, 2018) for millions of individuals who rely on wheat-based diets. It provides 40% of their dietary intake of crucial micronutrients such as zinc, iron, manganese, magnesium, and vitamins B and E (Velu *et al*, 2017). Wheat productivity must increase by more than 50% over existing levels by 2050 to meet demand as the human population continues to grow (Tadesse *et al*, 2019). Breeders face challenges in increasing current output levels of wheat production as the growing population of world require higher amount of production than is currently available (Rahman *et al*, 2021). It is not possible to expand the production area, hence, the only option is to boost the productivity by developing superior varieties and improving the agronomic production management in order to meet the rising food demands (Pradhan *et al*, 2015).

Plant breeding with the help of the information of quantitative genetics is considered as an active programme and selection is vastly effective when it depends on the presence of genetic variation within the accessions. Breeding for crop improvement is based on genetic richness and selection of suitable genotypes. Genetic variability study in plant is the significant argument and provides superlative resolution for plant breeders to develop novel improved genotypes with desired traits (Govindaraj *et al*, 2015). Before selecting the most desirable genotypes, breeders want to assess genetic components such as phenotypic coefficient of variation, genotypic coefficient of variation, environmental coefficient of variation, heritability, genetic advance etc. which are essential to estimate the genetic richness and superiority of any particular genotype (Atinafu *et al*, 2020). It also helps in determining the effectiveness of selection for a particular trait in that genotype to get improved cultivars (Atinafu *et al*, 2020).

The present study was conducted at Cis-Himalayan plains of West Bengal which is located in the terai region where rice is the predominant crop with the rice-wheat cropping system being prevalent in adjoining areas with rice prevailing during the monsoon season and wheat in the rabi season. The major problems for wheat cultivation in this region include short winter, early onset of monsoon, micro nutrient deficiency along with prevalence of foliar diseases such as spot blotch etc. To combat these issues the present study was undertaken with 254 wheat genotypes of both exotic and indigenous collection from NBPGR to study the genetic variability along with estimation of genetic components for selection of suitable genotypes in this region.

## **Martials and Methods**

Field research was conducted in Instructional Farm, Uttar Banga Krishi Viswavidyalaya, Pundibari, Coochbehar, West Bengal, during rabi season of 2018-19, for morphological characterization and screening for spot blotch of 254 wheat accessions. The experimental field is situated at 26° 24' 16.11" N latitude, 89° 23' 1.69" E longitude and at an altitude of 43 m above the mean sea level. The wheat collections were consisted with two hundred indigenous accessions and fifty-four exotic accessions which were collected from ICAR-National Bureau of Plant Genetic Resources, New Delhi, India. All the genotypes were evaluated in Augmented Block Design with four numbers of checks cultivars like HD 2967, BHU 35 and UBW 9 (highly resistant to moderately resistant to spot blotch) and Sonalika (susceptible to spot blotch disease). Each genotype was grown in three rows in 100 × 75 cm<sup>2</sup> plot size. The recommended cultural and agronomic practices were implemented to raise good crop (Mondal *et al.*, 2018). Field screening for spot blotch disease was done by creating disease pressure using Sonalika (susceptible check) in throughout the border. AUDPC and rAUDPC were calculated according to Chattopadhyay *et al.*, (2021). All the data were recorded for 11 morphological characters viz., flag leaf length (FL) and width (FW), days to 50 % heading (DF), plant height (PH), spike length (SL), peduncle length (PL), number of grains spike<sup>-1</sup> (NG), awn length (AL), thousand grain weight (TW), grain yield (YL) along with rAUDPC (AU) for spot blotch disease. All the statistical analysis were done by using "augmentedRCBD" package in R v. 3.6.0 software.

## **Result and discussion**

### *Mean performances and ANOVA*

The average days to 50% flowering were 77.65 days which varied from 57.13 (IC574387) to 120.38 (IC542063) days. Among the 254 genotypes, EC534432, IC527929, IC28622, IC28872, IC28554, IC252742 and IC335683 were showed early flowering. The plant height was ranged from 65.3 cm (IC576640) to 172.55 cm (EC576578) with average value 102.88. Some short height genotypes (IC530005, EC575981, IC416043, IC415870 and IC530086) were observed within the 254 accessions which are important for making dwarf population in breeding program. The average flag leaf length was 23.43 cm and it was varied from 14.35 to 46.92 cm. The highest flag leaf length was observed in IC539313 and the genotype EC187159 was lowest flag leaf length. The mean flag leaf width was 1.72 and it's ranged from 0.98 to 2.99. Genotype IC539313 showed the highest width length whereas EC575732 showed lowest width length. Increase in leaf area of plants had increased amount of photosynthates, which increased grain yield per plant (Weraduwege *et al.*, 2015). Variability in spike length ranged from 5.95 cm to 20.95 cm with mean value of 10.68 cm. EC445374 had the lowest spike length and

highest spike length was observed in IC531862. Variability in peduncle length ranged from 4.76 cm in IC445528 to 42.9 cm in IC529196 with mean value of 20.13 cm. The awn length was ranged from 0 to 16.31 cm with average value was 6.42. It was observed that, among the 254 genotypes only nine accessions (IC529052, IC530086, IC584159, EC576591, EC577738, EC10970, IC144903, EC576578 and IC406521) were awn less and noted as 0. Number of grains spike<sup>-1</sup> ranged from 41.65 to 109.57 with an average of 71.84. The genotype IC28755 showed lowest number of grain and IC539313 showed highest number of grains spike<sup>-1</sup>, respectively. The test weight of grain was ranged from 16.88 g in IC539314 to 49.54 g in IC542076 with the value mean was 29.72 g. Grain yield plot<sup>-1</sup> varied from 48.49 g to 537.24 g with average was 264.18 g. IC393877 and IC35715 possessed minimum and maximum number of grain yield plot<sup>-1</sup>, respectively. Yield was also high for IC530058 (503.49g), IC445425 (504.99g) and IC402042 (516.74g) which are the promising genotypes for future breeding programme. The average rAUDPC for spot blotch disease was 13.61 and it varies from 4.12 to 28.32. The low rAUDPC was recorded for IC527448, EC339632, EC187159, IC539313, EC463396 whereas IC252954 recorded the highest rAUDPC. This indicated higher degree of variability for disease resistance reaction against spot blotch. The mean performance of 254 genotypes for 11 characters were represented in the supplementary table 1.

The analysis of variance of 254 genotype of wheat for 11 quantitative characters viz. awn length, days


**Table 1: Analysis of variance for eleven quantitative characters in wheat**

Source	Df	DF	PH	FL	FW	EL	PL	AL	NG	TW	YL	AU
Block	24	374.7	1303.7	8.71	0.1	3.61	80.88	3.84	626.19	33.99	23452.9	27.02
		6 **	4 **	ns	1 **	**	**	**	**	**	4 **	**
Total entries	25	131.5	329.02	17.45	0.1	4.99	31.6	6.46	192.3	39.35	9943.17	26.61
		3	3 **	**	**	**	**	**	**	**	**	ns
Check	3	1001.	913.79	163.1	1.4	82.5	134.3	10.8	1632.2	768.0	36605.5	773.7
		4 **	**	8 **	3 **	3 **	7 **	7 **	5 **	9 **	7 **	8 **
Check x Accessions	25	121.0	322 **	15.71	0.0	4.06	30.37	6.4	175.02	30.6	9623.22	17.64
		0	9 **	**	9 **	**	**	**	**	**	**	ns
Residuals	72	19	9.48	6.08	0.0	1.43	1.91	0.3	48.48	2.66	7390.89	9.41
					3							

to 50% flowering, ear length, flag leaf length, flag leaf width, number of grains spike<sup>-1</sup>, plant height, length of peduncle, test weight, grain yield along with rAUDPC of spot blotch are presented in Table 1. Mean squares due to block for all the above characters except flag leaf length were highly significant indicating the adequacy of block for statistical analysis of the characters. Mean squares due to genotypes for all above the characters except grain yield plot<sup>-1</sup> were highly significant which indicated presence of genetic variability among the experimental materials. Mean squares due to checks for all above the characters were highly significant which indicated presence of genetic variability among the check materials too whereas interaction between check and accession were significant for all the above characters except grain yield plot<sup>-1</sup> indicated differential performance of the genotypes under different crop growing conditions.

#### *Estimation of Genetic components*


Estimated genotypic and phenotypic coefficient of variation (GCV and PCV) were analysed and grouped under three categories viz. low (<10%), moderate (10-20%) and high (>20%) according to Singh and Chaudhary, 1979. On the basis of the above grouping, days to 50% flowering, flag leaf length, flag leaf width, spike length, number of grains spike<sup>-1</sup>, test weight had moderate GCV and PCV whereas, awn length, peduncle length, plant height, rAUDPC showed high of GCV and PCV. Moderate GCV and high PCV were observed for grain yield plot<sup>-1</sup> (Supplementary table 2). Similar results were reported for moderate GCV and high PCV for spike length (Gerema, 2020), for number of grains spike<sup>-1</sup> (Dhananjoy *et al*, 2012; Singh *et al*, 2013), for flag leaf area (Singh *et al*, 2013), for days to 50% flowering (Kolakar *et al*, 2012). High GCV and PCV reported for plant height (Singh *et al*, 2013), for peduncle length (Kumar *et al*, 2016) and for AUDPC (De *et al*, 2014). Elevated magnitude between GCV and PCV were observed for flag leaf length, flag leaf width, ear length, number of grains spike<sup>-1</sup>, plot yield along with rAUDPC values for spot blotch indicating higher influence of environmental factors in expression of these characters (Figure 1). Very low difference was observed for awn length, days to flowering, plant height and test weight indicating low sensitivity to environment and consequently greater role of genetic factor influencing the expression of these characters. This also revealed the effectiveness of selection for genetic improvement of those characters which have lower impact of environmental factors.



**Fig 1.** Estimated GCV and PCV of eleven quantitative traits for 254 wheat accessions

The high heritability in broad sense (Figure 2) was found for characters such as plant height (97.83), awn length (95.49), peduncle length (94.88), test weight (90.98), days to 50% flowering (87.66), number of grains spike<sup>-1</sup> (70.33), spike length (62.3) whereas, moderate to low heritability (Figure 2) was for rAUDPC (48.59) and grain yield plot<sup>-1</sup>(27.06). Similar results have been reported for plant height and test weight (Singh and Upadhyay, 2013; Devesh *et al.*, 2018; Hossain *et al*, 2021); awn length (Teich, 1984); for peduncle length


(Baranwal *et al*, 2012; Shehrawat and Kumar, 2021), days to 50% flowering (Fellahi *et al*, 2013; Baranwal *et al*, 2012), number of grains spike<sup>-1</sup> (Ullah *et al*, 2012; Fellahi *et al*, 2013, Roy *et al*, 2021), spike length (Birhanu *et al*, 2016).



**Fig 2.** Estimated broad sense heritability of eleven quantitative traits for 254 wheat accessions

Genetic advance as a percent of mean (GAM) were classified as low (<15 %), moderate (15-20 %) and high (>20 %) according to Falconer and Mackay, 1996 and represented in the Figure 3. Based on the results, GAM was high for awn length (79.51), peduncle length (59.4), plant height (41.01), test weight (34.31), rAUDPC (31.49), days 50% to flowering (28.9), number of grains spike<sup>-1</sup> (25.81), spike length (23.29), grain yield plot<sup>-1</sup> (21.27) and flag leaf length (20.84) whereas, moderate for flag leaf width (19.78). Similar results were obtained earlier for peduncle length and number of grain spike<sup>-1</sup> (Baranwal *et al*, 2012), test weight and plant height (Fellahi *et al*, 2013), grain yield plot<sup>-1</sup> (Birhanu *et al*, 2016), spike length and days to 50% flowering (Belay and Fisseha, 2021), flag leaf width (Ahmad, 2017).

High genetic advance coupled with high heritability were observed in days to 50% flowering, number of grain spike<sup>-1</sup>, test weight, awn length, peduncle length and flag leaf length which are important characters for selection of genotypes in breeding programme. These results helped in good understanding of heritability and genetic advance present in different yield contributing characters which are the first option for crop improvement (Larik *et al*, 1999).



**Fig 3.** Estimated genetic advance in percent of mean of eleven quantitative traits for 254 wheat accessions

#### *Genetic variability*

Genetic variability was detected through dendrogram analysis for 254 genotypes utilizing 11 quantitative traits. In the present investigation, dendrogram analysis was performed with four different methods (“complete”, “average”, “single”, and “ward”) and finally “ward method” was considered based on the higher agglomerative coefficient value (Table 2).

Table 2: Different methods for dendrogram analysis with their agglomerative coefficient value

Method	Agglomerative coefficient
Complete	0.85
Average	0.75
Single	0.56
Ward	0.92

According to dendrogram (Figure 4), all the wheat accession were grouped into 3 distinct clusters. Cluster I (red) consisted with 132 genotypes, whereas cluster II (green) included 82 genotypes and cluster III (black) had 40 genotypes. This finding indicated that grate genetic variability was existed among the 254 wheat genotypes which is the important key factor for future breeding program. Similar result for dendrogram analysis was reported by Pujar *et al*, 2020; Sansaloni *et al*, 2020.




Fig.4. Dendrogram based on the ward method for 254 wheat genotypes utilizing eleven quantitative traits

### Conclusion

Overall, the findings indicated that eleven quantitative traits explored significant and extensive amount of genetic diversity among 254 wheat varieties. The dendrogram analysis will be helpful for selecting diverse parents in breeding programme and in preserving genetic diversity of the present wheat germplasm.

### COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

### References:

- Ahmad, T. 2017. Evaluation of genetic variability in bread wheat (*Triticum aestivum* L. em Thell) genotypes. *Bull. Env. Pharmacol. Life Sci.*, 6 (3): 309-313.
- Atinafu, D. M., Alayachew, S. A. and Heterat, K. Z. 2020. Study of genetic variability in some bread wheat accessions (*Triticum aestivum* L.) in Gurage zone. Ethiopia. *Asian J. Biol. Sci.*, 13: 309-317.

- Baranwal, D.K., Mishra, V.K., Vishwakarma, M.K., Yadav, P.S. and Arun, B. 2012. Studies on genetic variability, correlation and path analysis for yield and yield contributing traits in wheat (*T. aestivum* L. em Thell.). *Plant Archives*, 12 (1): 99 - 104.
- Belay, F. and Fisseha, K. 2021. Genetic variability, heritability, genetic advance and divergence in Ethiopian cowpea [*Vigna unguiculata* (L) Walp] landraces. *J Agric Sc Food Technol.*, 7(1): 138-146. DOI: 10.17352/2455-815X.000101.
- Birhanu, M., Sentayehu, A., Alemayehu, A., Ermias, A. and Dargicho, D. 2016. Genetic variability, heritability and genetic advance for yield and yield related traits in bread wheat (*Triticum aestivum* L.) genotypes. *Glob. J. Sci. Front. Res.*, 16:12-15.
- De, N., Baranwal, D. and Kumar, S. 2014. Genetic evaluation of spring wheat (*Triticum aestivum* L.) genotypes for yield and spot blotch resistance in Eastern Gangetic Plains of India. *Afr. j. biotechnol.*, 13. 1867-1875. DOI:10.5897/AJB2014.13787.
- Devesh, P., Moitra, P.K., Shukla, R., Shukla, S., Pandey, S. and Arya, G. 2018. Analysis of Variability, Heritability and Genetic Advance of Yield, Its Components and Quality Traits in Wheat. *Int. J. Agri. Env. Biotechnol.*, 855-859.
- Dhananjay, B. N., Singhraj, S., Bhushan, B. and Rahul, V. P. 2012. Genetic Variability in wheat (*Triticum aestivum*) under Normal and Timely Sown Condition. *Env. Eco.*, 30: 1085-1087
- Falconer, D.S. and Mackay, T.F.C. 1996. Introduction to quantitative genetics. (4th Eds.), Longman, Essex, England.
- Fellahi, Z., Hannachi, A., Guendouz, A., Bouzerzour, H. and Boutekrabi, A. 2013. Genetic variability, heritability and association studies in bread wheat (*Triticum aestivum* L.) genotypes. *Electron. J. Plant Breed.*, 4(2): 1161-1166.
- Gerema, G. 2020. Evaluation of durum wheat (*Triticum turgidum*) genotypes for genetic variability, heritability, genetic advance and correlation studies. *J. Agri and Nat. Resour.*, 3. 150-159. 10.3126/janr.v3i2.32497.
- Goel, S., Yadav, M., Singh, K., Jaat, R. S. and Singh, N. K. 2018. Exploring diverse wheat germplasm for novel alleles in HMW-GS for bread quality improvement. *J. Food Sci. Technol.*, 55: 3257–3262.
- Govindaraj, M., Vetriventhan, M. and Srinivasan, M. 2015. Importance of genetic diversity assessment in crop plants and its recent advances: An overview of its analytical perspectives. *Genet. Res. Int.*, 2015:1-14. DOI: 10.1155/2015/431487.
- Hossain, M., Azad, M., Alam, M. and Eaton, T. 2021. Estimation of Variability, Heritability and Genetic Advance for Phenological, Physiological and Yield Contributing Attributes in Wheat Genotypes under Heat Stress Condition. *Am. J. Plant Sci.*, 12: 586-602. DOI: 10.4236/ajps.2021.124039.
- Johnson, H.W., Robinson, H. and Comstock, R.F. 1955. Estimates of genetic and environmental variability in soybean. *Agron. J.*, 47:314-318.
- Kilian, B., Martin, W. and Salamini, F. 2010. In *Evolution in Action: Case studies in Adaptive Radiation, Speciation and the Origin of Biodiversity* (ed Glaubrecht, M.). 137–166.
- Kolakar, S. S., Hanchinal, R.R. and Nadukeri, S. 2012. Assessment of genetic variability in wheat genotypes. *Adv. Res. J. Crop Improv.*, 3 (2): 114-117.
- Kumar, A., Singh, D., Kerkhi, S., Chand, P., Sirohi, A., Shami, V., Kumar, N. and Kumar, A. 2016. Genetic variability, heritability and genetic advance for yield and its contributing traits in bread wheat (*Triticum aestivum* L.). *Res. Environ. Life Sci.*, 9(3): 317-319.
- Larik, A.S, Kakar, A.A., Naz, M.A. and Shaikh, M.A. 1999. Estimating genetic parameters in bread wheat (*Triticum aestivum* L.) crosses. *Sarhad J. Agric.*, 15: 230-204.
- Lerner, I.M. 1958. *The Genetic Basis of Selection*, John Willey and Sons, New York. Pp. 57-65.

- Panase, V.G. 1957. Genetics of Quantitative characters in relation to plant breeding. *Indian J. Genet.*, 17: 318-328.
- Roy, A., Kumar, A., Singh, A., Mandi, A. and Barman, M. 2021. Analysis of genetic diversity and correlation studies on grain yield and its component characters in bread wheat (*Triticum aestivum* L. em Thell) genotypes. *Pharm. Innov. J.*, 10(5): 341-345.
- Saari, E. E. and Prescott, J. M. (1975). A scale for appraising the foliar intensity of wheat disease. *Plant Disease Reporter*, 59, 377-380.
- Shehrawat, S. and Kumar, Y. 2021. Genetic Architecture of Morpho-Physiological Traits in Wheat Accessions under Terminal Heat Stress. *Ekin. J.*, 7(1):34-42.
- Singh, B. and Upadhyay, P.K. 2013. Genetic variability, correlation and path analysis in wheat (*Triticum aestivum* L.). *Internat. J. Plant Sci.*, 8 (2): 230-235.
- Singh, M. K., Sharma, P. K., Tyagi, B. S. and Singh, G. 2013. Genetic analysis for morphological traits and protein content in bread wheat (*Triticum aestivum* L.) under normal and heat stress environments. *Indian J. Genet.*, 73: 320- 324.
- Singh, R.K. and Chaudhary, B.D. 1979. Biometrical methods in quantitative genetic analysis. Kalyani Publication, India.
- Tadesse, W., Sanchez-Garcia, M., Assefa, S. G., Amri, A., Bishaw, Z., Ogonnaya, F. C. and Baum, M. 2019. Genetic gains in wheat breeding and its role in feeding the world. *Crop Breed. Genet. Genom.*, 1:e190005.
- Teich, A. H. 1984. Heritability of grain yield, plant height and test weight of a population of winter wheat adapted to Southwestern Ontario. *Theor. Appl. Genet.*, 68(1):21-3. DIO: 10.1007/BF00252304.
- Ullah, K & Khan, S. & Khan, Muhammad & Rahman, Habib. 2012. Genetic variability, correlation and diversity studies in bread wheat (*Triticum aestivum* L.) germplasm. *Journal of Animal and Plant Sciences*. 22. 330-333.
- Velu, G., Singh, R. P., Huerta, J. and Guzmán, C. 2017. Genetic impact of Rht dwarfing genes on grain micronutrients concentration in wheat. *F. Crop. Res.*, 214: 373–377.
- Weraduwege, S. M., Chen, J, Anozie Francisca, C., Morales, A., Weise, S. E. and Sharkey, T. D. (2015) The relationship between leaf area growth and biomass accumulation in *Arabidopsis thaliana*. *Frontiers in Plant Science*, 6:167. DOI=10.3389/fpls.2015.00167
- Mondal, T., Mitra, B. and Das, S. 2018. Precision nutrient management in wheat (*Triticum aestivum*) using Nutrient Expert®: Growth phenology, yield, nitrogen-use efficiency and profitability under eastern sub-Himalayan plains. *Indian Journal of Agronomy*, 63: 174-180.
- Rahman, M. M., Crain, J., Haghghattalab, A., Singh, R. P. and Poland J. 2021. Improving Wheat Yield Prediction Using Secondary Traits and High-Density Phenotyping Under Heat-Stressed Environments. *Frontiers in Plant Science*, 12:1977-1989. DOI=10.3389/fpls.2021.633651
- Pradhan, P., Fischer, G., van Velthuisen, H., Reusser, D. E. and Kropp, J. P. 2015. Closing Yield Gaps: How Sustainable Can We Be? *PLOS ONE*, 10(6): e0129487. DOI: 10.1371/journal.pone.0129487
- Pujar, M., Govindaraj, M., Kanatti, A., Shivade, H. and Gangaprasad, S. 2020. Genetic variation and diversity for grain iron, zinc, protein and agronomic traits in advanced breeding lines of pearl millet [*Pennisetum glaucum* (L.) R. Br.] for biofortification breeding. *Genetic Resources and Crop Evolution*. 67. 1-14. 10.1007/s10722-020-00956-x.
- Sansaloni, C., Franco, J., Santos, B. *et al.* Diversity analysis of 80,000 wheat accessions reveals consequences and opportunities of selection footprints. *Nat Commun* 11, 4572 (2020). <https://doi.org/10.1038/s41467-020-18404-w>