

Original Research Article

Assessment of genetic variability and diversity in rice cultivars of North Bank Plain Zone of Assam for traits associated with grain yield and adaptation under moisture stress

Abstract

A set of 54 rice cultivars from North Bank Plain Zone (NBPZ) of Assam were assessed for their genetic variability and diversity with respect to yield and traits associated with moisture stress tolerance. A higher magnitude of phenotypic and genotypic coefficient of variation was recorded for root volume, root dry weight, root length density, filled grains per panicle, shoot dry weight, root length, root-shoot ratio and grain yield per plant. Out of all the traits studied, high heritability coupled with high genetic advance was registered for root dry weight, root length density, root volume, root-shoot ratio, root length and grain yield per plant. D^2 analysis revealed that the genotypes could be distributed into a group of eight clusters with maximum distance between cluster III and cluster V (845.71). Root volume was found to contribute highest towards divergence (12.83%). Out of all the clusters, cluster IV with an intra cluster distance of 370.76 consisted of maximum number of accessions (17). Genotypes based on the inter se genetic distance and per se performances of specific traits were identified for undertaking hybridization programme. The implication of the diversity pattern elucidated for breeding varieties for moisture stress adaptation were discussed.

Keywords: Genetic variability, Genetic diversity, Rice cultivars, North Bank Plain Zone of Assam, moisture stress

Introduction

Rice (*Oryza sativa* L.) is a noble crop feeding almost 50 per cent of the world's human population. It is extremely dynamic with its potential to grow under extreme climatic conditions ranging from water sufficient to water deficient conditions. The crop is grown from about five meters water depth (below mean sea level) to even the hilltops at about 3050 meters above sea level having no stagnation of water (Paudel, 2013). Its adaptability ranges from the tropical climate of Sri Lanka and Africa to the temperate climate of South Korea and Japan. The crop is cultivated in diverse ecosystems like upland, rainfed lowland, flood-prone, and irrigated. Rice crop consumes a relatively huge quantity of water during its growth period compared to other crops (Pandey and Shukla, 2015). About

5000 litres of water is required to produce 1 kg of rice grain (Sarma *et al.*, 2015).

Under the current global water crisis, water is thus, considered as a major limiting factor in rice production. Reduced water supply to the crop or a condition of water stress not only affects the grain yield but also adversely affects the grain quality. With the advent of climate change, even in the high rainfall zone of Assam, intermittent water crisis has been affecting rice yield in recent years. Root characteristics are, therefore, appears to be of paramount consideration under such water deficit situation. This investigation was, therefore, carried out on the currently cultivated varieties of the North Bank Plain Zone of Assam with a view to assess their diversity with respect to some root traits which are related with moisture stress adaptation.

Materials and methods

The materials for the present investigation consisted of a set of 54 rice cultivars (**Table 1**) comprising of 51 winter rice cultivars from farmers' fields of North Bank Plain Zone of Assam representing all the administrative blocks of the zone along with three hill rice genotypes. The investigation was carried out at the experimental field of Biswanath College of Agriculture, Biswanath Chariali, represented by the latitude of 26⁰15'N, 77⁰45'N, the longitude of 92⁰42'E, 95⁰30'E and altitude of 104 m MSL. The crop for the present investigation was raised during the Kharif season of 2019. The experiment was carried out using PVC (Poly Vinyl Chloride) tubes of 1 m length and 20 cm in diameter. The experiment was conducted following Completely Randomized Design with two replications. To ensure the homogeneity of the experimental units, same soil media was used throughout the experimental units (PVC tubes). Five to six seeds were directly sown into each PVC pipes. Upon germination and seedling establishment, three well-spaced seedlings were retained and others were discarded. The experiment was carried out under rainfed condition to impose a natural moisture stress. To quantify and understand the genetic variability, observations were recorded for 15 quantitative traits viz., days to 50 *per cent* flowering, days to maturity, plant height (cm), effective tillers per plant, filled grains per panicle, root length (cm), root volume (cm³), root dry weight (g), shoot dry weight (g), root-shoot ratio, total chlorophyll content, chlorophyll stability index, root length density, proline content and grain yield per plant. The data obtained were subjected to analysis of variance following standard protocol given by Panse and Sukhatme (1967). The mean sum of squares obtained from the analysis of variance were subjected to estimation of genetic parameters of variation as per Singh & Choudhury (1988). Estimates of variability

parameters, heritability and genetic advance were calculated using standard methods of Burton and Devane (1953), Lush (1945) and Johnson *et al.*, (1955). In order to understand the nature and magnitude of diversity prevailing in the rice genotypes, the 15 quantitative characters involving five root related characters were assessed for diversity using the concept of Mahalanobis' generalized distance (D^2) as suggested by Rao (1952).

Results and discussion

The analysis of variance revealed that the mean sum of squares due to genotypes was significant for all the 15 traits. This indicated the presence of significant variation among all the genotypes and there is an ample scope for selection of promising lines from the present gene pool for yield and other drought adaptive traits. The magnitude of variability indicated that the cultivars in the region are diverse in nature.

Many workers in the past reported the presence of significant variability in rice germplasm for these traits. The present findings find co linearity with the findings of Zheng *et al.* (1996) and Hemamalini (1997), who also observed significant variability for grain yield and root related components in rice. No study was however conducted in the rice germplasm of the region under consideration of present study. Thus, further dissection of the nature and magnitude of this variation in genetic terms would be meaningful. The estimates of various genetic parameters of variation are presented in **Table 2**. Highest genotypic and phenotypic coefficient of variance (GCV and PCV) was observed for root volume followed by root dry weight, root length density and grain yield per plant. Similar results were recorded by Ganapathy *et al.* (2010) with regards to traits viz., root length, root volume, root length density, root dry weight and root-shoot ratio evaluated under conditions of PVC pipes.

Moderate estimates of GCV and PCV were observed for plant height, effective tillers per plant, proline content, days to 50 *per cent* flowering and days to maturity. Nithya *et al.* (2020) reported similar results for plant height, tiller number, days to 50 *per cent* flowering and relative water content. Total chlorophyll content and chlorophyll stability index on the other hand showed low estimates of GCV and PCV. Khriedino *et al.* (2011) also reported similar findings for chlorophyll stability index. Presence of high magnitude of variation is also not the sole criterion for effectiveness of selection (Burton, 1952). Therefore, estimation of heritability and genetic advance are also required to arrive at a more reliable conclusion (Sarma and Richharia, 1995).

Based on this consideration, high heritability coupled with high genetic advance as *per cent* mean was observed for traits viz., root dry weight, root length density, root volume, root-shoot ratio, root length and grain yield per plant. Hemamalini (1997) reported similar results with respect to high heritability and genetic advancement as *per cent* mean for root number, root volume and root dry weight. This suggested preponderance of additive gene action in the expression of these characters.

High heritability associated with moderate genetic advancement as *per cent* mean was observed for days to 50 *per cent* flowering and days to maturity which are in accordance with the findings of Islam *et al.* (2015) and Sabesan *et al.* (2009). This indicated possibility of obtaining a reasonable response to selection in these traits owing to their high transmissibility even if the coefficient of variability was moderate (Kumar *et al.*, 2001 and Osman *et al.*, 2012).

In order to understand the nature and magnitude of genetic diversity in 54 genotypes of rice, the data obtained for the 15 quantitative characters were subjected to the analysis of Mahalanobis' generalized distance

(D^2). Based on D^2 values, 54 genotypes were grouped into five clusters (**Figure 1**). Khatun *et al.* (2015) and Worede *et al.* (2014) calculated a similar clustering pattern for 43 and 24 upland rice genotypes, respectively. Out of the 54 genotypes, three genotypes were present in cluster I, eight genotypes in cluster II, 16 genotypes in cluster III, 17 genotypes in cluster IV and 10 genotypes in cluster V. The inter - cluster D^2 values also ranged widely with a minimum value of 192.42 between cluster I and cluster II and a maximum value of 845.71 between cluster III and cluster IV (**Table 3**). The intra - cluster distance varied from 62.83 in cluster I to a maximum of 629.93 in cluster V. This revealed the presence of considerable diversity existed even within a cluster. Cluster III with 16 genotypes and cluster IV with 17 genotypes were the most divergent groups with a maximum inter-cluster distance (845.71). Therefore, intercrossing between these genotypes may be expected to yield better recombinants. Sandhyakishore *et al.* (2007) and Chandra *et al.* (2007) gave similar suggestions regarding intercrossing between the genotypes of clusters separated by large inter-cluster distances to achieve higher heterotic recombinants.

Among the 15 quantitative characters studied, the most important character contributing to the divergence was root volume followed by grain yield per plant, root dry weight and root length density (**Table 4**).

The presence of the wide genetic diversity with respect to grain yield and the root traits related to adaptation for moisture stress in the cultivars of the NBPZ of Assam appears to be significance for breeding rice varieties for resilience to climate change leading to periodic moisture stress experienced recently. The spectrum and magnitude of genetic variability obtained in segregating generation is the function of genetic distance between the parents. Thus, the set of cultivars

showed potential for selection of desirable parents in the above direction. However, while selecting parents for hybridization, based on their inter se genetic diversity, the per se performances of the genotypes should not be overlooked. While planning a hybridization programme, therefore, we should emphasize in one hand, the inter se genetic distance and on the other hand the per se performance of the genotypes for specific traits (Sarma *et al.*, 1995).

In the above considerations genotypes were identified for inclusion in hybridization programme. Out of 54 rice genotypes evaluated, the genotype Haccha belonging to cluster V was the earliest to attain 50 *per cent* flowering. It also exhibited the highest root-shoot ratio. Saru Jahinga belonging to cluster IV had the highest number of effective tillers per plant along with the highest root dry weight and root length density. This cultivar being a traditional photoperiod sensitive variety adapted to the native environment appears to be a promising parent contributing towards moisture stress tolerance. Maximum root length and root volume were recorded for the genotypes Vasudev and Jaibangla respectively belonging to cluster IV. Filled grains per panicle was found to be highest in Bhogali followed by Bahadur. These two genotypes too belong to cluster IV. Ronga Joha of cluster II had the highest grain yield per plant followed by Bhogali.

Based on the mean performance of the genotypes, the earliness of Haccha appears to be advantageous and can be utilized in crossing programme with Saru Jahinga (effective tillers, root dry weight and root length density), Vasudev and Jaibangla (root length and root volume), Bhogali (filled grains per panicle) and Ronga Joha (grain yield) to produce recombinants with considerable higher yield and moisture stress tolerance. Better performances with respect to root length and filled grains per panicle was

recorded for Bairing and Guni Dhan belonging to cluster III which are having the highest inter cluster distance with cluster V ($D^2 = 845.71$). Hence, hybridization between these genotypes with Haccha (cluster V) would yield short duration heterotic segregants. In this case, higher root length of Bairing would add to better tolerance to moisture stress through greater moisture absorption during water deficit condition. Also, GuniDhan (cluster III) with higher number of filled grains per panicle could be used as parent in hybridization programme with genotypes having higher intensities of traits related to moisture stress tolerance. Likewise, utilization of Ronga Joha as one of the parents for crossing with the genotypes with better root traits might yield desirable high yielding segregants under moisture stress environment. Thus, an efficient hybridization programme may be formulated out of the above consideration which would pave the way for obtaining basic genetic materials to undertake further selection and crop improvement programme.

Conclusion

The present diversity analysis indicated the presence of significant variability and diversity in the currently cultivated rice varieties of a traditional rice growing pocket of Assam. In view of the advent of climate change and recently experienced drought spells, the elucidation of diversity and identification of genotypes as donors for different moisture stress tolerance traits appears to be worthwhile. The findings of the study had provided the benchmark, based on which efficient crop improvement programme could be undertaken for breeding rice varieties adapted under moisture stress situations resulted due to climate change.

Ethical Issues: There is no any involvement of ethical issues regarding the report on the manuscript.

References

- Burton GW. Quantitative inheritance in grasses. Proc. 6th Int. Grassland Cong. 1952; **1**:211-283.
- Burton GW and Devane EH. Estimating heritability in tall fescue (*Festuca arundinaceae*) from replicated clonal material. Agron. J. 1953; **45**:478-481.
- Chandra K., Gopi K.C, Rao DV and Valarmathi K. Current status of freshwater faunal diversity in India, Zoological survey of India 2007..
- Das S, Subudhi HN and Reddy JN. Genetic variability in Grain quality characteristics and yield in lowland rice genotypes. *Oryza* 2001; **44**(4): 343-346.
- Ganapathy S, Ganesh SK, Shanmugasundaram P and Chandrababu R. Studies on root traits for drought tolerance in rice (*Oryza sativa* L.) under controlled (PVC pipes) condition. *Electr. J. Plant Breed.* 2010;**1**(4): 1016-1020.
- Hemamalini GS. Molecular mapping of low-moisture stress induced response in rice (*Oryza sativa* L.) roots at peak vegetative stage. M.Sc. (Agri.) Thesis, University of Agricultural Sciences, Bangalore 1997
- Islam MA, Raffi SA, Hossain MA.and Hasan AK. Analysis of genetic variability, heritability and genetic advance for yield and yield associated traits in some promising advanced lines of rice. *Progress. Agric.* 2015; **26**: 26-31.
- Johnson HW, Robinson HF and Comstock R E. Estimates of genetic and environmental variability in Soybean. *Agron J.* 1955; **47**:1314-1318.
- Khatun T, Hanafi M, Rafii MY, Wong MY, Salleh FM and Ferdous J. Genetic Variation, Heritability, and Diversity Analysis of Upland Rice (*Oryza sativa* L.) Genotypes Based on Quantitative Traits. *Biomed Res Int.* 2015;**29**:0861.
- Khriedinuo P, Fukrei, Kumar A, Tyagi W, Rai M and Pattanayak A. Genetic Variability in Yield and its Components in Upland Rice Grown in Acid Soils of North East India. *J. Rice Res.* 2011; **4**(1&2).
- Kumar B, Thakur R, Mishra SB and Singh D N. Variability studies in segregating population of rice (*Oryza sativa* L.). *Ann. Biol.* 2001; **17**(1): 43-48.
- Lush JL. Intra- sire correlation on regression of off-spring on dams as a method of estimating heritability of characters. *Proc. Am. Soc. Animal Prod.* 1945; **33**: 292-301
- Nithya N, Beena R, Roy S, Abida PS, Jayalekshmi VG, Viji MM and Manju RV. Genetic Variability, Heritability, Correlation Coefficient and Path Analysis of Morpho-physiological and Yield Related Traits of Rice under Drought Stress. *Chemical Science Review and Letters. Chem. Sci. Rev. Lett.* 2020; **9**(33): 48-54.
- Osman KA, Mustafa AM, Ali F, Yonglain Z and Fazhan Q. Genetic variability for yield and related attributes of upland rice genotypes in semi-arid zone (Sudan). *Afri. J. Agric.* 2012; **7**(33): 4613-4619.
- Pandey V and Shukla A. Acclimation and Tolerance Strategies of Rice (*Oryza sativa* L.) under drought stress. *Rice Sci.* 2015; **22**(4): 147–161.
- Pandey VR, Singh PK, Verma OP and Pandey P. Interrelationship and path coefficient estimation in rice under

- salt stress environment. *Int. J. Agric. Res.* 2012; **7**(4): 169-184.
- Panse VG. and Sukhatme PV. *Statistical methods for agricultural workers.* ICAR Publications, New Delhi 1967.
- Paudel M. Rice (*Oryza sativa* L.) cultivation in the highest elevation of the world. *Agron. J. Nepal* 2013; **2**: 31-41.
- Price AH, Tomos AD and Virk DS. Genetic dissection of root growth in rice (*Oryza sativa* L.), I: a hydroponic screen. *Theor. Appl. Genet.* 1997; **95**: 132-142.
- Rao CR. *Advanced Statistical Methods in Biometrical Research.* John Wiley and Sons Inc., New York 1952; pp: 236-272.
- Sabesan T, Suresh R and Saravanan K. Genetic variability and correlation for yield and grain quality characters of rice grown in coastal saline low land of Tamilnadu. *Electron. J. Plant Breed.* 2009;**1**: 56-59.
- Sandhyakishore N, Babu VR, Ansari NA and Chandran R. Genetic divergence analysis using yield and quality traits in rice (*Oryza sativa* L.). *Crop Improv.* 2007;**34**: 12-15.
- Sarma MK and Richharia AK. Genetic variability and diversity in rice under irrigated transplanted condition. *JASS* 1995; **8**(2): 154-157.
- Sarma MK, Agrawal RK and Richaria AK. Genetic diversity in rice under rainfed and irrigated ecosystems. *Oryza* 1995; **34**(1): 19-24.
- Singh RK and Chaudhary BD. *Biometrical Methods in Quantitative Genetic Analysis.* Kalyani Publishers, New Delhi, 1988.
- Worede F, Sreewongchai T, Phumichai C and Sripichitt P. Multivariate analysis of genetic diversity among some rice genotypes using morpho-agronomic traits. *J. Plant Sci.* 2014; **9**(1): 14–24.
- Zheng J., Yuan Z., Yin G., He G., Yang Z., Jiang K., Zheng J. K., Yuan Z. L., Yin GD, He GH, Yang ZL and Jiang KF. Genetic studies on root growth ability in seedling stage of rice. *Chinese J. Rice Sci.* 1996; **10**(1): 51-53.

Table 1: List of 54 indigenous rice cultivars taken for the investigation

Sl. No.	Genotype	Place of Collection	Sl. No.	Genotype	Place of Collection
1	BaisMuthi	Udalguri	28	BiriabhongaB	Lakhimpur
2	ChaparAijung	Darrang	29	SokMaichu	KarbiAnglong
3	Kola Joha	Darrang	30	KomolDhan	Sonitpur
4	Ranjit 1	Darrang	31	Bora	Sonitpur
5	Solpona	Dhemaji	32	Ranjit 2	Udalguri
6	SulsuliBao	Dhemaji	33	Mala	Biswanath
7	Boga Joha	Dhemaji	34	Prasadbhog	Biswanath
8	BasantiSali	Dhemaji	35	Bishnuprasad	Biswanath
9	BiyoiSali	Dhemaji	36	TTB-404	Biswanath
10	KosuSali	Dhemaji	37	Kunkuni Joha	Biswanath
11	Kon Joha	Dhemaji	38	Bhogali	Biswanath
12	MaguriDhan	Dhemaji	39	Luit	Biswanath
13	KokuaBao	Dhemaji	40	Agni Sali	Biswanath
14	GuniDhan	Dhemaji	41	Ranikajol Joha	Biswanath
15	Jaldubi	Dhemaji	42	Moinagiri	Biswanath
16	Suwagmoni	Dhemaji	43	Bahadur	Biswanath
17	Ronga Joha	Dhemaji	44	Dehangi	Biswanath
18	MalbhoogDhan	Dhemaji	45	Kanaklata	Biswanath
19	Aampakhi	Dhemaji	46	Jaibangla	Biswanath
20	CheniSakowa	Lakhimpur	47	Panindra	Biswanath
21	Betguti	Dhemaji	48	Boga Aijung	Biswanath
22	Gumraj	Dhemaji	49	Haccha	Haflong
23	Saru Jahinga	Dhemaji	50	Vasudev	Biswanath
24	Bor Jahinga	Dhemaji	51	Maibee	KarbiAnglong
25	Nepali Sokowa	Dhemaji	52	Bairing	KarbiAnglong
26	RangaSali	Dhemaji	53	Vandana	KarbiAnglong

UNDER PEER REVIEW

Table 2: Genetic parameters of variation for 15 quantitative traits in rice under moisture stress situation

Character	Mean	Range	GV	PV	EV	GCV (%)	PCV (%)	H _{bs}	GA (%)
Days to 50per cent flowering	114.537	69 - 144	223.82	225.69	1.87	13.06	13.12	99.17	26.80
Days to maturity	146.787	102 - 174	222.91	226.48	3.56	10.17	10.25	98.43	20.79
Plant Height	114.662	64.28 - 161.94	508.95	514.96	6.01	19.68	19.79	98.83	40.29
Effective tillers per plant	9.503	4 - 16	3.42	4.96	1.54	19.47	23.44	69.03	33.33
Filled grains per panicle	84.111	22 - 182	1018.07	1134.79	116.72	37.93	40.05	89.71	74.02
Root length	37.402	15.44 - 63.42	109.66	119.55	9.88	28.00	29.23	91.73	55.24
Root volume	23.905	6.5 - 59.5	157.98	165.83	7.85	52.58	53.87	95.27	105.72
Root dry weight	14.562	4.18 - 30.36	42.95	44.04	1.09	45.01	45.57	97.53	91.56
Shoot dry weight	44.295	15.94 - 83.74	232.55	236.98	4.43	34.43	34.75	98.13	70.25
Root-Shoot ratio	0.325	0.19 - 0.52	0.01	0.01	0.00	26.98	27.82	94.07	53.91
Total Chlorophyll content	1.688	1.24 - 1.87	0.01	0.01	0.01	4.66	6.65	49.17	6.74
Chlorophyll Stability Index	76.269	66.92 - 83.62	7.46	11.02	3.56	3.58	4.35	67.72	6.07
Root length density	0.139	0.04 - 0.29	0.00	0.00	0.00	45.01	45.57	97.53	91.56
Proline content	52.611	32.00 - 97.95	85.78	91.78	6.00	17.60	18.21	93.46	35.06
Grain yield per plant	13.625	3.69 - 34.69	36.39	41.28	4.89	44.28	47.16	88.16	85.64

Where,

GV : Genotypic Variance

PV : Phenotypic Variance

EV : Environmental Variance

GCV : Genotypic Coefficient of Variance

PCV : Phenotypic Coefficient of Variance

H_{bs} : Heritability in Broad Sense

GA : Genetic Advance as *per cent* mean

Table 3: Clustering pattern along with average inter and intra cluster distances for yield and root related characters

Clusters	Genotypes	I	II	III	IV	V
I	Gumraj, Kanaklata, Kola Joha	62.83	192.42	400.82	468.41	632.68
II	Ronga Joha, Bor Jahinga, MalbhoogDhan, Solpona, Boga Joha, Bishnuprasad, Panindra, Aampakhi	192.42	138.34	490.68	362.59	467.15
III	BasantiSali, Jaldubi, MaguriDhan, KosuSali, Mala, SokMaichu, Kokuabao, SulsuliBao, Boga Aijung, Bairing, Maibee, Biriabhongabao, GuniDhan, Ranjit 1, Ranikajol Joha, Luit	400.82	490.68	223.66	818.93	845.71
IV	Ranjit 2, Kunkuni Joha, Kon Joha, Moinagiri, RongaSali, Biyoisali, Suwagmoni, Bora, Betguti, Til Bora, Saru Jahinga, Jaibangla, Prasadbhog, Bahadur, Komoldhan, Bhogali, Vasudev	468.41	362.59	818.93	370.76	807.09
V	Nepali Sokowa, Agni Sali, Vandana, TTB-404, CheniSakowa, BaisMuthi, Inglongkiri, ChaparAijung, Dehangi, Haccha	632.68	467.15	845.71	807.09	629.93

* Diagonal values indicate intra cluster distances

* Above and below diagonal values indicate inter cluster distances

Table 4. *Per cent* contribution of each character towards divergence in ricegenotypes evaluated under low-moisture stress condition using PVC Pipes

Sl. No.	Character	Contribution (%)
1	Days to 50 per cent Flowering	3.13
2	Days to Maturity	2.44
3	Plant Height	4.71
4	Effective Tillers per plant	5.58
5	Filled Grains per Plant	9.54
6	Root Length	6.96
7	Root Volume	12.83
8	Root Dry Weight	10.85
9	Shoot Dry Weight	8.28
10	Root-Shoot Ratio	6.63
11	Total chlorophyll content	1.58
12	Chlorophyll Stability Index	1.04
13	Root Length Density	10.85
14	Proline Content	4.34
15	Grain yield per Plant	11.23
Total contribution		100




Figure 1: Mahalanobis Euclidean Distance