

## Original Research Article

### **Assessing the single and combined toxicity of chlorantraniliprole with *Bacillus thuringiensis* against maize fall armyworm *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) under laboratory conditions**

#### **ABSTRACT**

Single and combined toxicity of chlorantraniliprole and *Bacillus thuringiensis* against the 2<sup>nd</sup> and 3<sup>rd</sup> larval instars of *Spodoptera frugiperda* (J. E. Smith) were examined. Single toxicity of chlorantraniliprole against 2<sup>nd</sup> and 3<sup>rd</sup> larval instars were 0.87 and 1.52 ppm (LC<sub>25</sub>); 4.08 and 6.50 ppm (LC<sub>50</sub>), respectively. With respect to *Bt*, single toxicity against 2<sup>nd</sup> and 3<sup>rd</sup> larval instars were 474.39 and 693.48 ppm (LC<sub>25</sub>); 1008.62 and 1228.62 ppm (LC<sub>50</sub>), respectively. Combination effect of chlorantraniliprole with *Bt* revealed that 2<sup>nd</sup> instar of FAW showed supplemental synergism at LC<sub>50</sub> of chlorantraniliprole + LC<sub>25</sub> of *Bt*. In the case of LC<sub>50</sub> of chlorantraniliprole + LC<sub>50</sub> of *Bt*, LC<sub>25</sub> of chlorantraniliprole + LC<sub>50</sub> of *Bt* and LC<sub>25</sub> of chlorantraniliprole + LC<sub>50</sub> of *Bt* combinations, they showed sub additive synergism. In 3<sup>rd</sup> instar larvae, the combined toxicity results were similar for all the combinations of chlorantraniliprole + *Bt* except LC<sub>25</sub> of chlorantraniliprole + LC<sub>50</sub> of *Bt* where it showed an antagonistic synergism. Activity of Carboxyl esterase (CarE), mixed function oxidase (MFO) and Glutathione-S-transferase (GST) were found to be lesser in chlorantraniliprole LC<sub>50</sub> + *Bt* LC<sub>25</sub> combination than single toxicity treatments. Therefore, combined use of chlorantraniliprole with *Bt* at LC<sub>50</sub> of chlorantraniliprole + LC<sub>25</sub> of *Bt* had supplemental synergism on fall armyworm under laboratory condition.

**Key words:** Chlorantraniliprole, *Bacillus thuringiensis*, *Spodoptera frugiperda*, combined toxicity, synergism, maize

#### **1. Introduction**

Maize is the third most important food crop next to rice and wheat in India. Fall armyworm (FAW), *Spodoptera frugiperda* (J. E. Smith) is a cosmopolitan insect native to the tropical and subtropical regions of Americas (Wiseman *et al.*, 1966). Larvae of *S. frugiperda* are known to feed on more than 353 plant species (Montezano *et al.*, 2018). Larvae feed on all growth stages of maize and cause significant yield losses. In India, its incidence was first

noticed at the farmer's field in Chikkaballapur, Karnataka on maize (Sharanabasappa *et al.*, 2018). They recorded more than 70% of crop damage. Further, it was reported in Tamil Nadu, Kerala, Karnataka, Maharashtra and Madhya Pradesh in India. In India other than maize, *S. frugiperda* damage was recorded in sugarcane (Srikanth *et al.*, 2018), pearl millet, finger millet, sorghum (Venkateswarlu *et al.*, 2018), paddy (Kalleshwaraswamy *et al.*, 2019), ginger (Shankar and Adachi 2019), barnyard millet (Roopika, 2020), para grass, guinea grass, fodder maize, green amaranth (Maruthadurai and Ramesh, 2020), sugar beet (Shanthi *et al.*, 2020) and sunflower (Bankar *et al.*, 2021). Faster spread in geographical distribution and wider host adaptability make FAW a promising pest in India. For the immediate control of FAW menace, farmers rely highly on synthetic pesticides than other management measures. Use of pesticide is inevitable in the current situation to control FAW. This may led to development of resistance in FAW. Therefore, it is necessary to use pesticides in a manner that will address both resistance and residue in due course of time. The present investigation was aimed at improving the toxicity of chlorantraniliprole through addition of *Bacillus thuringiensis* (*Bt*) in combination for the effective management of FAW.

## **2 MATERIAL AND METHODS**

### **2.1 Insect culture**

*S. frugiperda* culture was maintained at the FAW lab, Department of Agricultural Entomology, Tamil Nadu Agricultural University, Coimbatore. Adults were provided with potted maize seedlings for oviposition. Then the plants were removed from the oviposition cage and transferred to an insect-proof cage to let the eggs hatch and larvae develop. The continuous supply of FAW culture was maintained using semi-synthetic artificial diet (Ashok *et al.*, 2021). The 2<sup>nd</sup> and 3<sup>rd</sup> instar larvae were used for all bioassays.

### **2.2 Bioassay**

Bioassays were conducted using newly moulted second and third instar larvae of *S. frugiperda* obtained from laboratory cultures using leaf disc bioassay method with slight modification (Ahmad *et al.*, 2009). Newly moulted second and third instar larvae were pre-starved for 4 h before bioassay. Each treatments consisted six concentrations of selected insecticides and an untreated control. Leaves were treated with different concentrations of insecticides and shade dried for 1 h. Pre-starved larvae were individually placed into the six well culture plates poured with 1% agar, maintained at 28±1 °C and 12:12 h (light: dark). Mortality data were recorded at 24, 48 and 72 hours after treatment (HAT).

### **2.3 Classification of synergism (Burgess and Hussey, 1971)**

### **2.3.1 Independent synergism (= independent action with zero correlation)**

A system of two components acting independently and not interfering with each other. If  $P_M$  is the probability of death by microorganism taken alone, and  $P_I$  the corresponding value for the insecticide, the probability of death by the combined action is  $P_{I+M} = P_I + P_M(1-P_I)$  or if the corresponding values of mortality in % are used:  $M_{I+M} = M_I + M_M(1-M_I/100)$ .

### **2.3.2 Subadditive synergism**

A system of two components which together produce an effect greater than independent synergism but less than the algebraic sum of two single effects. A weak potentiating effect is necessary to produce such a result.

### **2.3.3 Supplement synergism**

A system of two components which together produce an effect greater than the algebraic sum of single effects ( $M_{I+M} > M_I + M_M$ ).

### **2.3.4 Potentiating synergism**

A system of a component "A" causing the effect  $M_A$  and a synergist ("S") which alone cause no effect ( $M_S=0$ ), but which in combination produce an effect which is significantly greater than  $M_A$ . This type of synergism may be found when non-lethal concentrations of an insecticide are combined with a microorganism.

## **2.4 Detoxifying enzymes assay**

It was carried out to study the induction of detoxifying enzymes such as carboxyl esterase (CE), mixed function oxidase (MFO) and Glutathione-S-transferase (GST) in *S. frugiperda* larvae after treating with different insecticide combinations. Untreated larvae were maintained as control. The experiment was replicated three times, each with ten larvae. The surviving larvae were used for enzyme analysis.

### **2.4.1 Enzyme homogenate preparation**

The larvae surviving after treatment as described earlier was used for enzyme homogenate preparation. larvae was weighed and homogenized in ice-cold 20 mM phosphate buffer (pH 8.0) containing 0.2 per cent triton X-100 using pre-chilled pestle and mortar. Five ml of phosphate buffer was used for extraction. The homogenate was centrifuged at 15,000 rpm for 10 minutes and the supernatant collected served as an enzyme source for the assay.

#### **2.4.2 Protein estimation (Bradford, 1976)**

To one ml of enzyme extract, 5 ml of Bradford reagent was added and allowed for colour development. The absorbance was read at 595 nm. Using the standard graph, quantity of protein in the enzyme extract was calculated.

#### **2.4.3 Carboxylesterase (CarE) assay (Khalifa *et al.*, 2017)**

Five ml of the working substrate solution ( $\alpha$  naphthyl acetate) was mixed with 1ml of enzyme homogenate. After 30 minutes of incubation at room temperature, 1ml of coupling reagent (1% fast blue B salt: 5% sodiumlauryl sulphate = 2: 5, v/v) was added. A red colour developed immediately, which changed to fairly stable blue colour, was measured at 600 nm. Specific activity (SA) of the enzyme was estimated using the formula, which was expressed as n moles of  $\alpha$ -naphthol released  $\text{minute}^{-1}\text{mg of protein}^{-1}$ .

#### **2.4.4 Mixed function oxidases (MFO) assay (Khalifa *et al.*, 2017)**

To 500  $\mu\text{l}$  of enzyme source, 500  $\mu\text{l}$  of tris buffer (pH 7.8) and 20  $\mu\text{l}$  of p-nitroanisole were added. To this 50  $\mu\text{l}$  of NADPH was added in dark at room temperature for 30 minutes. The reaction was stopped by adding 0.5 ml of sodium hydroxide. The reaction mixture was centrifuged at 10,000 rpm for 30 minutes. The absorbance of supernatant was determined at 400 nm. The specific activity (SA) of the enzyme was calculated using the formula and expressed as n moles of p-nitrophenol released  $\text{minute}^{-1}\text{mg of protein}^{-1}$ .

#### **2.4.5 Glutathione-S-transferase assay (Khalifa *et al.*, 2017)**

The total GST activity was determined using CDNB (1-chloro 2,4-dinitrobenzene) and glutathione reduced as substrates. Glutathione-s-transferases activities were measured spectrophotometrically. Potassium phosphate buffer (20 mM) of pH 8.0 was used in assaying GST. 0.1M Glutathione reduced and 0.1M CDNB substrate was prepared in pure ethanol. Activity of GST was analysed by addition of 0.1 ml of glutathione reduced, 0.1 ml CDNB and 0.1 ml gut homogenate solution to final volume of 3 ml. The enzymatic reaction was monitored for the optical absorbance increase at wavelength 340 nm at 37 °C for 10 min at 1 min interval in the spectrophotometer. Specific GST activity was calculated and articulated in  $\text{n moles min}^{-1}\text{mg protein}^{-1}$ .

### **2.5 Statistical analysis**

Probit analysis was done to calculate  $\text{LC}_{25}$ ,  $\text{LC}_{50}$ ,  $\text{LC}_{90}$  using SPSS. The log concentration probit (LCP) lines were drawn by plotting log concentrations on X-axis and probits on Y-axis. The response of test insect populations was studied at different concentrations of the test insecticides. The combined toxicity of chlorantraniliprole with *Bt* was studied by combining the different lethal concentrations ( $\text{LC}_{25}$  and  $\text{LC}_{50}$ ) at different

proportions (Finney 1971). Mortality was corrected by Abbott's formula (Abbott 1925) for each Probit regression analysis.

### 3 RESULTS AND DISCUSSION

From probit analysis (Table 1), it was found that  $LC_{25}$  of chlorantraniliprole for 2<sup>nd</sup> and 3<sup>rd</sup> instar of FAW were 0.87 ppm and 1.52 ppm, respectively.  $LC_{25}$  of *Bt* for 2<sup>nd</sup> and 3<sup>rd</sup> instar of FAW were 474.39 ppm and 693.48 ppm, respectively.  $LC_{50}$  of chlorantraniliprole for 2<sup>nd</sup> and 3<sup>rd</sup> instar of FAW was 4.08 ppm and 6.50 ppm respectively.  $LC_{50}$  of *Bt* for 2<sup>nd</sup> and 3<sup>rd</sup> instar of FAW were 1008.62 ppm and 1228.62 ppm, respectively. Results on combination effect of chlorantraniliprole with *Bt* revealed that 2<sup>nd</sup> instar of FAW showed supplemental synergism at  $LC_{50}$  of chlorantraniliprole +  $LC_{25}$  of *Bt* (Table 2) as the resulted mortality was higher than their single effect and their independent synergism. In case of  $LC_{50}$  of chlorantraniliprole +  $LC_{50}$  of *Bt*,  $LC_{25}$  of chlorantraniliprole +  $LC_{50}$  of *Bt* and  $LC_{25}$  of chlorantraniliprole +  $LC_{50}$  of *Bt* combinations, they gave sub additive synergism as the mortality was higher than independent synergism but lesser than their sum of single effect. With respect to of 3<sup>rd</sup> instar larvae, the combined toxicity results were similar for all the combinations of chlorantraniliprole + *Bt* except chlorantraniliprole  $LC_{25}$  + *Bt*  $LC_{50}$  which showed an antagonistic synergism.

Levels of detoxifying enzymes were assessed in second and third instar larva of *S. frugiperda* treated with different combinations of chlorantraniliprole with *Bacillus thuringiensis*. Carboxylesterase (CarE), mixed function oxidases (MFO) and Glutathione-S-transferase were the three detoxifying enzymes assessed in the present study. Comparing the single and insecticide combination treatments CarE activity was found to be lesser in CH  $LC_{50}$  + *Bt*  $LC_{25}$  (3.76 fold) followed by CH  $LC_{25}$  + *Bt*  $LC_{50}$  (4.09 fold) than the CH<sub>50</sub> (7.39 fold), CH<sub>25</sub> (6.42 fold) in relation to control in second instar larvae. Similar trend was observed in third instar larvae also. The MFO activity was also found be lesser in insecticide combination treatments CH  $LC_{50}$  + *Bt*  $LC_{25}$  (1.06 fold) followed by CH  $LC_{25}$  + *Bt*  $LC_{50}$  (1.22 fold) than single action treatments CH<sub>50</sub> (1.72 fold), CH<sub>25</sub> (1.62 fold) in the second instar larvae. As like preceding enzyme activities, GST activity was also low in insecticide combination treatment in CH  $LC_{50}$  + *Bt*  $LC_{25}$  (1.01 fold) followed by CH  $LC_{25}$  + *Bt*  $LC_{50}$  (1.13 fold) than the CH<sub>50</sub> (1.87 fold), CH<sub>25</sub> (1.78 fold) in relation to control in second instar larvae.

Combinations of insecticides play a meaningful role compared to single insecticide because they have multiple modes of action. Hence, chemical mixtures may be effective in

management of lepidopteran pests (Shabbir, 2021). Our results suggested synergistic, additive, and antagonistic effects among the combinations of insecticides and *Bt* used against *S. frugiperda*. According to Burges and Hussey (1971), supplemental synergism was considered as significant as the mortality caused by combined toxicity is higher than sum of two single effect and their independent synergism. Mixtures can be advantageous compared to individual constituents, because they may have different modes of action and may delay the development of resistance (Konecka *et al.*, 2020). *S. frugiperda* larvae showed subadditive or supplement synergism in combination treatments of Ch+Bt because the chemical insecticides act as stressors and make the larvae more susceptible to Bt. The stressed insects generally seem to be more susceptible to pathogens (Steinhaus, 1958). Koppenhöfer and Kaya (1996) accounted similar kind of interaction between entomopathogenic nematodes and Bt whereas Ansari *et al.* (2004) reported synergistic action between *Metarhizium anisopliae* (Metchnikoff) and entomopathogenic nematodes. The mixture of Bt + (Bt + Chl) (1:1:1; LC<sub>50</sub>:LC<sub>50</sub>) produced a synergistic activity compared to other combinations in the field-original highly resistant strain of *P. xylostella* (Shabbir, 2021).

It is speculated that detoxification enzymes, such as glutathione S-transferase, Mixed Function Oxidase and carboxyl-esterase, play an essential role in the metabolism of carbamates, pyrethroids, and novel insecticides in numerous insects (Oakeshott *et al.*, 2019; Feyereisen, 2021). Our enzyme assays showed that there was a high level of GST activity in larvae exposed to chlorantraniliprole alone compared to the CHBt combinations. This significant correlation between GST activity and chlorantraniliprole suggests that the enzymes contribute to the detoxification of chlorantraniliprole. Hu *et al.* (2014) and Nehare *et al.* (2010) reported increased GST activity in *P. xylostella* treated with exogenous combinations of pesticides, such as indoxacarb, acephate, and chlorantraniliprole. Furthermore, a positive correlation with GST activity was found in response to spinosad insecticide also (Wang *et al.*, 2009; Chen and Zhang, 2014; Yalçın *et al.* 2015). Similarly, carboxyl-esterase (CarE) activity was also more in chlorantraniliprole alone treated larva of *S. frugiperda* than the combinations. CarE has also been directly correlated with resistance to chlorantraniliprole and abamectin resistance in *P. xylostella* (Gong *et al.*, 2013). The synergistic combinations may represent a first step towards the utilization of Bt products and insecticides that will considerably delay resistance. Therefore, combination of chlorantraniliprole with *Bacillus thuringiensis* at LC<sub>50</sub> of chlorantraniliprole + LC<sub>25</sub> of *Bt*. ratio had supplemental synergism on fall armyworm under laboratory condition.

Table 1. Toxicity of tested insecticides against 2<sup>nd</sup> and 3<sup>rd</sup> instar of *S. frugiperda*

Instar	Insecticide	LC <sub>25</sub> ppm	LC <sub>50</sub> ppm	Slope	$\chi^2$
2 <sup>nd</sup> instar	Chlorantraniliprole	0.87 (0.373 – 2.007)	4.08 (2.582 – 6.432)	1.0017	0.1261
	<i>B. thuringiensis</i>	474.39 (349.57 – 643.76)	1008.62 (800.31 – 1271.14)	1.9799	5.0485
3 <sup>rd</sup> instar	Chlorantraniliprole	1.52 (0.79 – 2.91)	6.50 (4.28 – 9.89)	1.0748	0.8848
	<i>B. thuringiensis</i>	693.48 (549.57 – 743.76)	1228.62 (1137.31 – 1371.14)	1.9799	5.2185


Table 2. Interactive effects of chlorantraniliprole + *Bt* on 2<sup>nd</sup> instar of *S. frugiperda*

Treatments	2 <sup>nd</sup> instar			
	Mortality (%)	Sum of two single effect	Independent synergism	Synergism
CH LC <sub>25</sub>	33.33			
CH LC <sub>50</sub>	57.14			
BT LC <sub>25</sub>	19.05			
BT LC <sub>50</sub>	28.57			
CH LC <sub>50</sub> + BT LC <sub>50</sub>	76.19	85.71	69.39	Subadditive synergism
CH LC <sub>50</sub> + BT LC <sub>25</sub>	80.95	76.19	65.31	Supplement synergism
CH LC <sub>25</sub> + BT LC <sub>50</sub>	57.14	61.90	52.38	Subadditive synergism
CH LC <sub>25</sub> + BT LC <sub>25</sub>	47.62	52.38	46.03	Subadditive synergism
Control	6.66			

Table 3. Interactive effects of chlorantraniliprole + *Bt* on 3<sup>rd</sup> instar of *S. frugiperda*

Treatments	3 <sup>rd</sup> instar			
	Mortality (%)	Sum of two single effect	Independent synergism	Synergism
CH LC <sub>25</sub>	38.10			
CH LC <sub>50</sub>	61.90			
BT LC <sub>25</sub>	9.52			
BT LC <sub>50</sub>	19.05			
CH LC <sub>50</sub> + BT LC <sub>50</sub>	71.43	80.95	69.16	Subadditive synergism
CH LC <sub>50</sub> + BT LC <sub>25</sub>	76.19	71.43	65.53	Supplement synergism
CH LC <sub>25</sub> + BT LC <sub>50</sub>	52.38	57.14	49.89	Subadditive synergism
CH LC <sub>25</sub> + BT LC <sub>25</sub>	42.86	47.62	43.99	Antagonistic effect
Control	3.33			

Fig. 1. Levels of detoxifying enzymes in Chlorantraniliprole + *Bacillus thuringiensis* treated 2<sup>nd</sup> and 3<sup>rd</sup> instar *S. frugiperda*



## COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## REFERENCES

- Ashok K, Balasubramani V, Kennedy J S, Geethalakshmi V, Jeyakumar P, Sathiah N. Evaluating artificial diets for the fall armyworm, *Spodoptera frugiperda* (JE Smith) (Lepidoptera: Noctuidae) through nutritional indices and an age-stage, two-sex life table approach. *African Entomology*. 2021: 29(2), 620-634.
- Ahmad M, Saleem M A, Sayyed A H. Efficacy of insecticide mixtures against pyrethroid and organophosphate-resistant populations of *Spodoptera litura* (Lepidoptera: Noctuidae). *Pest Management Science: formerly Pesticide Science*. 2009: 65(3), 266-274.
- Ansari M A, Tirry L, Moens M. Interaction between *Metarhizium anisopliae* CLO 53 and entomopathogenic nematodes for the control of *Hoplia philanthus*. *Biological Control*. 2004: 31(2), 172-180.
- Bankar D R, Bhamare V K, Mahajan R S and Kulkarni S. R. Biology of *Spodoptera frugiperda* (J. E. Smith) on Sunflower, *Helianthus annuus* (Linn.). Paper presented in International conference on global perspectives in crop protection for food security December 8-10, 2021, TNAU, Coimbatore, Tamil Nadu, India. 2021: pp 52-53.
- Bradford M M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*. 1976: 72(1-2), 248-254.
- Burges H D, Hussey N W. *Microbial control of insects and mites*. New York, Academic press. 1971: pp 327-353.

- Chen X E, Zhang Y L. Identification and characterisation of multiple glutathione S-transferase genes from the diamondback moth, *Plutella xylostella*. *Pest management science*. 2015: 71(4), 592-600.
- Feyereisen R. Insect CYP genes and P450 enzymes. In *Insect molecular biology and biochemistry*, Academic Press. 2012: 236-316.
- Gong Y J, Wang Z H, Shi B C, Kang Z J, Zhu L, Jin G H, Wei S J, Liu T X. Correlation between pesticide resistance and enzyme activity in the diamondback moth, *Plutella xylostella*. *Journal of Insect Science*. 2013: 13(1).
- Habig W H, Pabst M J, Fleischner G, Gatmaitan Z, Arias I M, Jakoby W B. The identity of glutathione S-transferase B with ligandin, a major binding protein of liver. *Proceedings of the National Academy of Sciences*. 1974: 71(10), 3879-3882.
- Hu Z D, Feng X, Lin Q S, Chen H Y, Li Z Y, Yin F, Liang P, Gao X W. Biochemical mechanism of chlorantraniliprole resistance in the diamondback moth, *Plutella xylostella* Linnaeus. *Journal of Integrative Agriculture*. 2014: 13(11), 2452-2459.
- Kalleshwaraswamy C M, Asokan R, Mahadevaswamy H M M, Sharanabasappa C M. First record of invasive fall armyworm, *Spodoptera frugiperda* (JE Smith) (Lepidoptera: Noctuidae) on rice (*Oryza sativa*) from India. *Journal of Entomology and Zoology Studies*. 2019: 7, 332-337.
- Konecka E, Kaznowski A, Grzesiek W, Nowicki P, Czarniewska E, Baranek J. Synergistic interaction between carvacrol and *Bacillus thuringiensis* crystalline proteins against *Cydia pomonella* and *Spodoptera exigua*. *Biocontrol*. 2020: 65(4): 447-460.
- Maruthadurai R, Ramesh R. Occurrence, damage pattern and biology of fall armyworm, *Spodoptera frugiperda* (JE smith) (Lepidoptera: Noctuidae) on fodder crops and green amaranth in Goa, India. *Phytoparasitica*. 2020: 48(1), 15-23.
- Montezano, D G, Sosa-Gómez D R, Specht A, Roque-Specht V F, Sousa-Silva J C, Paula-Moraes S D, Peterson J A, Hunt TE. Host plants of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the Americas. *African entomology*. 2018: 26(2), 286-300.

- Nehare S, Moharil M P, Ghodki B S, Lande G K, Bisane K D, Thakare A S, Barkhade U P. Biochemical analysis and synergistic suppression of indoxacarb resistance in *Plutella xylostella* L. Journal of Asia-Pacific Entomology. 2010: 13(2), 91-95.
- Oakeshott J, Claudianos C, Campbell P M, Newcomb R D, Russell R. Biochemical genetics and genomics of insect esterases. Comprehensive molecular insect science. 2010: Volume, 5.
- Roopika M, Srinivasan G, Shanthi M, Moorthy AV, Prabhu R. First record on barnyard millet as a new host for fall armyworm, *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) using novel insecticides and their residues on maize foliage. Paper presented in International seminar on Transboundary Pest Management March 4–5 2020, TNAU, Coimbatore, Tamil Nadu, India. 2020: p 45
- Rwomushana I. *Spodoptera frugiperda* (fall armyworm). Invasive Species Compendium. 2019: (29810).
- Shankar G, Adachi Y. First report of the occurrence of fall armyworm, *Spodoptera frugiperda* (JE Smith) (Lepidoptera: Noctuidae) on Ginger (*Zingiber officinale*) in Haveri district, Karnataka, India. J. Entomol. Zool. Stud. 2019: 7(5), 78-80.
- Shanthi M, Srinivasan G, Nivetha T K, Gurusamy A. First report on sugarbeet, as a new host for fall armyworm, *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) in Madhurai, Tamil Nadu. Paper presented in International Seminar on Transboundary Pest Management March 4–5 2020, TNAU, Coimbatore, Tamil Nadu, India. 2020: p 43
- Sharanabasappa, Kalleshwaraswamy C M, Asokan R, Swamy H M, Maruthi M S, Pavithra H B, Hegbe K, Navi S, Prabhu S T, Goergen G E. First report of the fall armyworm, *Spodoptera frugiperda* (JE Smith) (Lepidoptera: Noctuidae), an alien invasive pest on maize in India, Pest Management in Horticultural Ecosystem. 2018: 24(1): 23-29.
- Srikanth J, Geetha N, Singaravelu B, Ramasubramanian T, Mahesh P, Saravanan L, Salin K P, Chitra N, Muthukumar, M. First report of occurrence of fall armyworm *Spodoptera frugiperda* in sugarcane from Tamil Nadu, India. Journal of Sugarcane Research. 2018: 8(2), 195-202.

- Venkateswarlu U, Johnson M, Narasimhulu R, Muralikrishna T. Occurrence of the fall armyworm, *Spodoptera frugiperda* (JE Smith)(Lepidoptera, Noctuidae), a new pest on bajra and sorghum in the fields of agricultural research station, Ananthapuramu, Andhra Pradesh, India. Journal of Entomology and Zoology Studies. 2018: 6(6), 811-813.
- Wang D, Qiu X, Ren X, Niu F, Wang K. Resistance selection and biochemical characterization of spinosad resistance in *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae). Pesticide Biochemistry and Physiology. 2009: 95(2), 90-94.
- Wiseman B R, Painter R H, Wassom C E. Detecting corn seedling differences in the greenhouse by visual classification of damage by the fall armyworm. Journal of Economic Entomology. 1996: 59(5), 1211-1214.
- Yalcin M, Mermer S, Kozaci L D, Turgut C. Insecticide resistance in two populations of *Tuta absoluta* (Meyrick, 1917) (Lepidoptera: Gelechiidae) from Turkey. Türkiye Entomoloji Dergisi. 2015: 39(2), 137-145.
- Finney D J. Probit analysis 3rd edition. Cambridge University, London, UK. 1971: 333.
- Abbott W S. A method of computing the effectiveness of an insecticide. J. econ. Entomol. 1925: 18(2), pp.265-267.