

## Original Research Article

### Studies on Comparative Susceptibility of Chickpea Cultivars to *Callosobruchus chinensis* L. and *Callosobruchus analis* F.

#### ABSTRACT

Chickpea is highly preferred and is the important pulse crop of India. Bruchid (*Callosobruchus chinensis* L) is one of the most important insect pests of pulses in storage. In the present study eight cultivars (BG-256, Pusa-372, BGD-72, Pusa-1088, Pusa-5023, Pusa-5028, Pusa-1103, BG-1108) of chickpea were selected and evaluated against *Callosobruchus chinensis* and *Callosobruchus analis*. The choice and no choice methods were adopted to evaluate oviposition, emergence, mean development period (MDP) and growth index (GI) of selected chickpea cultivars on two years produce. The results revealed that varietal difference, old and fresh variety, different conditions i.e. (choice and no choice) affected the bruchid suitability or unsuitability. However, one year old varieties were found more suitable than fresh variety in case of *Callosobruchus analis* but in case of *Callosobruchus chinensis* except few varieties maximum fresh varieties were found susceptible than one year old varieties. These varietal characteristics responsible for such a behavior may be used and incorporated in breeding programme to develop high yielding varieties which are also resistant to insect infestation during storage.

*Keywords: Chickpea; cultivars; growth index; mean development period.*

## 1. INTRODUCTION

Chickpea (*Cicer arietinum* L.) is an ancient crop that has been grown in Asia, the Middle East and parts of Africa for many years [1,2]. India is a major chickpea-producing country with production 9.9 million tons and 10.041 t/ha productivity from 9.5 million ha area. According to Lale [3], grain storage has often resulted in quantitative and qualitative losses due to physical, chemical, and most important biological factors such as pests which may be birds, rodents, fungi and insects of which storage insect pests are the most important. Bruchid (*Callosobruchus chinensis* L) is one of the most important insect pests of pulses in storage. The larvae of this species feed and develop exclusively on the seed of chickpeas, while the adults do not require food or water and spend their limited lifespan (1–2 weeks) in mating and laying eggs on seeds. It cause damage during storage resulting in losses in quantity and nutritional quality. Apart from their direct losses by consumption of kernels, they accumulate frass, exuviae, and insect cadavers which may result in grain that is unfit for human consumption and/or induced changes in the storage environment warm, moist ‘hot-spots’ that are suitable for the development of storage fungi that cause further losses [2].

Chickpea is consumed as food and because of residue problem of insecticides, chemical control is not acceptable. Insecticide residues may remain on the treated crops, making them unfit for human consumption. In order to reduce both over dependence on chemicals for control and seed loss due to the bruchid attack, the search for host plant resistance in leguminous crops has increasingly become the option of choice in recent years [4]. The development and use of tolerant/resistant chickpea cultivars offer a simple, cheap and attractive way for the reduction of bruchid damage since it requires little knowledge by farmers, free of extra cost to farmers and also enhances the effectiveness of other pest control tactics such as cultural and biological means [5]. Hence, many studies were conducted periodically to evaluate seeds of many leguminous accessions for resistance against different bruchid species [6]. The present study was conducted to evaluate eight cultivars of chickpea for resistance against *Callosobruchus chinensis* and *Callosobruchus analis*.

## 2. MATERIALS AND METHODS

Eight varieties of chickpea (BG-256, Pusa-372, BGD-72, Pusa-1088, Pusa-5023, Pusa-5028, Pusa-1103, BG-1108) were selected for present study and 200 g cultivars produce of each were procured for two consecutive years from the pulse laboratory, Division of Genetics, Indian Agricultural Research Institute, New Delhi and the experiment was conducted in Division of Entomology, Indian Agricultural Research Institute, New Delhi. Each cultivar was kept separately in a muslin cloth tightened with rubber band and kept in desiccators maintaining 60% relative humidity by potassium hydroxide. Insect cultures of *Callosobruchus chinensis* and *Callosobruchus analis* were maintained separately as a stock culture in a glass jar (15.5 cm × 10.5 cm) on chickpea. Glass jar covered with markin cloth tightened with rubber band and kept at  $27\pm 1.5^{\circ}\text{C}$  and 60% relative humidity according to Solomon [7].

Free choice test was carried out in olfactory chamber and four grains were kept in a single hole and each cultivar randomly replicated thrice in the chamber. Five pairs of 3-4 days old adult insect were transferred in a central hole and fixed with lid. Like these three chambers were used which comprised nine replications. Forced choice test was carried in small petri-dish (7.5 cm dia.). Around 36 grains of each cultivar were kept in which 5 pairs of adult insects (3-4 days old) were transferred. Insects were separated after 24 hrs. in both (free and forced) tests. Four grains of each cultivar in both conditions were kept in a homeopathic vial plugged with cotton. These vials were kept in BOD chamber at  $27\pm 1.5^{\circ}\text{C}$  and 60% RH. The number of eggs laid on the surface were counted and after a week number of hatched eggs were also recorded. Similar procedures were followed in case of *Callosobruchus analis*. The adult emergence was recorded up to complete emergence. From the recorded data, growth index value Howe [8] was calculated from per cent emergence of adult beetle and mean development period.

### **3. RESULTS AND DISCUSSION**

#### **3.1 Susceptibility of Chickpea Produces to *Callosobruchus chinensis* during First Year**

**Free choice:** - Out of eight cultivars most suitable cultivar recorded Pusa-372 with growth index value 0.1461. Least susceptibility recorded in Pusa 5023 with GI value 0.1096. However maximum egg laying 5.88 was found in Pusa1088 on four grains followed by 5.55 eggs on Pusa372. The maximum emergence 90.09% recorded in Pusa-372. Mean development period varied from 30.0 days to 32.0 days.

**Force choice:** - In this method maximum eggs 9.00 recorded on BG1103 while the maximum per cent emergence 59.02 was found in BG-1108 and this variety was found most suitable with GI value 0.1743. Pusa1088 was found least susceptible cultivar with GI value 0.1099. However, mean development period (MDP) ranged between 23.4 (BG-1108) to 29.6 (BGD-72) (Table 1a).

### **3.2 Susceptibility of Chickpea Produces to *Callosobruchus chinensis* during Second Year**

**Free choice:** - In this fresher's cultivar, the minimum egg (0.33) was observed in Pusa 5023 and that too were 100 per cent emerged. Therefore, this variety observed the most susceptible with GI value (0.1550). Least susceptible variety recorded as Pusa1103 with GI value 0.0947. Mean development period ranged from 29.6 (Pusa-1103) to 31.2 (BG-1108).

**Force choice:** - Maximum (10.4) egg laying was found in BG1108 and maximum per cent emergence (82.21) recorded in this cultivar that is why this cultivar showed most susceptible with GI value 0.1959. Minimum (0.55) egg laying was observed on 4 grain and recorded minimum 20.00 per cent emergence in BGD-72. However, Pusa-372 showed least susceptibility of growth index value 0.0922 followed by BG-256 (0.1022) and BGD-72 (0.1185) (Table 1b).

### **3.3 Susceptibility of Chickpea Produces to *Callosobruchus analis* during First Year**

**Free choice:** - Egg laying was not minimum but emergence was minimum 35.11% recorded in Pusa1088, mean development period was maximum 34.00 days, but this cultivar was found to be least susceptible with growth index 0.1047. This cultivar was followed by Pusa5028 (0.1213) and BG256 (0.1233). Most susceptible cultivar noted as BG1108 as the highest per cent emergence 88.5 and maximum growth index (GI) 0.1443.

**Force choice:** - More than 80 per cent emergence observed with the minimum development period of 28.2 days in BG1108. Therefore, in this condition also maximum growth index 0.1570 observed and cultivar found to be most susceptible for *Callosobruchus analis*. Least susceptible cultivar was differed with the free choice condition and recorded minimum (28.57) per cent emergence and minimum growth index value (0.1117) in case of BGD-72 followed by Pusa1088 (0.1326) and BG-256 (0.1366) (Table 2a).

### **3.4 Susceptibility of Chickpea Produces to *Callosobruchus analis* during Second Year**

**Free choice:** - Maximum (5.00) eggs laid on Pusa5023 and the maximum per cent emergence (81.35) was observed in BGD-72 that is why this cultivar showed maximum susceptibility to *Callosobruchus analis* with the maximum value (0.1293) of growth index. The minimum (0.0988) value of growth index and the minimum per cent emergence 22.68 was observed in Pusa5028 showed least susceptibility. This variety followed by Pusa1103 (0.1062), Pusa1088 (0.1221) and BG1108 (0.1224) (Table 2b).

**Force choice:** - The result of force choice was similar as like as number of eggs laid on grain surface. The maximum 14.22 eggs per 4 grain was observed in pusa5023, but the maximum (80.14) per cent emergence was recorded in Pusa 1088. Mean development period (MDP) varied from 30.3 (Pusa-1103) to 34.00 (Pusa-1088). In this condition the least susceptibility was observed in BGD-72 (0.1093) and the most susceptible cultivar was recorded Pusa5023 with growth index value (0.1549).

Khanna et al. [9] reported BGD-72 and BG-1103 the most susceptible for *Callosobruchus chinensis* and *Callosobruchus analis* in the testing of 7 cultivars in free choice condition. They also recorded BG-1101 cultivar was least susceptible with growth index (GI) value 0.91 and 1.13, respectively, for *Callosobruchus chinensis* and *Callosobruchus analis*. Jha et al. [10] reported that Pusa-72 was better with GI value 0.153 and 0.377 for free and force choice condition, respectively. However, *Callosobruchus analis* showed that cultivar Pusa-1088 was least susceptible and GI values found 0.0259 and 0.107 for force and free choice condition, respectively. Similarly, Jha et al. [11] in different combinations of eight cultivars of chickpea found BGD-72, BG -1103 and Pusa-372 showed GI value 0.082, 0.073 and 0.101, respectively, for *Callosobruchus analis* in free choice conditions. However, in case of force choice conditions no development recorded in two varieties under force choice conditions only BGD-72 showed 0.107 GI value. In case of *Callosobruchus chinensis* BGD-72, BG-1103 and Pusa-372 responded 0.159, 0.156 and 0.153 GI value, respectively, for free choice while 0.186, 0.187 and 0.018 recorded for force choice test.

#### 4. CONCLUSION

The bruchid suitability or unsuitability was influenced by varietal differences, old and fresh varieties, and varied situations (choice and no option). In the case of *Callosobruchus analis*, one

year old variations were found to be more suited than fresh variants, however in the case of *Callosobruchus chinensis*, fresh varieties were shown to be more susceptible than one year old varieties, with the exception of a few types.

## REFERENCES

1. Tesfu FT. Evaluation of *Parthenium hysterophorus* powder against *Callosobruchus chinensis* L (Coleoptera; Bruchidae) on chickpea under laboratory condition. A Thesis submitted to the School of Graduate Studies in partial fulfilment of the requirements for the Degree of Masters of Science in Zoological Sciences program Units (Insect Science Stream), Addis Ababa university, Ethiopia. 2011.
2. FAO. Year Book. Food and Agriculture Organization of the United Nations. 2005.
3. Lale N. Stored product Entomology and Acarology. Mole Publications, Maiduguri. 2002; pp. 264.
4. Shaheen FA, Khaliq A, Aslam M. Resistance of chickpea (*Cicer arietinum* L.) cultivars against pulse beetle. Pakistan Journal of Botany. 2006; 38(4): 1237-1244.
5. Thomas MB, Waage JK. Integration of biological control and host plant resistance breeding for control of insect pests. In CTA-IAF. IIBC Seminar. 1995; pp. 9-14.
6. Raghuwanshi PK, Sharma S, Bele M, Kumar D. Screening of certain gram genotypes against *Callosobruchus chinensis* L. (Coleoptera: Bruchidae). Legume Research. 2016; 39(4): 651-653.
7. Solomon ME. Control of humidity with potassium hydroxide, sulphuric acid and other solutions. Bulletin of Entomological Research. 1951; 42:543-554.
8. Howe RW. A parameter for expressing the suitability of an environment for insect development. Journal of Stored Product Research. 1971; 7:63-65.
9. Khanna SC, Jha AN, Yadav SS. Response of chick pea cultivars to *Callosobruchus chinensis* and *Callosobruchus analis* infestation during storage. In: Proceeding of National Symposium on Frontier Areas of Entomological Research held during 5-7 Nov., 2003; pp.206.
10. Jha AN, Srivastava C, Kumar J. Host susceptibility of chick pea (*Cicer areitinum*) cultivars to *Callosobruchus chinensis* and *Callosobruchus analis*. Indian Journal of Agricultural Sciences. 2009; 79(5):372-374.
11. Jha AN, Srivastava C, Kumar J. Host susceptibility of chick pea cultivars to *Callosobruchus chinensis* and *Callosobruchus analis*. Indian journal of Entomology. 2009; 71(1):72-75.

**Table 1a. Mean value of oviposition, emergence, per cent emergence, mean development period (MDP) and growth index (GI) of *Callosobruchus chinensis* in different cultivars of chickpea during first year under free and force choice condition.**

S. No.	Cultivars	Oviposition	Emergence	Per cent emergence	MDP	GI
1.	BG-256	2.10 (3.77)	1.33 (2.00)	63.33 (53.05)	31.00 (28.9)	0.1338 0.1374
2.	Pusa-372	5.55 (6.66)	5.00 (2.11)	90.09 (31.68)	30.8 (28.4)	0.1461 0.1217
3.	BGD-72	0.44 (1.11)	0.22 (0.44)	50.00 (39.64)	30.00 (29.6)	0.1305 0.1243
4.	Pusa-1088	5.88 (5.22)	4.66 (1.11)	79.25 (21.26)	30.84 (27.8)	0.1418 0.1099
5.	Pusa-5023	0.66 (0.66)	0.22 (0.11)	33.33 (16.66)	32.00 (25.00)	0.1096 0.1125
6.	Pusa-5028	0.22 (4.77)	1.11 (2.11)	50.00 (44.23)	30.00 (24.4)	0.1304 0.1553
7.	Pusa-1103	2.22 (9.00)	1.66 (4.11)	74.77 (45.66)	30.5 (24.3)	0.1414 0.1572
8.	BG-1108	2.11 (4.88)	1.44 (2.88)	68.24 (59.02)	30.13 (23.4)	0.1402 0.1743

Figures in parentheses are force choice condition

**Table 1b. Mean value of oviposition, emergence, per cent emergence, mean development period (MDP) and growth index (GI) of *Callosobruchus chinensis* in different cultivars of chickpea during second year under free and force choice condition.**

S. No.	Cultivars	Oviposition (No.)	Emergence	Per cent emergence	MDP	GI
1.	BG-256	1.11 (1.00)	0.77 (0.33)	69.36 (33.00)	30.6 (34.2)	0.1385 0.1022
2.	Pusa-372	1.66 (1.00)	1.11 (0.23)	66.86 (22.00)	30.6 (33.5)	0.1373 0.0922
3.	BGD-72	1.11 (0.55)	0.88 (0.11)	79.27 (20.00)	30.6 (33.0)	0.1429 0.1185
4.	Pusa-1088	6.11 (6.00)	5.33 (2.33)	87.23 (38.83)	30.6 (28.8)	0.1446 0.1270
5.	Pusa-5023	0.33 (3.22)	0.33 (2.44)	100.00 (75.77)	29.7 (22.2)	0.1550 0.1949
6.	Pusa-5028	2.00 (6.33)	1.66 (2.11)	83.00 (33.33)	30.6 (23.5)	0.1444 0.1492
7.	Pusa-1103	2.00 (8.44)	0.33 (4.88)	16.50 (57.82)	29.6 (22.8)	0.0947 0.1779
8.	BG-1108	5.00 (10.4)	3.66 (8.55)	72.20 (82.21)	31.2 (22.5)	0.1372 0.1959

Figures in parentheses are force choice condition.

**Table 2a. Mean value of oviposition, emergence, per cent emergence, mean development period (MDP) and growth index (GI) of *Callosobruchus analis* to chickpea cultivars during first year under free and force choice condition.**

S. No.	Cultivars	Oviposition (No.)	Emergence	Per cent emergence	MDP	GI
1.	BG-256	1.33 (3.77)	0.88 (1.88)	66.16 (49.86)	34.00 (28.6)	0.1233 0.1366
2.	Pusa-372	1.00 (0.77)	0.77 (0.22)	77.00 (85.71)	34.00 (28.4)	0.1277 0.1567
3.	BGD-72	0.11 (0.77)	0.11 (0.22)	100.00 (28.57)	34.00 (30.00)	0.1354 0.1117
4.	Pusa-1088	1.88 (0.44)	0.66 (0.22)	35.11 (50.00)	34.00 (29.5)	0.1047 0.1326
5.	Pusa-5023	1.55 (11.77)	1.00 (8.88)	64.52 (75.45)	32.1 (29.4)	0.1384 0.1470
6.	Pusa-5028	1.88 (6.11)	0.88 (3.77)	46.81 (61.70)	31.7 (28.4)	0.1213 0.1451
7.	Pusa-1103	3.66 (5.77)	1.77 (2.88)	48.36 (49.91)	30.9 (28.5)	0.1255 0.1372
8.	BG-1108	2.00 (12.33)	1.77 (10.33)	88.5 (83.78)	31.0 (28.2)	0.1443 0.1570

Figures in parentheses are force choice condition

**Table 2b. Mean value of oviposition, emergence, per cent emergence, mean development period (MDP) and growth index (GI) of *Callosobruchus analis* in different cultivars of chickpea during second year under free and force choice condition.**

S. No.	Cultivars	Oviposition (No.)	Emergence	Per cent emergence	MDP	GI
1.	BG-256	0.77	0.55	71.43	34.00	0.1255
		(3.66)	(2.55)	(69.67)	(32.1)	0.1322
2.	Pusa-372	1.55	1.11	71.61	34.00	0.1256
		(1.00)	(0.44)	(44.00)	(32.00)	0.1182
3.	BGD-72	1.77	1.44	81.35	34.00	0.1293
		(1.00)	(0.33)	(33.00)	(32.00)	0.1093
4.	Pusa-1088	2.44	1.55	63.52	34.00	0.1221
		(2.77)	(2.22)	(80.14)	(32.1)	0.1365
5.	Pusa-5023	5.00	2.55	51.00	30.7	0.1281
		(14.22)	(10.22)	(71.87)	(27.6)	0.1549
6.	Pusa-5028	3.88	0.88	22.68	31.6	0.0988
		(4.88)	(2.44)	(50.00)	(27.1)	0.1443
7.	Pusa-1103	4.44	1.11	25.00	30.3	0.1062
		(11.66)	(5.11)	(43.82)	(28.2)	0.1340
8.	BG-1108	3.88	1.77	45.62	31.2	0.1224
		(5.33)	(3.22)	(60.41)	(28.0)	0.1465

Figures in parentheses are force choice condition