

Effect of Micronutrients on physico-chemical parameters of fruit and yield of Guava (*Psidium guajava* L.) cv. L-49.

Abstract

A field experiment entitled “Effect of Micronutrients on physico-chemical parameters of fruit and yield of Guava (*Psidium guajava* L.) cv. L-49.” was carried out at Horticulture Research Farm, Department of Horticulture, Babasaheb Bhimrao Ambedkar University, Vidya-Vihar, Rae Bareilly Road, Lucknow 226 025 (U.P.) during the summer season of 2019-20. The experiment consisted of eight treatment combinations having Zn, Mn and Fe concentration of 0.5, 0.5 and 0.4 % respectively, in Randomized Block Design with three replications. There was significant effect of micronutrients (Zn, Mn and Fe) on physico-chemical parameters of fruit and yield of guava respectively, alone and combination over control. The main objectives of the present investigation were to be find out the effect of Zn, Fe, and Mn on flowering, fruiting, physico- chemical composition of Guava fruit and yield.

Key words: Guava, Micronutrients, Physico-Chemical Parameters, Yield, Zn, Mn and Fe

Introduction

Guava (*Psidium guajava* L.) “Apple of the tropics” is an important fruit crop of country, not because of large area and production but due to its wider edapho-climatic adaptability, hardly to various biotic and abiotic stresses, precocious and prolific bearing habit, quality fruit with high nutritive value, medicinal attribute, use both as fresh fruit and after processing in different value added products and considered as multipurpose tree due to its utility as a fruit, fuel, fodder, timber and it is highly remunerative crop. It is a very rich source of vitamin C and A along with minerals like iron, calcium, and phosphorus. Guava is one of the fourth most important fruit crop in India after Mango, Banana and Citrus (Ray 2002). It also contains substantial quantities of carbohydrates, sugars and pectin. Owing to excellent taste and flavour, high nutritional value and wide availability at moderate price the fruit is often called as “Poor man’s apple”. The conventional products of guava are jelly, jam, pulp, concentrate, juice, cheese, toffee, dehydrated guava and canned guava.

Guava is commercially grown in throughout the country particularly Maharashtra, Uttar Pradesh, Bihar, Orissa, Punjab, Uttarakhand, Gujarat, Madhya Pradesh, and west Bengal. Uttar Pradesh is considered as the most important guava producing state of India, in which Allahabad (Prayagraj) region has the reputation of growing the best quality guava in the country as well as all over the world.

It covers around 3.7% (2.7 lakh ha) of total area under fruit crops and contributes 3.3% (41.07 MT) of total fruit production (NHB 2017-18).

The spring flowering is called “Ambe Bahar”, June or monsoon flowering is called “Mrig Bahar” and third flowering which comes in October is called “Hast Bahar”. Ambe Bahar fruits ripen from July to September and Mrig Bahar fruits ripen from November to February however, Hast Bahar fruits ripen in spring season which is also known as summer season.

The fruit is an excellent source of vitamin C (210 mg/100g) and pectin (0.60 %) but has low energy (66 cal /100g), protein content (1%) and has dry matter (17%) and moisture (83%). The fruit is also rich in minerals like phosphorus (24- 37mg/100g), calcium (14-30 mg/100g) and iron (0.6-1.4mg/100g) as well as vitamins like niacin, pantothenic acid, thiamine, riboflavin and vitamin A (Bose *et al.*, 1990).

Foliar application is based on the principle that the nutrients are quickly absorbed by leaves and transported to different parts of the plant to fulfil the functional requirement of nutrition. This method is highly helpful for the correction of element deficiencies to restore disrupted nutrient supply, overcome stress factors limiting their availability and it plays an important role in improving fruit set, productivity and quality of fruits and recovery of nutritional and physiological disorders in fruit trees.

Zinc is the important constituent of several enzyme systems which regulate various metabolic reactions associated with water relations in the plant. Zinc is essential for auxin and protein synthesis, seed production and proper maturity. It also increases fruit size as well as yield. Zinc is essential for improving the vegetative growth of guava trees obtained in terms of terminal shoots, shoot diameter and number of leaves per shoot (Price *et al.*, 1972). Boron is a constituent of cell membrane and essential for cell division. It acts as a regulator of potassium / calcium ratio in the plant and helps in nitrogen absorption and translocation of sugar in plant. Boron increases nitrogen availability to plant. It is involved in the synthesis of cell wall components. It has a central role in pollen viability and good fruit set. It increases the elongation growth of primary and lateral roots (O' Kelley, 1957).

Iron plays a critical role in metabolic processes such as DNA synthesis, respiration, and photosynthesis. It serves as a component of many vital enzymes such as cytochromes of the electron transport chain, and it is thus required for a wide range of biological functions. In plants, iron is involved in the synthesis of chlorophyll, and it is essential for the maintenance of chloroplast structure and function (Rout and Sahoo, 2015).

Manganese plays an essential role in respiration and N metabolism in plants. Also acts as an activator for the enzymes, nitrite reductase and hydroxylamine reductase. Manganese is directly or

indirectly involved in chloroplast formation and probably in their multiplication and also involved in various oxidation-reduction reactions in photosynthesis (Nijjar, 1990).

Material and methods

The present investigation was carried out at Horticulture Research Farm, Department of Horticulture, Babasaheb Bhimrao Ambedkar University (A Central University), Vidya-Vihar, Rae Bareli Road, Lucknow 226 025 (U.P.), India during the 2019-20. The Horticulture Research Farm, Department of Applied Plant Science Horticulture, Babasaheb Bhimrao Ambedkar University, Vidya- Vihar, Rae Bareli Road, Lucknow is situated at elevation of 111 meter above mean sea level in the subtropical tracts of Central U.P. at 260-560 North latitude. The experiment is laid out in randomized block design with factorial concept containing 8 treatment combinations (T₀, T₁, T₂, T₃, T₄, T₅, T₆, T₇) in which T₀: Control (Water Spray). T₁: Zinc Sulphate 0.5%, T₂: Manganese Sulphate 0.5%, T₃: Iron Sulphate 0.4% T₄: Zinc Sulphate + Manganese Sulphate 0.5% + 0.5%, T₅: Zinc Sulphate + Iron Sulphate 0.5% + 0.4%, T₆: Manganese Sulphate + Iron Sulphate 0.5% + 0.4%, T₇: Zinc Sulphate + Manganese Sulphate + Iron Sulphate 0.5% + 0.5% + 0.4%. Application of treatment was given in two doses 1st spray at Fruit Set and 2nd Spray 15 days after Fruit Set.

Table:1. Treatment details

S. No.	Treatment notation	Treatment content
1.	T ₀	Control (Water Spray)
2.	T ₁	Zinc Sulphate (0.5 %)
3.	T ₂	Manganese Sulphate (0.5 %)
4.	T ₃	Iron Sulphate (0.4%)
5.	T ₄	Zinc Sulphate + Manganese Sulphate (0.5 % + 0.5 %)
6.	T ₅	Zinc Sulphate + Iron Sulphate (0.5 % + 0.4%)
7.	T ₆	Manganese Sulphate + Iron Sulphate (0.5 % + 0.4%)
8.	T ₇	Zinc Sulphate + Manganese Sulphate + Iron Sulphate (0.5 % + 0.5 % + 0.4%)

Plant parameters recorded during the course of experiment are Physico-chemical characteristic of fruits i.e., fruit size (cm), fruit weight (gm), volume of fruit (cm³), specific gravity and chemical parameters are Total soluble solids (TSS ⁰Brix), acidity (%), ascorbic acid (Vitamin 'C')

(mg/100gm pulp), reducing sugars (%), non-reducing sugar (%) and total sugar (%) are recorded. The data mended during the experimentation were statistically analysed by randomized block design as suggested by Panse *et al.* (1995).

Results and Discussion

fruit characteristics (physical) of guava

The result revealed that application of micronutrients on Physico-chemical parameters and yield of guava. The parameters *viz.* Fruit length, Fruit breadth, Fruit weight, Fruit volume, Fruit set, Fruit drop, Total soluble solids (TSS), total sugar, ascorbic acid content, number of fruits per plant, yield and yield efficiency.

Fruit length and breadth the data reveal that different micronutrients application exhibited a significant influence on fruit length and breadth during the course of study. Maximum fruit length (64.33 mm) was recorded in T₇. Whereas, the minimum fruit length (54.73 mm) was observed under treatment T₂. Maximum fruit breadth (68.03 mm) was recorded in T₇. Minimum fruit breadth (61.12 mm) was, however, observed in T₀ (control).

The data on fruit weight as influenced by different micronutrients application reveals that different micronutrients application had a significant effect on fruit weight which ranged from 116.93 g – 165.37 g. Maximum fruit weight (165.37 g) was recorded in T₇. The minimum fruit weight (116.93 g) was, however, observed in T₀ (control).

Different micronutrients effected fruit volume considerably whose value varied from 121.67 cc to 182.67 cc. Highest fruit volume (168.33 cc) was in T₇ and minimum (121.67 cc) in T₀ (control). As indicated above, the application of different micronutrients alone or in combination has significant effect on fruit size (length and breadth), weight and volume. The possible reason for this effect might be due to the fact that mineral nutrients appear to have indirect role in hastening the process of cell division and cell elongation due to which the size of fruit might have improved. The increase in fruit weight may also be due to the rapid increase in the size of cells or it is also due the fact that foliar application of boron increased the fruit weight eventually by maintaining lighter level of auxins in various parts of the fruits which helped in increasing the fruit growth (Kaur, 2017). The findings are similar to those reported by Bagali *et al.*, (1993), Pal *et al.*, (2008), Kumawat *et al.*, (2012) in guava and Meena *et al.*, (2014) in aonla who recorded maximum fruit size and weight with the foliar application of micronutrients.

The fruit set was not much influenced by micronutrients application during the course of study (Table 2). However, the maximum fruit set (61.32 %) was recorded in T₇ and minimum (60.03%) in T₁. As is evidenced from the data in Table 3, the micronutrients alone or in combinations had no significant effect on fruit set. The possible reason for non- significant effect on fruit set in the present study might be because of the foliar application of micronutrients started after the fruit set.

It is evident from the data presented in Table 2 those different treatments of micronutrient application exhibited marked influence on the extent of fruit drop during the course of study. Minimum fruit drop (42.08 %) was recorded in T₇. Maximum fruit drop (58.60 %) was recorded in T₀ (control).

The data in Table 2 indicate that the foliar application of different micronutrients alone or in combination resulted in significant effect on fruit drop. The minimum fruit drop was observed in trees sprayed with zinc sulphate @ 0.5% + manganese sulphate @ 0.5% + iron sulphate @ 0.4% (T₇). The results are in conformity with those of Hada *et al.* (2014); Bagali *et al.* (1993); Balakrishnan (2000) who found reduction in fruit drop in guava with foliar application of micronutrients. The reduction in fruit drop might be due to the fact that the foliar application of micronutrients affected metabolic activities of the tree, improved the source sink relationship and favourably influenced the metabolic status resulting in better control of drop and enhancing the fruits retention (Katiyar *et al.*, 2008). Wright (1956) also suggested that the primitive effect of growth substances attributed in greater retention of fruit and reduction in fruit drop. There is correlation between fruit drop and endogenous hormonal status, and existence of high level of internal auxin is useful for preventing fruit drop. Since high level of endogenous hormones might help in building up endogenous hormone at appropriate level that might be potent enough to reduce the fruit drop. Application of zinc could have promoted the auxin synthesis in the plant system which might have delayed the formation of abscission layer during early stages of fruit development (Nason and McElroy, 1963).

Treatments	Fruit length (mm)	Fruit breadth (mm)	Fruit weight (g)	Fruit volume(cc)	Fruit set(%)	Fruit drop(%)
T ₁	59.46	64.86	136.43	140.00	60.03	45.66
T ₂	54.73	63.22	123.33	127.33	60.08	48.84
T ₃	55.98	63.29	120.70	126.33	60.89	49.16
T ₄	62.65	66.63	155.07	158.00	61.01	43.62
T ₅	59.14	65.08	140.10	145.67	60.66	44.29
T ₆	55.12	63.45	125.53	130.67	60.33	48.52
T ₇	64.33	68.03	165.37	168.33	61.32	42.08
T ₀	55.16	61.12	116.93	121.67	61.34	58.60
CD _{0.05}	4.87	2.49	2.41	4.88	NS	1.91

Table 2. Effect of micronutrients on fruit characteristics (physical) of guava

Chemical parameters of fruits

The result revealed that application of micronutrients on chemical parameters of guava. Data pertaining to the effect of micronutrients application on total soluble solids in fruits are given in Table 3. It is evident from the data that the application of micronutrients exerted a significant influence on total

soluble solid contents of fruits. Maximum total soluble solids (10.63° B) were recorded in fruits of the plants treated with. The minimum total soluble solids (8.45° B) was, however, found in T₀ (control). The increase in TSS has also been recorded by Yadav *et al.* (2018), Awasthi and Lal (2009) and Singh *et al.* (2004) in guava. Increase in total soluble solids might be due to the fact that boron helped in trans-membrane sugar transport. Micronutrients play important role in vital plant metabolic functions such as chlorophyll synthesis, various enzymatic reaction, respiration and photosynthesis. Given that the main product of photosynthesis is sugar, so increasing the photosynthesis, lead to increase the sugar compounds and cause more soluble solids in fruit juice (Ram and Bose, 2000).

The scrutiny of data given in Table 3 indicate that application of different micronutrients and their combinations exerted significant effect on total sugars during the course of present study. Highest total sugars (7.51%) were induced in fruits under T₇. While, lowest total sugars (5.74%) was recorded with T₀ (control).

Statistical analysis of data in relation to ascorbic acid content of guava provides substantiation that different micronutrients application individually or in combinations pronouncedly increased the ascorbic acid content as per Table 3. Maximum ascorbic acid content (161.77 mg/100 g pulp) was recorded with T₇. Whereas, minimum ascorbic acid content (140.04 mg/100 g pulp) was found in T₀ (control).

Above observations of the present study indicate that the total sugars, reducing sugars, non-reducing sugars and ascorbic acid content were significantly affected by micronutrients applications alone or in combination (Table 3). These results are in conformity with the results achieved by Kumar *et al.*, (2015) and Jat and Kacha (2014). According to them micronutrients application increased the quality parameters of guava fruit. The increase in the sugars content might be due to the more rapid translocation of sugars from leaves to developing fruits. Boron facilitated sugar transport within the plant. It was also reported that borate reacted with sugar to form a sugar-borate complex. Boron acted as a switcher in the degradation of glucose either by glycolysis or by pentose sugar path way (Singh and Kaur 2016). The improvement in quality of fruit might be due to the catalytic action of micronutrients particularly at higher concentrations. The foliar application of micronutrients quickly increased the uptake of macronutrients in the tissues and organs and improves fruit quality. The boron also plays an important role in activating the synthesis of ascorbic acid (Baranwal *et al.*, 2017). The present results on ascorbic acid is in conformity with the results achieved by Jeyabaskaran and Pandey (2008).

Table 3. Effect of micronutrients on TSS and titratable acidity of guava

Treatments	TSS (° B)	Total sugar (%)	Ascorbic acid
------------	-----------	-----------------	---------------

			(mg/100 g pulp)
T₁	10.63	7.12	164.24
T₂	9.66	6.69	147.66
T₃	9.50	6.76	146.79
T₄	10.26	7.05	159.49
T₅	9.90	6.66	156.40
T₆	9.75	6.53	152.79
T₇	10.56	7.51	161.77
T₀	8.45	5.74	140.04
CD_{0.05}	0.40	0.35	6.46

UNDER PEER REVIEW

YIELD PARAMETERS:

Data pertaining to the effect of different micronutrients and their combinations on number of fruits per plant in guava is summarized in Table 4. The data indicated that the number of fruits per plant in guava increased significantly with the foliar application of micronutrients. The maximum number of fruits per plant (45.33) was recorded under treatment T₇. The minimum number of fruits per plant (37.00) was recorded under treatment T₀ (control).

It is clear from the data given in Table 4 that the application of micronutrients exerted a significant effect on fruit yield per plant during the course of study. Maximum fruit yield (7.50 kg/ tree) was observed under T₇. While the minimum fruit yield (4.33 kg/ tree) was recorded in treatment T₀ (control), which was significantly lower than all other treatments.

The application of micronutrients exerted significant influence on yield efficiency (kg/cm² TCSA) during the course of study. The data in the Table 3 showed that the highest yield efficiency of 0.35 kg/cm² TCSA was recorded in T₇ and the lowest yield efficiency of 0.16 kg/cm² TCSA was recorded in T₅. As far as the effect of micronutrients on guava yield and yield efficiency is concerned, the increase in fruit yield with the application of micronutrients may be attributed to increased fruit size, fruit weight and minimum fruit drop resulting from the effects of micronutrients on cell division, cell elongation and translocation of photosynthetic and metabolites from leaves and others parts of plants to the developing fruits. The highest fruit yield which was obtained by foliar spray of micronutrients may be attributed to better uptake and mobilization of nutrients to the sink which caused better fruit development. These findings are also supported by earlier reports of Bagali *et al.* (1993), Rajkumar *et al.* (2014), Jat and Kacha (2014), and Gaur *et al.* (2014a) who also found that foliar application of micronutrients increases the yield of guava.

Table 4: Effect of micronutrients on number of fruits per plant, yield and yield efficiency of guava

Treatments	Number of fruits/plants	Yield (kg/tree)	Yield efficiency (kg/cm ² TCSA)
T ₁	42.67	5.82	0.23
T ₂	40.67	4.91	0.20
T ₃	44.33	6.87	0.27
T ₄	43.33	6.07	0.24

T ₅	41.33	5.19	0.16
T ₆	46.00	7.35	0.34
T ₇	45.33	7.50	0.35
T ₀ (control)	37.00	4.33	0.25
CD (5%)	1.91	0.31	0.02

1. CD has been calculated based on percentage value

Conclusion

On the basis of the results obtained in the present investigation, it is concluded the combined foliar application of zinc sulphate (ZnSO₄) @ 0.5% + manganese (MnSO₄) (0.5%) + iron sulphate (FeSO₄) (0.4%) (T₇). Therefore, combined spray of ZnSO₄ (0.5%) + MnSO₄ (0.5%) + FeSO₄ (0.4%) can be advocated to guava growers for maximum physical, chemical and yield parameters of guava.

References:

- Ali, W.; Pathak, R.A. and Yadav, A.L. (1993). Effect of foliar application of nutrients on guava (*Psidium guajava* L.) cv. Allahabad Safeda. *Prog. Hort.*, 23 (1-4): 14-21.
- Aulakh, P.S. (2004). Effect of seasonal variation on yield and fruit quality of some promising guava cultivars under arid irrigated region of Punjab. *Haryana J. Hort. Sci.* **33**: 170-71.
- Awasthi P and Lal S. 2009. Effect of calcium, boron and zinc foliar sprays on the yield and quality of guava (*Psidium guajava* L.). *Pantnagar Journal of Research* 7:223-25.
- Bagali AN, Hulamani NC and Sulikeri GS. 1993. Effect of foliar application of zinc, magnesium and boron on growth and yield of guava (*Psidium guajava* L.) cv. Sardar. *Karnataka Journal of Agricultural Sciences* 6:137-41.
- Baranwal D, Tomar S, Singh JP and Maurya JK. 2017. Effect of foliar application of zinc and boron on fruit growth, yield and quality of winter season guava (*Psidium guajava* L.). *International Journal of Current Microbiology and Applied Sciences* 6:1525-29.
- Bose TK, Mitra SK and Sanyal D. 1990. Fruits: tropical & sub-tropical. Naya Prakashan, Calcutta, West Bengal. 838p.
- Das BC, Chakraborty A, Chakraborty PK, Maiti A, Mandal S and Ghosh S. 1995. Comparative performance of guava cultivars under red and laterite soils of West Bengal. *Horticultural Journal* 8:141-46.

- Dhaliwal GS, Nanra NK and Rattanpal HS. 2002. Effect of chemicals on flower drop, fruit set and yield on rainy and winter season crops of guava. *Indian Journal of Horticulture* 59:31-33.
- Gaur B, Beer K, Hada TS, Kanth N and Syamal MM. 2014a. Studies on the effect of foliar application of nutrients and GA₃ on fruit yield and quality of winter season guava. *The Ecoscan* 6:479-83.
- Gaur B, Hada TS, Beer K, Kanth N and Syamal MM. 2014b. Studies on the effect of foliar application of nutrients and GA₃ on yield and reproductive parameters of winter season guava. *Trends in Biosciences* 7:3386-89.
- Ghosh SN. 1986. Effect of magnesium, zinc and manganese on yield and fruit quality of guava. *South Indian Horticulture* 34:327-30.
- Jat G, Kacha HL. 2014. Response of guava to foliar application of urea and zinc on fruit set, yield and quality. *Journal of AgriSearch* 1:86-91.
- Jeyabaskaran KJ and Pandey SD. 2008. Effect of foliar spray of micronutrients in banana under high soil pH condition. *Indian Journal of Horticulture* 65:102-05.
- Kaur G and Dhillon WS. 2006. Effect of foliar application of chemicals on physico and chemical characters of variety Allahabad Safeda of guava during winter. *Journal of Reseach* 43:114-16.
- Kaur S. 2017. Effect of micronutrients and plant growth regulators on fruit set, fruit retention, yield and quality attributes in litchi cultivar Dehradun. *Chemical Science Review and Letters* 6:982-86.
- Kumawat KL, Sarolia DK and Shukla AK. 2012. Growth, yield and quality of rejuvenated guava as influenced by thinning-bending and micronutrients. *Indian Journal of Horticulture* 69:478-83.
- Manoj Gaund, Dr. D Ram, Anand Singh Rawat and Ajendra Kumar 2022. Response of foliar application of micronutrients and plant growth regulator on yield and economic feasibility of guava (*Psidium guajava* L.) CV. Shweta and Lalit. *The Pharma Innovation Journal*, 11(3): 1752-1756.
- Meena D, Tiwari R and Singh OP. 2014. Effect of nutrient spray on growth, fruit yield and quality of aonla. *Annals of Plant and Soil Research* 16:242-45.
- NHB. State wise area, production and productivity of guava. 2017. www.nhb.gov.in.
- Nijjar GS. 1990. *Nutrition of fruit trees*. 2nd ed. Kalyani Publishers, Ludhiana. 311p.
- O' kelley JC. 1957. Boron effects on growth, oxygen uptake and sugar absorption by germinating pollen. *American Journal of Botany* 44:239-44.
- Pal A, Pathak RK, Pal K and Singh T. 2008. Effect of foliar application of nutrients on yield and quality of guava (*Psidium guajava* L.) fruit cv. Sardar. *Progressive Research* 3:89- 90.
- Panse, V.G. and Sukhatme, P.V. (1995). *Statistical methods for agricultural workers*. Indian Council of Agricultural Research Publications, New Delhi. 330

- Price CA, Clark HE and Funkhouser EA. 1972. Functions of micronutrients in plants. *In: Micronutrients in agriculture. Soil Science Society America, Madison, Wisconsin.* pp. 173-88.
- Rout GR and Sahoo S. 2015. Role of iron in plant growth and metabolism. *Reviews in Agriculture Science* 3:1-24.
- Ruby Rani and Brahmachari, V.S. (2004). Effect of growth substance and calcium compounds on fruit retention, growth and yield of Amrapali mango. *Orissa J. Hort.*, 32 (1): 15-18.
- Shukla AK, Sarolia DK, Kumari B, Kaushik RA, Mahawer LN and Bairwa HL. 2009. Evaluation of substrate dynamics for integrated nutrient management under high density planting of guava cv. Sardar. *Indian Journal of Horticulture* 66:461-64.
- Singh R, Chaturvedi OP, Singh R. 2004. Effect of pre-harvest spray of zinc, boron and calcium on the physico-chemical quality of guava fruit (*Psidium guajava* L.). *Internal Seminar on Recent Trend in Hi-Tech Horticulture and Post-Harvest Technology* pp. 4-6.
- Yadav A, Ram RB, Verma RS, Kumar V and Yadav RK. 2018. Effect of foliar application of micronutrients on bio-chemical attributes of winter season guava (*Psidium guajava* L.) cv. Lalit. *Journal of Pharmacognosy and Phytochemistry* 7:3196-97.

UNDER PEER REVIEW