

Original Research Article

Thirteen Year Long Term Fertilization Effect on Soil Phosphorus (P) Fractions of an Acid *Inceptisol* and Their Contribution to P Uptake by a Double Crop of Rice under Sub-Tropical Climate

ABSTRACT

A 13 year old long term fertilizer experiment conducted on an acid soil with rice-rice system was used to study the impact of continuous application of fertilizer nutrients and amendments on changes in soil phosphorous in soil phosphorous fractions and their relative contribution to P nutrition. There was accumulation of total P on surface soil in all P fertilized treatments and depletion in P minus treatments. Depending on the P balance the treatments differed significantly with respect to P accumulation. 150%NPK treatment had highest P accumulation ($841.92 \text{ kg ha}^{-1}$) in surface soil. The high yielding treatments viz. 100% NPK + FYM and 100% NPK + FYM + Lime has relatively less P built up than 100% NPK. The total P comprised of inorganic P 64.43 - 89.60% and organic P 10.40–35.57%. The inorganic fraction in terms of their abundance were in the order Sal-P (42.13 kg ha^{-1}) < Occl-P (52.16 kg ha^{-1}) < Al-P (52.98 kg ha^{-1}) < Ca-P (55.44 kg ha^{-1}) < Red-P ($134.22 \text{ kg ha}^{-1}$) < Fe-P ($218.70 \text{ kg ha}^{-1}$). Olsen P which was significantly influenced by the fertilizer treatments had a very strong positive correlation ($p < 0.01$) with Sal-P ($r = 0.469^{***}$), Occl-P ($r = 0.639^{***}$) and Ca-P ($r = 0.739^{***}$). Among the inorganic P fractions, Ca-P contributed highest of 42.57% to P uptake, followed by Occl-P (28.59%), Sal-P (18.56%) and Al-P (7.63%). The fractions in order of their relative importance for P uptake under submerge rice-rice situation are Ca-P > Occl-P > Sal-P > Al-P > Red > Fe-P and application of P fertilizer not only increased the rice yield, but also improved yield stability. The study indicated that 100% NPK + FYM and 100% NPK + FYM + Lime are the best treatments for maintaining P more active fraction of P required for P nutrition of rice-rice system under the tropical acid situation.

Keywords: Rice-rice system; Phosphorous fractions; Phosphorous uptake; long term fertilization; FYM.

1. INTRODUCTION

Rice–rice cropping system is most prevalent across a major portion of India as well as South Asia, especially among small and marginal farmers. It is usually practiced by farmers where either sufficient irrigation is available or in favorable lowland rain fed areas [1, 2]. In addition to irrigation water availability, high consumer demand, a relatively stable market price, and assurance of a

minimum support price by the government encourage the farmers to grow two crops of rice continuously in consecutive seasons.

In many of these rice production areas, particularly in highly weathered soils of low latitude regions phosphorus(P) has become a limiting nutrient after nitrogen (N) [3]. Although the total P content of soils may be large, only a small part of it is available for plant uptake [4, 5]. Most arable soils cannot supply sufficient amount of the element to the crops. Therefore P fertilizer is applied every season to overcome the deficiency. When fertilizer is applied to soil, only a small amount(10-30%) is taken up by the crop and a major part remains in soil which is not available to the present crop [6,7]. This has led to P enrichment of the top soil of agricultural lands [8, 9] as P is almost immobile in soil. The residual P however, can gradually be available to crop depending upon soil type and soil properties [10], weather condition [4], cropping system and management practices.

Phosphorus in soil is present in both inorganic and organic forms. The inorganic P constitutes a major part of around 58-84% of total P present in soil [11] and comprises of many forms which remain in association with various mineral elements like Al, Fe, Ca and in many physical positions like occluded P and reductant P which unequally contribute to P nutrition depending on the soil and plant environment. In soils, rice crop is grown mostly under submerged condition where there is transformation of P fraction from one form to other depending on soil characteristics which are also influenced by the management practices including application of fertilizers and manure's [12]. In acid soils, P is mainly sorbed to iron (Fe) and aluminium (Al) oxides and hydroxides, and in calcareous soils to calcium (Ca) carbonates [10]. For optimizing P management in crops in a particular agro ecosystem, it is very important to characterize the residual P remaining in the soil after repeated fertilizer P application. Long term field experiments are essential for providing key information on the impacts of management practices in soil and for assessing the sustainability of agro-ecosystem [13]. However, limited long term experimental work has been done in relation to the impacts of fertilization soil P-changes and its dynamics in low land paddy fields.

Information on the composition of all chemical forms of P is fundamental to understanding of P dynamics and interactions in paddy soils, which in turn are necessary for its effective management. However, little is known about the cycling of P in soil, including the fate of P in long term fertilizer and manure applied to lowland paddy soils. Therefore, this paper aims to evaluate the effect of 13 years of Fertilizer and manure application on the transformation of the inorganic P fractions in intensively cultivated lowland paddy soil and their relative contribution to P nutrition using a long term fertilizer experiment with rice-rice production system under subtropical climatic situation of coastal Odisha.

2. MATERIALS AND METHODS

2.1 Experimental site

The study was conducted in the experimental field of All India Coordinated Research Project (AICRP) on LTFE of ICAR at OUAT, Bhubaneswar, India (20°17' N, 85°49' E and 30 m above mean sea level) which was started since 2005-06. The location of the experimental site is characterized as sub-humid subtropical climate with dry season from October to June and wet season from July to September.

The average annual rainfall is 1628 mm, and the mean maximum and minimum temperatures are 31.40 °C and 21.10 °C, respectively. The experimental soil is a pale yellow (10Y RR6/8), lateritic *Inceptisols* (*Udic Ustochrept*) sandy loam. The initial soil properties of 0-15cm layer were pH 5.3, Bulk density 1.55 g cc⁻¹, cation exchange capacity 3.75 cmol(+) kg⁻¹, soil organic carbon (SOC) 4.4 g kg⁻¹, total P, 632 kg ha⁻¹, and available (Olsen) P, 19.7 kg ha⁻¹.

2.2 Experimental details

The experiment consisted of 12 manurial treatments viz., 100% PK, T₂=100% NPK, 150% NPK, 100% NPK + Zn, 100% NPK + FYM, 100% NPK + Lime + FYM, 100% NPK + B + Zn, 100% NPK + S + Zn, 100% N, 100% NP, 100% NPK + Lime and control, where 100% NPK correspond to 80-40-60 kg of N, P₂O₅ and K₂O ha⁻¹. The experiment was laid out in randomized block design (RBD) with four replications. Rice cultivar Swarna (MTU 7029) was grown in wet season and Lalat in dry season of every year. Twenty five days old rice seedlings were transplanted at a spacing of 20 cm × 10 cm with 2-3 seedlings per hill to puddled field in both the seasons. Nitrogen (N) was applied in three splits i.e. 25% at puddling as basal, 50% top dressing at 18 days after transplanting and 25% top dressing at panicle initiation stage. Entire dose of phosphorus (P) was applied during puddling as basal and potassium (K) was applied in two splits, 50% at puddling as basal and 50% top dressing at panicle initiation (PI) stage. Entire FYM (5 t ha⁻¹ season⁻¹) was applied at the time of puddling. Necessary uniform intercultural, water management and plant protection measures were undertaken in general until the crop was matured for harvesting. For the research work, intensive study was conducted on crops of two season dry, 2017-18 and wet, 2018 where 2 different rice varieties Lalat (120 days duration) and Swarna (145 days duration) were grown. Before harvest of crop grain yield was monitored through crop cutting.

2.3 Biomass yield and P content in biomass

After harvesting in both the seasons, biomass yield and phosphorus content in both grain and straw was determined. Both grain and straw yields were estimated after weighing of air dried sample and making adjustment for moisture content. Laboratory analysis for P content was done after drying the sample at 70 °C in oven for 72 hours. The dried plant sample was digested in di-acid for determination of phosphorus content [14].

For determination of P content in grain and straw, plant sample (1g) was poured to 10ml HNO₃ taken in a 150 ml conical flask, which was kept overnight for pre digestion and then heat digested the next day on a hot plate for which 10ml of di-acid was added and gradually heated until just 2-3 ml of samples remained in the flask. The flask was rinsed with distilled water, and the solution was filtered into a 100 ml of volumetric flask then distilled water used for volume makeup. Vanado-molybdo-phosphoric acid reagent was used to estimate total P of soil /grain/straw, 5ml aliquot of 25 ml volumetric flask, and the strength of yellow colour (Vanado-molybdo-phosphoric) was measured at 470nm in a spectrometer and content of P was estimated. Total phosphorous removed by the crop (grain and straw) in both the seasons was calculated separately and then total P removed from the soil in the year was calculated.

2.4 Soil analysis

Total P in soil was also determined following the method similar to plant P estimation outlined by Jackson [14]. Besides total P, fractions of inorganic P (Saloid-P, Fe-P, Al-P, Ca-P, Reductant soluble-P, Occluded bound-P), were determined directly by sequential extraction method given by Peterson and Corey [15] modified by Kuo [16].

2.5 Extraction of P

2.5.1 Saloid bound Phosphorous (Sal-P): 1g of soil was placed in a 50 ml centrifuge tube, 25ml of 1M NH₄Cl solution was added, and shaken for 30 minutes and centrifuged for 10 minutes @ 2000 rpm. Aliquot from the supernatant solution was taken for spectroscopic measurement.

2.5.2 Aluminium bound Phosphorous (Al-P): The residue left was shaken for an hr with 25ml of NH₄F solution (pH 8.2) and then centrifuged. The residue was washed twice with 20ml of saturated NaCl solution, centrifuged each time to recover the soil. The washing was discarded.

2.5.3 Iron bound Phosphorous (Fe-P): The soil was shaken for 17 hours with 50ml of 0.1 M NaOH solution and then centrifuged. In the supernatant solution, concentrated H₂SO₄ was added until the organic colloids begin to flocculate and the suspension was taken for measurement. The precipitate was washed twice in saturated NaCl and the washings were discarded.

2.5.4 Reductant soluble P (Red-P): After that, the soil was suspended in 25ml of 0.3 M sodium citrate solution and 0.3g of sodium di-thionate was added, and the mixture was agitated for 10 minutes. The solution was heated in a hot water bath to 80°C, diluted to 50 mL, agitated for 5 minutes, and centrifuged. Excess citrate and dithionate were oxidised with 0.25 M KMnO₄.

2.5.5 Occluded bound Phosphorous (Occl-P): The residue was washed again, and the washing was disposed in the same manner as before. Shaking the soil with 0.1 M NaOH for an hour.

2.5.6 Calcium bound Phosphorous (Ca-P): The remaining residue was washed twice more with saturated NaCl. Ca-P was collected by shaking the soil for an hour with 0.5ml of 0.25 M H₂SO₄. Total organic P was indirectly determined from difference between total P and inorganic P.

2.6 Estimation

5ml of the extracted aliquot was taken in 25ml of volumetric flask with 2-3 drops of the p-nitrophenol indicator were added and the pH adjusted with 2 M H₂SO₄ until the indicator colour just changed and volume makeup by distilled water. Phosphorus concentrations were estimated using the ascorbic acid method at 480nm [17].

The amount of P in each fraction was calculated using the following equation:

P concentration in given fraction (mg kg⁻¹) =

$$[\text{Conc. of P (mg L}^{-1}\text{)}] \times [\text{R factor}] \times [\text{Volume of extractant (L)} \div \text{mass of soil (kg)}]$$

The total organic P was indirectly determined by from difference between total P and inorganic P.

$$\text{Total organic P (kg ha}^{-1}\text{)} = \text{Total P (kg ha}^{-1}\text{)} - \text{Total inorganic P (kg ha}^{-1}\text{)}$$

$$\text{Total inorganic P (kg ha}^{-1}\text{)} = \text{Sal-P} + \text{Fe-P} + \text{Al-P} + \text{Red-P} + \text{Occl-P} + \text{Ca-P}$$

Besides total and fractions of P soil was also analyzed for Olsen's P, pH, SOC, CEC, and Clay % by standard methods [14].

2.7 Statistical analysis

Two –way analysis of variance was carried out to that the effect of various treatments on the study variables. DMRT was performed to that the significance of pairwise mean difference among various treatments. Karl Pearson's product moment correlation analysis was carried out to find out the relationship between various parameters soil P fractions with Grain yield, P uptake and soil properties. The correlation coefficient were tested at 1% and 5% level of significance. A multiple linear regression analysis was carried out to establish underline relationship between P uptake (dependent variable) and various P fractions (Independent variables)

$$U = b_{Sal} + b_{Al} + b_{Fe} + b_{Red} + b_{Occl} + b_{Ca}$$

Where, **U** stand for phosohorous uptake , **Sal** stand for saloid bound phosphorous, **Al** stand for Aluminium bound phosphorous, **Red** stands for Reductant bound phosphorous, **Occl** stands for Occluded bound phosphorous, and **Ca** stand for Calcium bound phosphorous.

Furthermore, a Hierarchical partitioning analysis [18, 19] was carried out to find out relative importance of various P fractions on uptake. All the statistical analysis was carried out using R statistical package.

3. RESULTS AND DISCUSSION

3.1 Grain yield

3.1.1 Dry season, 2017-18

Result on grain yield of dry season 2017-18 showed [Table1. ; Fig.1.(a-c)] that highest yield of 4453 kg ha⁻¹ was produced in 100% NPK + Lime + FYM treatment which was at par with 100% NPK + FYM (4347 kg ha⁻¹) demonstrating no effect of lime in presence of FYM applied @ 5 t ha⁻¹. But in absence of FYM, liming has significant effect 3525kg ha⁻¹. P or K also significantly increased the grain yield. Application of 50% more NPK also was effective in producing 13.26% significantly higher yield than 100% NPK. Application of zinc did not produce any significant effect over 100% NPK. There was no significant response to application of B and S. Grain yield was significantly lower in control plot (1325 kg ha⁻¹) than all other treatments except 100% N that also produced vary low yield of 1751 kg ha⁻¹.

3.1.2 Wet season, 2018

The data revealed that the yield of wet season 2018 (Table 1.; Fig.1.b) rice (cv. Swarna) varied from a minimum of 17.77 q ha⁻¹ recorded in control to a maximum 4891 kg ha⁻¹ in 100% NPK + FYM + Lime treatments. Application of FYM @ 5 t ha⁻¹ was very effective in significantly increasing 37.44% raise in the grain yield over 100% NPK, whereas lime @1 t ha⁻¹ has no effect when applied along with FYM. Application of 50% more NPK also was effective in producing 10.04% significantly higher yield than 100% NPK. Application of zinc did not produce any significant effect over 100% NPK. Further, in the present investigation, application of B and S did not produce any significant effect over 100% NPK + Zn. Working on similar soil, [20] also found that continuous addition or exclusion of some secondary

(S) and micro-nutrients (Zn and B) did not make any significant difference on grain yield. From the results it is also clear that FYM has significant effect on grain yield in both the seasons.

Combination of FYM + Lime to 100% NPK resulted in significant increase in grain yield in both the seasons in all the years. However, the response to FYM was more in wet season than in dry season. Higher response to FYM in the wet season than dry season has also been reported by Majhi and Rout [21]; Shahid et al [22]. Significant yield increase by 22.10% was also caused by 150% NPK over 100% NPK. In a study conducted on *typic Ustochrept* on a clayey soil of Andhra Pradesh. Srilatha et al. [23] also reported more yield with 150% NPK than 100% NPK + FYM.

3.2 Phosphorous (P) -uptake

The crop yield and uptake of nutrients are interdependent. The total uptake of nutrients for rice was calculated by adding the nutrient uptake [Table 1; Fig. 2. (a-c)] by both grain and straw yield of individual season. Results pertaining to total P uptake of dry season 2017-18 wet season 2018 and are presented in Table 1. Fig. 2. (a and b). Total P uptake in dry season (2017-18) varied from 4.00 kg ha⁻¹ in control to 16.89 kg ha⁻¹ and in wet season 2018 varied from 3.87 kg ha⁻¹ in control to 20.55 kg ha⁻¹ in 100% NPK+ Lime + FYM. 100% NPK + FYM was at par with 100% NPK + Lime + FYM as presented in Table.1. FYM amended plots resulted in more P uptake by releasing the organic acids during its decomposition [24] P uptake was more in all P treated plots than P minus treatments. Significantly higher uptake was recorded with super optimal dose (150% NPK) than 100% NPK. No significant effect was observed for secondary (S) and micronutrients (Zn and B) applied with optimal dose. Uptake of nutrients was lower in the control plot due to absence of external source of nutrient to the plants [25].

3.3 Available Phosphorous (Olsen's P)

Results on available P (Olsen's P) presented in Table 1. reveal that continuous application of FYM along with optimal level of NPK either with or without lime maintained significantly higher quantity of Olsen's P than non FYM and non-lime treatments. Highest available P (53.41 kg ha⁻¹) was measured in NPK + FYM + Lime treatment which was at par with NPK + FYM treatment (49.72 kg ha⁻¹). P content increase in pH with application of FYM and lime favors P desorption from clay, iron and aluminum oxides. Such pH changes when found to favor desorption of freshly applied P only [26].

3.4 Changes in P fractions

Data pertaining to various P fractions recorded in Table 2. reveal that after 13 years of continuous cropping the total inorganic P (Pi) varied from 337.73 kg ha⁻¹ in control plot to a highest of 684.96 kg ha⁻¹ in 150% NPK treatment and organic P (Po) varied from a lowest of 48.58 kg ha⁻¹ in 100% N plot to a highest of 204.29 kg ha⁻¹ in 100% NPK + FYM treatment and control 186.45 kg ha⁻¹. The Pi fraction constituted 64.43-89.60% where Po fraction 10.40-35.57% of the total phosphorous. The inorganic fractions on an average thus constituted 76.46 % and organic P, 23.53 % of the total P under the rice –rice ecosystem of eastern India (Fig.3). The organic P constituted one third of the total inorganic P in the intensively rice grown and soil of subtropical condition. In a study under similar

situation reported total inorganic phosphorous (Sal-P, Fe-P, Al-P, Red-P, Occl-P, Ca-P) constituting 58-84% of total P present in soil [11]. The lowest value of all inorganic fraction of P observed under control i.e. cultivation without fertilizers might be due to continuous removal of P from soil P reserve without any replenishment through fertilizers [25].

After 13 years Sal-P ranged from 20.35 to 62.98 kg ha⁻¹, Fe-P 152.89 to 288.38 kg ha⁻¹, Al-P, 35.07 to 61.75 kg ha⁻¹, Ca-P, 24.87 to 89.32 kg ha⁻¹, Red-P 79-152 kg ha⁻¹, Occl-P 25.19 to 77.86 kg ha⁻¹. From the result it is observed that the content of Sal-P is the least among all the fractions which has also been reported by many workers [27, 28]. The fractions in order of their content are: Fe-P > Red-P > Ca-P > Al-P > Occl-P > Sal-P, contradictory trend of P fraction, Fe-P > Red-P > Al-P > Ca-P > Occl-P > Sal-P under *Inceptisol* [29]. The increase of Fe-P than Al-P due to the availability of NaOH extractable inorganic P serves as primary sink for applied P fertilizer in tropical soil and also in other cropping systems in various soils under continuous cropping system [30]. After 13 years of continuous cropping the available P (Olsen's P) ranged between 7.89 and 53.41 kg ha⁻¹ as compared to the initial P of 19.14 kg ha⁻¹.

The higher amount of Ca-P in inorganic P fraction with respect to Al-P due to application of N, K and organic fertilizer will accelerate the P fixation equally in Fe-P and Ca-P sites at nearly neutral pH under flooded condition [31]. The fractions of inorganic P in soil and its availability thus varied to a great extent due to different long term manorial management practices followed on a particular soil and climatic situation.

3.5 Long term treatments effects on various P fractions in an acidic inceptisol under wet land rice-rice system

3.5.1 Saloid bound Phosphorous (Sal-P)

Saloid P is the smallest fraction among all inorganic P fractions in surface soil. It constituted 6.13-11.31% of total inorganic P and 3.88-8.23% of total P. After 13 years of continuous cropping and comparisons of mean of major effect of different manorial treatments on Sal-P (Table 2; Fig.4.) showed that the highest content is recorded in 100% NPK + FYM (62.98%) and lowest in 100% N (27.16%) as compared to the initial status of 22.35 kg ha⁻¹ which is 5.83% of total inorganic P and 3.54% of total P present in soil this increase in Sal-P concentration was more with the application of organic manure than inorganic fertilizers due to release of P from organic matter through mineralization [32]. Application of FYM @ 5 t ha⁻¹ season⁻¹ increased the Sal-P by 54.17% in absence of lime and 34.26% in presence of lime applied @ 1 t ha⁻¹ indicating negative interaction effect of lime and FYM on Sal-P. Application of Zn @ 0.4% ZnO root dip also increased Sal-P by 9.67%. Zinc and phosphorus interaction lead to highly significant increase in the uptake of phosphorus and zinc in wheat and rice crops [33, 34]. Combination of S or B with Zn however, in our study had negative effect on Sal-P. In contrast both N and K had positive effect on Sal-P. Thirteen years of continuous application of N and K caused 33.46% and 26.74% increase in Sal-P respectively. Similar effect of N and K was also reported by Kaur [35].

3.5.2 Iron bound Phosphorous (Fe-P)

Results on Fe-P content of surface soil measured after 13 years of cropping (Table. 2) revealed that it is the largest fraction among all the inorganic fractions of P constituting 30.53 - 45.75 % of total inorganic P and 22.22 - 37.46% of total P in soil. The results of comparison of mean values and the ANOVA test result showed that the treatments had significant effect on Fe-P content under the prevailing soil situation. There was significant increase in its content in all fertilized treatments except 100% N, 100% NPK + FYM and 100% NPK + FYM + Lime treatments where the initial content (172 kg ha^{-1}) remained almost unchanged. P added through mineral fertilizer to acidic soils gradually reacts with Fe and Al-P compound and is transformed into relatively insoluble P compounds [36]. The treatment 150% NPK registered highest Fe-P of 288 kg ha^{-1} which was 21.01 % more than the optimum level (100% NPK) 88% more than control ($152.89 \text{ kg ha}^{-1}$) and 67.66% over the initial 172 kg ha^{-1} . Combination of FYM (5 t ha^{-1}), however, resulted in decrease in Fe-P content by 28.55% over 100% NPK. Organic matter in soil causes ferric iron reduction through its promoting influence on the bacterial activity in flooded soil. Humus and humus forming materials also help to decrease the fixation of P ions on Fe and Al oxides and maintain a steady availability of P to plants [37]. Application of compost decreased the residual P and Fe-P fraction in acid soil [38].

3.5.3 Aluminum bound phosphorous (Al-P)

Results on Al-P content of surface soil measured after 13 years of cropping (Table 2.) revealed that it is the 4th largest fraction among all the inorganic fractions of P that varied from 35.07 kg ha^{-1} in control plot to a highest of 61.75 kg ha^{-1} in 150% NPK treatment constituting 8.36-11.15 % of total inorganic P and 6.20-9.05% with total P in soil. The results of comparison of mean values and the ANOVA test showed that the treatments had significant effect on Al-P content under the prevailing soil situation. Continuous application of N fertilizer on rice field increased the Al-P content by 12.71% over control. Potassium and zinc increased the Al-P content by 24.37% and 2.86% respectively over 100%NPK. In contrast, secondary and micro nutrients (S and B) had negative impact on Al-P availability in rice soil. Continuous application of lime also caused reduction in Al-P content.

It is interesting to note that unlike Fe-P, Al-P was not influenced by application of FYM. In a laboratory study however, showed that FYM @ 20 t ha^{-1} significantly increased saloid-P, Al-P and Fe-P but showed non-significant increase in Ca-P [39].

The Al-P and Fe-P pools are expected to have very low bio-availability but can be used by plants when available soil P is severely low [40].In acidic soils, the original superficial, loosely bound phosphates to Fe and Al-oxides available to plant are converted gradually via a re-precipitation process into highly crystalline Fe and Al-P (not available to plant) [38].

3.5.4 Calcium bound phosphorous (Ca-P)

Ca-P varied from a lowest of 24.87 kg ha^{-1} in control constituting 7.36% total inorganic P to a highest of 89.39 kg ha^{-1} in 100%NPK+FYM+Lime treatment contributing 16.15% and 12.15% of total inorganic

P and total P. Application FYM and lime significantly increased the Ca-P by 64.50% and 45.07% respectively over 100%NPK. Application of Zn as ZnO (0.4%) root dipping increased the Ca-P by 4.91% while secondary and micronutrients reduced the Ca-P content. Addition of K increased the Ca-P by 27.49% while continuous application of N alone in form of urea increased Ca-P by 47.08%. Similar result was reported by Kaur [35].

3.5.5 Reductant soluble Phosphorous (Red-P)

The reductant and occluded P are dominant in acid red laterite soils than neutral-alkaline or black clay soil [41] or due to weathered condition of the soil studied in the soils of Karnataka, Tamilnadu, Gujrat and AP [42]. The results of ANOVA test showed that the treatments had significant effect on Red-P after 13 years of cropping. Red-P varied from lowest of 79 kg ha⁻¹ in control plot to a highest of 152.48 kg ha⁻¹ in 150%NPK treatment contributing 23.92%, 15.14% with control and 18.57%, 14.07 % with 150%NPK to total inorganic P and total P percentage for Red-P respectively due to application of graded dose of NPK fertilizer caused in increase in reductant P over control [43].

Within 13 years, there was increase in Red-P in all fertilized treatments over the initial status, 85.50 kg ha⁻¹ whereas, there was significant decrease in control. With application of organic manure, @5t ha⁻¹ season⁻¹ there was significant decrease in Red-P as compared to 100% NPK due, to interaction of FYM chemicals bind the reductant P strongly [43]. There was decrease by 28.44% to 31.09 % in FYM and FYM + lime amended treatments indicating lesser accumulation in less active form of P in soil. Increase the addition of organic fertilizer will enhance the reductant P fraction in acid soil [44].

With application of lime @1 t CaCO₃ ha⁻¹, Red-P also decreased to a greater extent in absence of FYM. Application of Zn as 0.4% ZnO solution root dipping, there was slight decrease (0.72%) in Red-P. Joint application of B or S with Zn however further reduce the Red-P by 0.84%, 4.43% respectively indicating formation of more less active form. Application P or K also had significant effect on built up of this form of less active / recalcitrant form of P on the surface soil. Continuous application of N enhanced the Red-P bound P by 30.62 %. Amount of Red-P in surface soils could be attributed to the weathering of soil.

3.5.6 Occluded bound P (Occl-P)

Occluded P is another less active fraction of P in soil which constituted 7.74-13.98 % of inorganic P and 4.81-10.18% of total P with lowest found with control and highest with 100%NPK+FYM in acid soil. After 13 years of continuous cropping and manuring there was increase in occluded P in all P fertilized treatments with highest recorded in 100% NPK +FYM (77.86 kg ha⁻¹) and lowest in 100% NP treatment (40.53 kg ha⁻¹) as compared to the initial status of 28.31 kg ha⁻¹. Continuous application of lime with FYM (0.34%) and lime alone (4.09%) have much effect in increasing the Occl-P. Amount of occluded Fe-P/Al-P could be attributed to the weathering of the soils. Addition of K and continuous application of N increased Occl-P by 15.74% and 45.45% respectively. Super optimal dose of P enhanced the occluded P by 33.23%. Higher accumulation in 150%NPK and relatively lower occluded

P in high yielding treatments suggest that P bound to Occl-P is used more by rice through higher biomass production in high yielding treatments viz. 100% NPK + FYM and 100% NPK + FYM + Lime.

3.5.7 Total Phosphorous (Total P)

Data on changes in total P of surface soil layer (0-15cm) (Table 2.) show that there is accumulation of total P in all P treated soils and depletion in P minus treatments. Within 13 years of continuous cropping without fertilizer resulted in a decrease of 108 kg P ha⁻¹ or 17.08% from the initial 632 kg ha⁻¹. Among the P applied treatments, highest accumulation (841.92 kg ha⁻¹) was found with the super optimal dose (150% NPK) which was 9.77% more than that with optimal dose (100% NPK). Total P content in high yielding treatments such as 100% NPK + FYM and 100% NPK + FYM + lime was less than that of 100% NPK treatment. Addition of recommended dose of K to 100% NP increased the total P by 5.35%. On the other hand, the total P content decreased with application of micro and secondary nutrients (Zn, S and B) in conjunction with 100% NPK. Less accumulation in high yielding treatments is due to more removal by the above ground crop biomass which is displaced from the field.

3.6 Inter relationship of seasonal crop yield (dry and wet) with P uptake in terms of correlation coefficient (R²)

Correlation studies made on the crop yields and P uptake of both dry season (2017-2018) and wet season (2018) depicted in Fig. 5. (a to c) showed highly significant correlation between crop yields and P uptakes of both the seasons and system as whole.

Relationship of various P fractions in soil with crop yield and P uptake P uptake of both the seasons was presented in Table.3. The results reveal that grain yield of rice in dry and wet season have highly significant correlation, with Sal-P (r= 0.580***, 0.642***), Al-P (0.372**, 0.413**) and Ca-P (0.819***, 0.795***) respectively. Similar result was reported by Kaur [35]. Among the various P fractions Ca-P show strongest positive correlation with both grain yield (r= 0.819***, 0.795***) and P uptake (r= 0.810***, 0.857***) in both dry and wet season respectively followed by Occl-P (0.659***, 0.673***) , (0.697***, 0.774***) than Sal-P (r= 0.580***, 0.642***), (0.581***, 0.640***), than Al-P (0.372**, 0.413***) , (0.387**, 0.457***) respectively but, Fe-P and Rd-P and organic P had non-significant correlation.

3.7 Correlation of soil properties with various P fractions measured on postharvest surface soil of wet season 2018

Results of correlation between various fractions of soil P and soil properties (pH, SOC and CEC) in Table 4. among the fractions, Sal-P, Occl-P and Ca-P have significant positive correlation with pH, SOC and CEC, whereas, Fe-P and Red-P have negative correlation with pH and SOC. Total inorganic P, organic P and total P represent non-significant correlation with pH and available P. CEC have strong correlation with inorganic P and total P, while SOC show 5% level of significance with total P.

3.8 Relative contribution of surface soil P fractions to P uptake by rice in different seasons

A multiple regression analysis was carried out taking dry season P uptake as dependent variable and P fractions as independent variable. The overall variation explained by all fractions was 69.75%. From the regression equation it is revealed that among the inorganic P fractions, the effect of Ca-P is highly significant on P uptake (Table 6.; Fig.6.). Furthermore, a relative importance analysis was carried out to find the independent effect percentage of each fraction on dry season P uptake. The results revealed that, Ca-P has maximum independent contribution to P uptake accounting for 45.41 % followed by Occl-P 25.86%, and then Sal-P 18.68% and Al-P 6.90% suggesting these four fractions to be the maximum contributing fractions on dry season P uptake variation. Ca-P > Occl-P > Sal-P > Al-P > Fe-P > Red-P.

Results of multiple regression analysis carried with wet season P uptake showed that the overall variation explained by all fractions was 78.79%. From the regression equation it was revealed that the effect of Ca-P is highly significant on P uptake. Results of relative importance analysis carried out to find the independent effect percentage of each fraction on wet season P uptake showed that, Ca-P has maximum independent contribution accounting for 42.57% followed by occluded P 28.59% and then Sal-P 18.56% and Al-P 7.63% to wet season uptake variation. Table 6. and Fig. 7 shows that independent effect (%) contribution of each fractions on wet season uptake. The regression equation obtained shown Table 5.

A multiple regression analysis was carried out with various inorganic P fractions of surface soil for total P uptake as independent variables showed Table 6; Fig. 8 that the overall variation explained by all fractions was 77.28%. From the regression equation it is revealed that the effect of Ca-P is highly significant on P uptake. Furthermore a relative importance analysis carried out to find the independent effect percentage of each fraction on P uptake revealed that, Ca-P has maximum independent contribution accounting for 43.90%, followed by Occl-P 27.39%, Sal-P 18.64% and Al-P 7.28% to system P uptake. In rice soil, availability of Ca-P has been found in abundant quantity and also required by crop as P uptake, due to submergence condition of rice field raise the soil pH from acidic to slightly acidic to neutral thereby liberate Ca-P as labile form of P as compare to non labile form (Al/Fe oxide and hydroxides) of P present in acid soil [45, 46]. The indirect and direct effect of occluded P as source of P on the available P pool for P nutrition can be attributed to the resolution of the stable forms of inorganic P found as occluded P fraction [47]. Reductant soluble P is consider to be recalcitrant and hence not extractable as is assumed to be unavailable [48].

Thus it is inferred that among the total inorganic P fractions of surface soil, Ca-P contributes maximum to P uptake in both the seasons followed by occluded P then Sal-P and Al-P. Relative contribution of Fe-P and reductant P is minimum. The fractions in order of their relative importance for P uptake under submerged rice-rice situation are: Ca-P > Occl-P > Sal-P > Al-P > Red-P > Fe-P.

4. CONCLUSION

Combination of FYM to 100% NPK resulted in significant increase in grain yield in both dry and wet season. However the response to FYM was more in wet season than in dry season. Application of Zn or Zn+B or Zn+S did not have any significant impact on grain yield. Continuous application of FYM along with optimum level of NPK either with or without lime maintained significantly higher level of Olsen P than non FYM and non Lime treatments. Results showed that there was build-up of total P in all P treated plots and depletion in P minus treatments. Highest accumulation of 841.92 was found in super optimal dose (150% NPK) which was 9.77 more than the optimal dose (100% NPK) but total P content in high yielding treatment such as 100% NPK + FYM and 100% NPK + FYM + Lime was less than that of 100% NPK treatment. Continuous cropping without P resulted in decrease of total P in surface soil. In control the decrease was 17.08% from the initial 632 kg ha⁻¹. The inorganic P fraction on surface soil constituted 64.43-89.60% whereas the organic P fraction 10.40-35.57% of total Phosphorous. Organic P thus constituted about 1/3rd of total inorganic P in the rice-rice system of the eastern India. The inorganic fraction in terms of their abundance were in the order Sal-P (20.35-62.98 kg ha⁻¹) < Occl-P (25.19-77.86 kg ha⁻¹) < Al-P (35.07-61.75 kg ha⁻¹) < Ca-P (24.87-89.39 kg ha⁻¹) < Red -P (79.36-167.83 kg ha⁻¹) < Fe-P (152.89-288.38 kg ha⁻¹). Olsen P which was significantly influenced by the fertilizer treatments had a very strong positive correlation (p<0.01) with Sal-P (r= 0.469***), Occl-P (r=0.639***) and Ca-P (r=0.739***). Among the inorganic P fractions, Ca-P contributed highest of 42.57% to P uptake, followed by Occl-P (28.59%), Sal-P (18.56%), and Al-P (7.63%). Relative contribution of Fe-P and red-P is negligible. So, the fractions in order of their relative importance for P uptake under submerge rice-rice situation are Ca-P > Occl-P > Sal-P > Al-P > Red-P > Fe-P.

REFERENCES

1. Deep M, Kumar RM, Saha S, Singh A. Rice-based cropping systems. Indian Farming. 2018; 68 : 27–30.
2. Lal B, Gautam P, Panda BB, Raja R, Singh T, Tripathi R, Shahid M, Nayak AK. Crop and varietal diversification of rainfed rice based cropping systems for higher productivity and profitability in Eastern India. PLoS ONE. 2017 ; 12 : 175-709.
3. Nishigaki T, Tsujimoto Y, Rinasoa S, Rakotoson T, Andriamananjara A, Razambelo T. Phosphorus uptake of rice plants is affected by phosphorus forms and physicochemical properties of tropical weathered soils. 2019; 435 : 27–38.
4. Syers JK, Johnston AE, Curtin D. Efficiency of soil and fertilizer phosphorus use. Reconciling changing concepts of soil phosphorus behaviour with agronomic information. FAO Fertilizer and Plant Nutrition Bulletin. Food and Agricultural Organization of the United Nations, Rome, Italy. 2008.
5. Krupenikov IA, Boincean BP, Dent DL. The black earth: ecological principles for sustainable agriculture on chernozem soils. Springer, Dordrecht, the Netherlands. 2011.

6. Manske CGB, Ortiz-Monasterio JJ, Van Ginkel M, Gonzalez RM, Rajaram S, Molina E, Vlek PLG. Traits associated with improved P-uptake efficiency in CIMMYT's semi dwarf spring bread wheat grown on an acid Andisol in Mexico. *Plant Soil*. 2000 ; 221 : 189–204.
7. Sarkar S, Bhaduri D, Chakraborty K. Plant adaptation mechanisms in phosphorus-deprived soil: mitigation of stress and way to balanced nutrition. *Adv. Plant Physiol*. 2014; 15: 254–282.
8. Kashem MA, Akinremi OO, and Racz GJ. Phosphorus fractions in soil amended with organic and inorganic phosphorus sources. *Can. J. Soil Sci*. 2004 ;84 : 83–90.
9. Qian P, Schoenau JJ, Wu T, and Mooleki P. Phosphorus amounts and distribution in a Saskatchewan soil after five years of swine and cattle manure application. *Can. J. Soil Sci*. 2004;84 (3):275–281.
10. Brady NC, Weil RR. The nature and properties of soils, in Chapter 14: soil phosphorus and potassium. 11th ed. Prentice-Hall International Inc., Upper Saddle River, NJ, USA. 1999 ;Pp 445–486.
11. Sanyal SK, Dwivedi BS, Singh VK, Majumdar K, Datta SC, Pattnaik SK, and Annapurna K. Phosphorous in relation to domain cropping sequences in India: chemistry, fertility relations and management options. *Current Science*. 2015; 108(7): 1262-1270.
12. Singh M, Reddy KS, Singh VP, Rupa VP. Phosphorous availability to rice (*Oriza sativa* L.) wheat (*Triticum estivum* L.) in a vertisol after eight years of inorganic and organic fertilizer additions. *Bioresource Technology*. 2007; 98:1474-1481.
13. Rasmussen PE, Goulding KWT, Brown JR, Grace PR, Janzen HH, Korschens M. Long-term agroecosystem experiments assessing agricultural sustainability and global change. *Science*. 1998; 282 : 893-896.
14. Jackson ML. Soil chemical analysis. Prentic Hall of India Pvt. Ltd., New Delhi. 1973.
15. Petersen GW, Corey RB. A modified Chang and Jackson procedure for routine fractionation of inorganic soil phosphates. *Soil Science Society of America Proceedings* 1966 ; 30:563-565.
16. Kuo S. Phosphorus. In: Sparks D.L. (Eds.), *Methods of Soil Analysis. Part 3. Chemical Methods. Agronomy Monograph, Vol.9.* Soil Science Society of America, Madiston, WI. 1996 ; 869-919.
17. Murphy J, Riley JP. A modified single solution method for determination of phosphate in natural waters. *Analytica Chimica Acta* 1962 ;27: 31-36.
18. Chevan Albert, Sutherland Michael. Hierarchical Partitioning. *The American Statistician*. 1991; 45(2) : 90-96.
19. Gromping Ulrike. Relative Importance for Linear Regression in R. *Journal of Statistical Software*. 2006; 17: 1-27.
20. Majhi P, Rout KK, Sahoo D, Behra M, Behra BB. Nitrogen fertility of an Inceptisol planted to rice-rice as influenced by three years of contunoius manuring with various organic and inorganic sources, *Journal of Research , OUAT*, 2003; 21(1) :55-61.
21. Majhi P, Rout KK, Mandal M, Singh M, Sasmal N. Soil enzyme activity of anintesively grown rice field as influenced by long term fertilization and manuring under sub tropical climatic condition, *International Journal of Farming and Allied Science*. 2016; 6(3): 172-179.

22. Shahid M, Nayak AK, Shukla AK, Tripathi R, Kumar A, Mohanty S, Bhattacharyya P, Raja R, Panda BB. Long-term effects of fertilizer and manure applications on soil quality and yields in a sub-humid tropical rice-rice system. *Soil Use Management*. 2013; 29: 322–332.
23. Srilatha M, Sharma SHK, Devi M Uma, and Rakha K. Bhanu. Grain yield and soil nutrient status of rice-rice cropping system as influenced by nutrient management under long term fertilizer experimentation. *Journal of Progressive Agriculture*. 2014; 5(1): 85-89.
24. Arulmozhiselvan K, Elayarajan M, and Sathya S. Effect of long term fertilization and manuring on soil fertility, yield and uptake by finger millet on Inceptisol. *Madras Agricultural Journal*. 2013; 100: 490-494.
25. Ghosh Delina, Mandal Mitali, and Pattanayak Sushanta Kr. Long term effect of integrated nutrient management on dynamics of phosphorous in an acid Inceptisols of Tropical India. *Communications in Soil Science and Plant Analysis*. 2021; 52(19): 1-15. DOI:10.1080/00103624.2021.1924186.
26. Willet IR. Causes and prediction of changes in extractable phosphorous during flooding. *Aust J Soil Res*. 1989; 27: 45-54.
27. Pati Ram, Mukhopadhyay D. Inorganic phosphorus fractions in some acid soils under terai situations of West Bengal. *Research on Crops*. 2008; 9:57-60.
28. Adhikari M, Si SK. Distribution of inorganic phosphate fractions in some acid soils of West Bengal. *Journal of the Indian Society of Soil Science*. 1994; 42: 459-61.
29. Prajapati Jaya, Pattanayak SK. Phosphorous Fractionation under Intensive Cropping System in Acidic Soil of Odisha and Correlation Matrix Study with Different Forms of P, *Communications in Soil Science and Plant Analysis*. 2019; 50(14):1-11. DOI:10.1080/00103624.2019.1635142.
30. Motavalli PP, Miles RJ. Soil phosphorus fractions after 111 years of animal manure and fertilizer applications, *Biol. Fert. Soils*. 2002; 36: 35–42.
31. Bhattacharyya P, Nayak AK, Shahid M, Tripathi R, Mohanty S, Kumar A, Swain CK. Effects of 42-year long-term fertilizer management on soil phosphorus availability, fractionation, adsorption–desorption isotherm and plant uptake in flooded tropical rice. *The Crop Journal*. 2015; 3(5): 387-395.
32. Bahl GS, and Singh NT. (1997) The inorganic soil phosphorous fractions and available P as influenced by green manuring cropping and P fertilization. *J Indian Soc. Soil Sci*. 1997; 45: 19-23.
33. Khan AA, Zende GK. Effect of zinc and phosphorus fertilization on the content and uptake of N, P, K, Ca, Mg and Zn by maize and wheat. *Mysore J. Agr. Sci*. 1976;10: 574–584.
34. Tiwari KN, Pathak AN, Upadhyay RL. Studies on Fe and Zn nutrition of rice at varying moisture regimes in a black clay soil of Uttar Pradesh. *J.Indian Soc. Soil Sci*. 1976; 24(3):303-307.
35. Kaur Sukhvir, Brar BS, Dheri GS. Effect of long term use of inorganic and integrated fertilization on soil phosphorous fractions in rice-wheat cropping system. *Agric Res J*. 2015; 52(4): 39-43.

36. Verma S, Subehi SK, Sharma SP. Phosphorus fractions in an acid soil continuously fertilized with mineral and organic fertilizers. *Bio. Fertil. Soils*. 2005;41: 295-300.
37. Bhattacharyya P, Chakrabarti K, Chakraborty A, Nayak DC. Effect of municipal solid waste compost on phosphorous content of rice straw and grain under submerged condition. *Archives of Agronomy and Soil Science*. 2005; 51(4): 363-370.
38. Chang HL, Park CY, Park KD, Jeon WT, Kim PJ. Long-term effects of fertilization on the forms and availability of soil phosphorous in rice paddy. *Chemosphere*. 2004; 56: 299-304.
39. Kumar Sarvesh, Srivastava Ajaya, and Gupta Amit. (2015) Effect of organic amendments on availability of different chemical fractions of phosphorus. *Agric. Sci. Digest*. 2015; 35 (2): 83-88.
40. Shen J, Li R, Zhanga F, Fan J, Tang C, Rengel Z. Crop yields, soil fertility and phosphorous fractions in response to long term fertilization under the rice monoculture system on a calcareous soil. *Field Crop Research*. 2004; 86: 225-238.
41. Kothandaraman GV, and Krishnamurthy KR. Forms of inorganic phosphorus in Tamil Nadu soils. *Phosphorus in Soils, Crops and Fertilizers*. *Bulletin of the Indian Society of Soil Science*. 1979; 12: 608.
42. Kalivanan D, Sudhir K. Phosphorus fraction of selected banana growing soils of India and their relationship with soil characteristics. *Mysore Journal of Agriculture*. 2012; 45: 73-79.
43. Nayak Tripti, Bajpai RK, Sharma Priyanka. Forms of soil phosphorous and depth wise distribution under organic and inorganic nutrient management in a Vertisol planted rice, *Asian J. Soil Sci*. 2015; 10(1): 47-54.
44. Abolfazli F, Forghani A, Norouzi M. Effect of phosphorous and organic fertilizers on phosphorous fractions in submerged soil. *Journal of Soil Science and plant nutrition*. 2012; 12(2): 349-362.
45. Zhang Q, Wang GH, Feng YK, Sun QZ, Witt C, Dobermann A. Changes in soil phosphorous fractions in a calcareous paddy soil under intensive rice cropping. *Plant Soil*. 2006; 288: 141-154.
46. Sah RN, and Mikkelsen DS. Transformation of inorganic phosphorous during the flooding and draining cycles of soil. *Soil Sci, Soc. AM, J*. 1986; 50:62-67.
47. Costa MG, Gama-Rodrigues AC, Gonçalves JLD, Gama-Rodrigues EF, Sales MVD, Aleixo S. Labile and non-labile fractions of phosphorus and its transformations in soil under Eucalyptus plantations, Brazil. *Forests*. 2016.
48. Condon LM, Newman S. Revisiting the fundamentals of phosphorus fractionation of sediments and soils. *Journal of Soils and Sediments*. 2011; 11 (5): 830–840.




Fig. 1 (a)




Fig. 1 (b)




Fig. 1 (c)

Fig.1 Grain yield of both the season dry (a) and wet (b) and total yield per year (c) under various manurial practice




Fig. 2 (a)




Fig. 2 (b)




Fig. 2 (c)

Fig. 2 P uptake of rice in both the season dry (a), wet (b) and total P uptake/year (c) under various manurial practice




Fig. 3 Percentage distribution of organic and inorganic fractions in surface soil




Fig. 4 Composition percentage of various inorganic fractions of phosphorous to total inorganic P and total P in surface soil




Fig. 5 (a) Interaction between dry season yield and dry season P uptake




Fig. 5 (b) Interaction between wet season yield and wet season P uptake




Fig. 5 (c) Interaction between system yield and system P uptake




Fig. 6 (a) Percentage independent effect of fractions of P on dry season uptake




Fig. 6 (b) Percentage independent effect of fractions of P on system P uptake




Fig. 6 (c) Percentage independent effect of fractions of P on total uptake

Table 1. Long term effect of 13 years of cropping and manorial treatments on grain yield and P uptake.

Treatments	Grain yield (kg ha ⁻¹)		P uptake (kg ha ⁻¹)		Olsen's P (kg ha ⁻¹)
	Dry season 2017-18	Wet season 2018	Dry season 2017-18	Wet season 2018	Post-harvest surface soil
100% PK	2291 ^{de}	2873 ^e	7.66 ^{ef}	8.52 ^e	8.27 ^b
100% NPK	3063 ^{bc}	3456 ^{bc}	10.01 ^{cd}	11.97 ^{cd}	9.24 ^b
150% NPK	3469 ^b	3803 ^b	12.44 ^b	14.78 ^b	11.16 ^b
100% NPK + Zn	2625 ^{cde}	3353 ^{bcd}	8.66 ^{de}	11.14 ^b	9.85 ^b
100% NPK + FYM	4347 ^a	4750 ^a	16.62 ^a	19.79 ^a	49.72 ^a
100% NPK + Lime + FYM	4453 ^a	4891 ^a	16.89 ^a	20.55 ^a	53.41 ^a
100% NPK + B + Zn	2560 ^{de}	3556 ^{bc}	8.90 ^e	12.42 ^{cd}	9.10 ^b
100% NPK + S + Zn	2763 ^{cde}	3582 ^{bc}	10.20 ^{cd}	11.43 ^d	9.82 ^b
100% N	1751 ^{fg}	2970 ^{de}	5.08 ^{gh}	7.15 ^e	7.89 ^b
100 % NP	2186 ^{ef}	3132 ^{cde}	6.75 ^{fg}	8.83 ^e	8.84 ^b
100% NPK + Lime	3525 ^b	3807 ^b	11.61 ^{bc}	13.58 ^{bc}	10.59 ^b
Control	1325 ^g	1777 ^f	4.00 ^h	3.87 ^f	7.90 ^b
Initial					19.14

LSD (p<0.05%): in each column the values (mean of four replicates observations) followed by common letters are not significantly different (p, 0.05%) between treatments by DMRT

Table 2. Effect of long term manurial practice and nutrient content under long term manurial experimental treatments on different phosphorus fraction of surface soil (0-15 cm) in *Kharif* 2018

Treatments	Sal-P (kg ha ⁻¹)	Fe-P (kg ha ⁻¹)	Al- P (kg ha ⁻¹)	Red-P (kg ha ⁻¹)	Occl-P (kg ha ⁻¹)	Ca-P (kg ha ⁻¹)	Total Pi (kg ha ⁻¹)	Total Po (kg ha ⁻¹)	Total P (kg ha ⁻¹)
100% PK	43.34 ^{bc}	251.21 ^b	58.85 ^b	150.36 ^{abcd}	48.67 ^{def}	47.50 ^c	599.93 ^{bcd}	199.08 ^{ab}	799.02 ^a
100% NPK	40.85 ^{cd}	238.00 ^{bc}	56.17 ^b	151.93 ^a	46.91 ^{ef}	51.89 ^{bc}	585.74 ^{cde}	181.62 ^{ab}	767.37 ^{ab}
150% NPK	48.57 ^b	288.38 ^a	76.19 ^a	152.48 ^a	62.50 ^{bc}	56.84 ^b	684.96 ^a	156.96 ^{ab}	841.92 ^a
100%NPK + Zn	44.80 ^{bc}	270.16 ^a	54.56 ^b	150.83 ^a	60.96 ^{bc}	54.44 ^b	652.75 ^{ab}	152.35 ^{ab}	805.10 ^a
100% NPK + FYM	62.98 ^a	170.04 ^{cd}	55.93 ^b	108.71 ^{bc}	77.86 ^a	85.36 ^a	560.87 ^{cde}	204.29 ^a	765.17 ^{ab}
100% NPK+ Lime+ FYM	57.44 ^a	171.56 ^{cd}	61.75 ^a	104.69 ^c	68.75 ^{ab}	89.39 ^a	533.57 ^{de}	182.11 ^{ab}	735.69 ^{abc}
100% NPK+B+Zn	43.75 ^{bc}	234.79 ^{bc}	57.05 ^b	150.65 ^a	58.33 ^{cd}	54.63 ^b	612.40 ^{bc}	151.35 ^{ab}	763.75 ^{ab}
100%NPK+S+Zn	37.19 ^{de}	239.36 ^{bc}	48.97 ^c	145.19 ^a	50.72 ^{de}	47.79 ^c	585.21 ^{cde}	187.02 ^{ab}	772.24 ^{ab}
100% N	27.16 ^f	174.98 ^{cd}	39.53 ^d	103.66 ^c	36.64 ^g	36.58 ^d	418.55 ^f	48.58 ^b	467.13 ^c
100 %NP	32.23 ^{ef}	239.43 ^{bc}	45.16 ^c	129.49 ^{ab}	40.53 ^{fg}	40.70 ^d	527.53 ^e	200.08 ^{ab}	727.62 ^{abc}
100% NPK+lime	46.91 ^{bc}	193.61 ^{bcd}	46.54 ^c	137.11 ^a	48.83 ^{def}	75.28 ^a	548.28 ^{cde}	201.95 ^{ab}	750.19 ^{abc}
Control	20.35 ^g	152.89 ^d	35.07 ^d	79.36 ^d	25.19 ^h	24.87 ^e	337.73 ^g	186.45 ^{ab}	524.18 ^{bc}
Initial	22.35	172.0	50.62	85.50	28.31	24.87	383.65	248.35	632

LSD ($p < 0.05\%$): in each coloum the values (mean of four replicates observations) followed by common letters are not significantly different ($p, 0.05\%$) between treatments by DMRT

Table 3. Correlation of various P fractions of surface soil with crop yield and P uptake expressed in term of correlation coefficient (r)

P-fractions	Grain Yield		P uptake	
	Dry season 2017-18	Wet season 2018	Dry season 2017-18	Wet season 2018
Sal-P	0.580***	0.642***	0.581***	0.640***
Fe-P	0.042 ^{NS}	-0.058 ^{NS}	0.004 ^{NS}	-0.047 ^{NS}
Al-P	0.372**	0.413***	0.387**	0.457***
Red-P	0.079 ^{NS}	0.128 ^{NS}	0.079 ^{NS}	0.092 ^{NS}
Occl-P	0.659***	0.673***	0.697***	0.774***
Ca-P	0.0.819***	0.795***	0.810***	0.857***
Total P	0.394**	0.356*	0.375**	0.383**
Total Pi	0.395**	0.350*	0.373**	0.377**
Total Po	0.084 ^{NS}	0.083 ^{NS}	0.077 ^{NS}	0.093 ^{NS}

Note: p > 0.05 'NS' (non significant), p < (0.05) = '', p < (0.01) = '**' and p < 0.001 = '***'*

Table 4. Correlation of soil properties with various P fractions measured on postharvest surface soil of wet season 2018 expressed in term of correlation coefficient (r)

P-fractions	pH	SOC	CEC	Olsen's P
Sal-P	0.424**	0.594***	0.608***	0.469***
Fe-P	-0.290*	-0.155 ^{NS}	0.081 ^{NS}	-0.148 ^{NS}
Al-P	0.057 ^{NS}	0.374**	0.529***	0.318*
Red-P	-0.179 ^{NS}	-0.117 ^{NS}	0.154 ^{NS}	-0.273 ^{NS}
Occl-P	0.391**	0.485***	0.696***	0.639***
Ca-P	0.682***	0.604***	0.634***	0.739***
Total Pi	-0.070 ^{NS}	0.136 ^{NS}	0.472***	0.087 ^{NS}
Total Po	0.159 ^{NS}	0.266 ^{NS}	0.145 ^{NS}	-0.121 ^{NS}
Total P	0.087 ^{NS}	0.343*	0.485***	0.185 ^{NS}

Note: p > 0.05 'NS' (non-significant), p < (0.05) = '', p < (0.01) = '**' and p < 0.001 = '***'*

Table 5. Regression equation representing contribution of P fractions as independent variable on uptake of phosphorous as dependent variable

	Regression equation of independent fractions and uptake as dependent factor
Dry season uptake	$U = -1.332 + 0.077P_{Sal} - 0.056P_{Al} + 0.002P_{Fe} + 0.016P_{Red} + 0.026P_{Occl} + 0.123P_{Ca}$
Wet season uptake	$U = -1.762 + 0.081P_{Sal} - 0.060P_{Al} - 0.004P_{Fe} + 0.021P_{Red} + 0.601P_{Occl} + 0.139P_{Ca}$
Total uptake	$U = -3.097 + 0.157P_{Sal} - 0.116P_{Al} + 0.002P_{Fe} + 0.037P_{Red} + 0.087P_{Occl} + 0.262P_{Ca}$

Table 6. Relative contribution of inorganic P fractions to the total explained variation (%)

Variables	Relative Contribution of uptake on fractions		
	Dry season uptake	Wet season uptake	Total uptake
Sal-P	18.68	18.56	18.64
Fe-P	1.74	1.20	1.35
Al-P	6.90	7.63	7.28
Red-P	1.39	1.43	1.41
Occ-P	25.86	28.59	27.39
Ca-P	45.41	42.57	43.90
R ²	69.75	78.79	77.28