

# **A review on origin and genetic analysis of Wheat genes conferring resistance to wheat rust in common bread wheat (*Triticum aestivum* L.)**

## **Abstract**

Wheat rusts, produced by three species of *Puccinia*, are severe diseases and important biotic impediments in efforts to keep wheat production going around the world. The most commercially significant and prevalent diseases among global wheat varieties includes leaf rust also known as brown rust, caused by *Puccinia graminis triticea* Eriks, stem or black rust caused by *Puccinia graminis f. sp. tritici*, and stripe or yellow rust caused by *Puccinia striiformis f. Tritici*. Currently, more than 200 rust resistance genes (100 leaf rust, 60 stem rust, and 51 stripe rust) have been identified by using various molecular approaches from several wheat crop species and related wild species. This review highlights various aspects of *Lr* (Leaf rust), *Sr* (Stem rust) and *Yr* (yellow rust) resistance genes, their primitive origins, and their prevalence in globally wheat cultivars.

Kewwords: Wheat production, Rust, Resistant genes, Molecular approaches.

## **Introduction**

Wheat is one of the major cereal crops grown worldwide and one of the most important staple food crop for feeding 2.5 billion world population. Wheat cultivation represents approximately 19 % of global major cereal production (**FAOSTAT, 2020-21**). It is agronomically and nutritionally most important cereal essential for the food security, poverty alleviation and improved livelihoods. Wheat production is hampered by many biotic and abiotic stresses, resulting in yield disparity. Biotic stresses, including rust diseases, pose a serious danger to future food security since they reduce cereal crop yields drastically. Rust fungi are one of the most important diseases for cereals. Rust pathogens are obligatory parasitic organisms that have a macrocyclic and heteroecious life cycle (**Kolmer 2013**). Controlling of rust infections is very difficult due to the airborne nature of their spores, which can travel large distances and the fast emergence of highly aggressive races of rust pathogens. Rust infections have become more common in recent years, causing epidemics in Australia (**Murray et al., 2009**), in Europe (**Volkova et al., 2009**) and North and South American wheat-growing regions (**Germán et al.,**

2007). Ug99, a novel strain of stem rust discovered in Uganda in 1999, and its variants are pathogenic on around 80–90% of wheat cultivars and germplasm ( **Goyal and Manoharachary 2014**).

As a result, managing rust diseases is important for increasing world wheat production on a long-term basis. The most efficient and long-term strategies of managing diseases in crop plants are the development and deployment of resistant cultivars. **Biffen 1905**, made the first attempt to implement rust resistance breeding programmes in wheat. Modern breeding procedures, which are based on advanced genetic engineering and molecular biology tools, enable fresh and efficient techniques for developing new improved varieties.

Leaf rusts are the most severe rust disease in global wheat production, with yield losses of more than 50% reported in susceptible cultivars (**Draz et al., 2015**). Recently, many genes and QTLs governing rust resistance genes have been identified in wheat (**Buerstmayr et al., 2014**). A brief description about different identified rust resistant genes present below.

#### **Leaf rust resistance genes derived from various wheat species**

Over 100 leaf rust resistance genes have been isolated from common wheat, including *Lr 1* (**Roelfs et al., 2000**). *Lr2, Lr2a, Lr2b, Lr2c, Lr3, Lr3a, Lr3b, Lr3c* and *Lr4* to *Lr6* and also *Lr9* which was derived from *Ae.umbellulata*, *Lr14* gene that confers long term resistance in durams (**McIntosh et al., 2003**). *Lr20* (**Neu et al., 2002**), *Lr23* and *Lr30* (**Nelson et al., 1997**), *Lr49* and *Lr67* ( **shahin et al., 2015**), *Lr68* (**Herrera et al., 2012**), *trp1* and *trp2* recessive genes that conferred adult plant resistance in Brazilian wheat cultivar Toropi (**da SILVA et al., 2012**). Many resistance genes of major wheat diseases come from wild wheat cultivars like *Aegilops tauschii*, *Aegilops squarrosa.*, and *Triticum tauschii*. 3 leaf rust resistance genes *Lr19, Lr24* and *Lr29* was derived from *Aegilops elongatum*, 4 genes namely *Lr 28, Lr36, Lr47* and *Lr51* derived from *Aegilops speltoides* and one single gene from *Aegilops ventricosa* also isolated (**Todorovska et al., 2009**). *Lr41, Lr42,* and *Lr43* leaf rust resistance genes that transferred to common wheat from a wild species known as *Triticum tauschii* (**Cox et al., 1994**). 2 another leaf rust resistance genes *Lr14a* from durum wheat cultivars Hope and Gaza transferred into common wheat (**Aktar et al., 2017**).

#### **Identified stem rust resistance genes against Ug99 race**

In Uganda, a novel aggressive stem rust strain known as Ug99 was discovered in 1999 (Pretorius *et al.*, 2000). It is currently a major danger to global wheat production. Only 25% of India's total wheat-growing region is susceptible to stem rust (Nandni *et al.*, 2020). Approximately 60 genes for stem rust resistance (*Sr*) has been discovered so far (McIntosh *et al.*, 2016). *Sr13* was the first resistance gene identified in the germplasm of *Triticum dicoccom* but it also present in several *Triticum durum* species (Huerta *et al.*, 1992). Jyoti *et al.*, 2022 conducted genetic and phenotypic analysis on F2 and doubled haploid population from the cross between RL6071 (wheat line) and Tr129. Tr129 was also a wheat line which consist stem rust resistance gene derived from *Aegilops triuncialis*. Tr129 has 4 stem rust resistance (*Sr*) genes *Sr9b* , *Sr7b*, *Sr8a* and *SrTr129*, according to mapping and phenotyping. *Sr21* resistance gene derived from *Triticum monococcum* showed resistance against Ug99 race (Chen *et al.*, 2018). *Sr35* is a coiled-coil, nucleotide-binding, leucine-rich repeat gene from *Triticum monococcum* that exhibited resistance against Ug99 race (Saintenac *et al.*, 2013). *Sr2*, *Sr24*, *Sr25*, *Sr26*, *Sr31*, and *Sr38* resistant genes identified using linked molecular markers (Lin *et al.*, 2021). Some other resistance genes like *Sr24*, *Sr36*, *Sr7b* and *Sr9b* identified by using gene specific markers in Nebraska bread wheat germplasm conferred resistance against stem rust (Wu *et al.*, 2014). In recent years, new stem rust races like TKTTF, TTRTF have arisen, triggering a pandemic. As a result, stem rust has resurfaced as a danger to global wheat production (Olivera *et al.*, 2019).

### **Identified stripe rust resistance genes**

Stripe rust was common in the northwest United States and infection was mainly associated with cool weather (Johnson 1988). Approximately 51 genes for stripe rust resistance (*Yr*) have been discovered so far in which *Yr5* and *Yr15* are extremely resistant to all races found thus far in the United States (Mu *et al.*, 2020). *Yr10* was the first identified seedling resistance gene against stripe rust by using a map-based technique obtained from Moro wheat (Liu *et al.*, 2014). *Yr 15* resistant gene derived from wild emmer wheat (Yaniv *et al.*, 2015), *Yr51* and *Yr57* genes have been recently discovered to be effective in Australia against pre- and post-2002 *Pst* pathotypes (Randhawa *et al.*, 2019). *Yr81* from common wheat land race Aus27430 (Gessese *et al.*, 2019), *Yr82* from Aus27969 (Pakeerathan *et al.*, 2019), *Yr83* (Li *et al.*, 2020) also identified.

## **MAS and Modern approaches for Gene Pyramiding, mapping and gene Deployment for rust resistance genes**

Marker-assisted selection is a time-saving method of selecting desirable characters in an indirect manner. The primary prerequisite for marker assisted selection is the selection and detection of markers connected to the gene of interest. By the use of MAS it can be possible to release resistant variety quickly by doing gene deployment and gene pyramiding. Molecular markers are important in both genetic analysis of new sources of resistance genes and transferring two or more resistance genes in a single line (**Haile et al.,2013**). Molecular markers such as SSR( simple sequence repeat), AFLP (amplified fragment length polymorphism) and RAPD (random amplified polymorphic DNA) can be used to detect the resistance gene and compare rust populations. Some modern breeding tools include TILLING-targeting induced local lesions in genomes (**Comai et al., 2006**), transgenic technology (**Periyannan et al., 2013**), cisgenesis, reverse breeding, RNA-dependent DNA methylation (**Becker et al.,2012**), genome editing technology (GET), and genomic selection (**Desta et al.,2014**), have been added to molecular breeding techniques that already exist (**Savadi et al.,2018**). Recently, numerous genes and QTLs involving rust resistances have been identified in wheat (**Buerstmayr et al., 2014**).

STS marker was first used for detection of *Lr9* gene derived from *Aegilops umbellulata* (**Schachermayr et al., 1994**). STS, SCAR and CAPS markers was used for identification of *Lr19* (**Prabhu et al.,2004**), *Lr51* (**Helguera et al., 2005**) and *Lr52* (**Hiebert et al., 2005**), *Sr24* and *Sr26* (**Mago et al.,2005**), *Lr1* (**Cloutier et al., 2007**) and *Lr10* (**Feuillet et al., 2003**) have been identified by cloning and gene sequencing. *Lr36*, *Lr37* and *Lr38* by PCR assay (**Helguera et al.,2003**). Some stem rust resistance genes like *Sr39* resistant gene was identified by using SCAR marker in Canadian wheat (**Zhang et al., 2012**). *Sr9a* stem rust resistant gene was also identified by Microsatellite markers (**Tsilo et al., 2007**). *Sr6*, *Sr40* by Microsatellite marker (**Ejaz et al., 2012**). Simple sequence repeat SSR markers used to identify and map the APR (adult plant resistance) stem rust resistant gene *Sr56* in a winter wheat variety (**Bansal et al., 2014**). *Sr58*, *Lr46* and *Yr29* by CAPS molecular marker (**Yu et al., 2014**). Other relevant markers, EST-STS diagnostic markers, which were used for the resistance gene YrSM139-1B derived from *Triticum dicoccoides* against stripe rust in bread wheat (**Zhang et al., 2016**). Therefore, these technologies enable us to modify crop plants more precisely than ever before,

accelerating crop enhancement efforts for long-term food production while also approaching safety concerns about food crops.

### **Conclusion**

Knowledge about origin, distribution and deployment of resistance genes is essential for developing new wheat varieties with desirable characters. Gene pyramiding by MAS and the application of other molecular methods are essential for maintaining the long-term or durable resistance against rust disease. Overall, the goal of this review paper is to improve understanding of the present status of wheat research among wheat researchers, students, academics, plant breeders, and pathologists.

### **References**

- Aktar-Uz-Zaman, M. D., Tuhina-Khatun, M. S. T., Hanafi, M. M., & Sahebi, M. (2017). Genetic analysis of rust resistance genes in global wheat cultivars: an overview. *Biotechnology & Biotechnological Equipment*, 31(3), 431-445.
- Bansal, U., Bariana, H., Wong, D., Randhawa, M., Wicker, T., Hayden, M., & Keller, B. (2014). Molecular mapping of an adult plant stem rust resistance gene Sr56 in winter wheat cultivar Arina. *Theoretical and applied genetics*, 127(6), 1441-1448.
- Becker, C., & Weigel, D. (2012). Epigenetic variation: origin and transgenerational inheritance. *Current opinion in plant biology*, 15(5), 562-567.
- Biffen, R. H. (1905). Mendel's laws of inheritance and wheat breeding. *The Journal of Agricultural Science*, 1(1), 4-48.
- Buerstmayr, M., Matiasch, L., Mascher, F., Vida, G., Ittu, M., Robert, O., ... & Buerstmayr, H. (2014). Mapping of quantitative adult plant field resistance to leaf rust and stripe rust in two European winter wheat populations reveals co-location of three QTL conferring resistance to both rust pathogens. *Theoretical and Applied Genetics*, 127(9), 2011-2028.
- Buerstmayr, M., Matiasch, L., Mascher, F., Vida, G., Ittu, M., Robert, O., ... & Buerstmayr, H. (2014). Mapping of quantitative adult plant field resistance to leaf rust and stripe rust in two European winter wheat populations reveals co-location of three QTL conferring resistance to both rust pathogens. *Theoretical and Applied Genetics*, 127(9), 2011-2028.

- Chen, S., Zhang, W., Bolus, S., Rouse, M. N., & Dubcovsky, J. (2018). Identification and characterization of wheat stem rust resistance gene Sr21 effective against the Ug99 race group at high temperature. *PLoS genetics*, *14*(4), e1007287.
- Cloutier, S., McCallum, B. D., Loutre, C., Banks, T. W., Wicker, T., Feuillet, C., ... & Jordan, M. C. (2007). Leaf rust resistance gene Lr1, isolated from bread wheat (*Triticum aestivum* L.) is a member of the large psr567 gene family. *Plant molecular biology*, *65*(1), 93-106.
- Comai, L., & Henikoff, S. (2006). TILLING: practical single-nucleotide mutation discovery. *The Plant Journal*, *45*(4), 684-694.
- Cox, T. S., Raupp, W. J., & Gill, B. S. (1994). Leaf rust-resistance genes Lr41, Lr42, and Lr43 transferred from *Triticum tauschii* to common wheat. *Crop science*, *34*(2), 339-343.
- da SILVA, P. R., Brammer, S. P., Guerra, D., Milach, S. C. K., Barcellos, A. L., & Baggio, M. I. (2012). Monosomic and molecular mapping of adult plant leaf rust resistance genes in the Brazilian wheat cultivar Toropi. *Embrapa Trigo-Artigo em periódico indexado (ALICE)*.
- Desta, Z. A., & Ortiz, R. (2014). Genomic selection: genome-wide prediction in plant improvement. *Trends in plant science*, *19*(9), 592-601.
- Draz, I. S., Abou-Elseoud, M. S., Kamara, A. E. M., Alaa-Eldein, O. A. E., & El-Bebany, A. F. (2015). Screening of wheat genotypes for leaf rust resistance along with grain yield. *Annals of Agricultural sciences*, *60*(1), 29-39.
- Ejaz, M., Iqbal, M., Shahzad, A., Ahmed, I., & Ali, G. M. (2012). Genetic variation for markers linked to stem rust resistance genes in Pakistani wheat varieties. *Crop Science*, *52*(6), 2638-2648.
- Food and Agriculture Organization Corporate Statistical Database (FAOSTAT 2020-21) [Internet] <https://www.fao.org/faostat/en/#home>
- Feuillet, C., Travella, S., Stein, N., Albar, L., Nublát, A., & Keller, B. (2003). Map-based isolation of the leaf rust disease resistance gene Lr10 from the hexaploid wheat (*Triticum aestivum* L.) genome. *Proceedings of the National Academy of Sciences*, *100*(25), 15253-15258.
- Germán, S., Barcellos, A., Chaves, M., Kohli, M., Campos, P., & de Viedma, L. (2007). The situation of common wheat rusts in the Southern Cone of America and perspectives for control. *Australian Journal of Agricultural Research*, *58*(6), 620-630.

- Gessese, M., Bariana, H., Wong, D., Hayden, M., & Bansal, U. (2019). Molecular mapping of stripe rust resistance gene Yr81 in a common wheat landrace Aus27430. *Plant disease*, 103(6), 1166-1171.
- Goyal, A., & Manoharachary, C. (Eds.). (2014). *Future challenges in crop protection against fungal pathogens*. Springer New York.
- Haile, J. K., & Rouml, M. S. (2013). Status of genetic research for resistance to Ug99 race of Puccinia graminis f. sp. tritici: A review of current research and implications. *African Journal of Agricultural Research*, 8(50), 6670-6680.
- Helguera, M., Khan, I. A., Kolmer, J., Lijavetzky, D., Zhong-Qi, L., & Dubcovsky, J. (2003). PCR assays for the Lr37-Yr17-Sr38 cluster of rust resistance genes and their use to develop isogenic hard red spring wheat lines. *Crop science*, 43(5), 1839-1847.
- Helguera, M., Vanzetti, L., Soria, M., Khan, I. A., Kolmer, J., & Dubcovsky, J. (2005). PCR markers for Triticum speltoides leaf rust resistance gene Lr51 and their use to develop isogenic hard red spring wheat lines. *Crop science*, 45(2), 728-734.
- Herrera-Foessel, S. A., Singh, R. P., Huerta-Espino, J., Rosewarne, G. M., Periyannan, S. K., Viccars, L., ... & Lagudah, E. S. (2012). Lr68: a new gene conferring slow rusting resistance to leaf rust in wheat. *Theoretical and Applied Genetics*, 124(8), 1475-1486.
- Hiebert, C., Thomas, J., & McCallum, B. (2005). Locating the broad-spectrum wheat leaf rust resistance gene Lr52 (LrW) to chromosome 5B by a new cytogenetic method. *Theoretical and Applied Genetics*, 110(8), 1453-1457.
- Huerta-Espino, J., & Roelfs, A. P. (1992). Leaf rust on durum wheats. *Vortraege fuer Pflanzenzuechtung (Germany)*.
- Johnson, R. (1988). Breeding strategies for resistance to the rusts of wheat. *DF CIMMYT: Mexico City, Mexico*, 63-75.
- Kolmer, J. (2013). Leaf rust of wheat: pathogen biology, variation and host resistance. *Forests*, 4(1), 70-84.
- Li, J., Dundas, I., Dong, C., Li, G., Trethowan, R., Yang, Z., ... & Zhang, P. (2020). Identification and characterization of a new stripe rust resistance gene Yr83 on rye chromosome 6R in wheat. *Theoretical and Applied Genetics*, 133(4), 1095-1107.
- Liu, W., Frick, M., Huel, R., Nykiforuk, C. L., Wang, X., Gaudet, D. A., ... & Laroche, A. (2014). The stripe rust resistance gene Yr10 encodes an evolutionary-conserved and unique CC-NBS-LRR sequence in wheat. *Molecular plant*, 7(12), 1740-1755.

- Mago, R., Bariana, H. S., Dundas, I. S., Spielmeier, W., Lawrence, G. J., Pryor, A. J., & Ellis, J. G. (2005). Development of PCR markers for the selection of wheat stem rust resistance genes Sr24 and Sr26 in diverse wheat germplasm. *Theoretical and Applied Genetics*, 111(3), 496-504.
- McIntosh, R. A., Dubcovsky, J., Rogers, W. J., Morris, C., Appels, R., & Xia, X. C. (2016). Catalogue of gene symbols for wheat: 2015–2016 supplement. *Komugi Wheat Genet. Resour. Database*.
- McIntosh, R. A., Yamazaki, Y., Dubcovsky, J., Rogers, J., Morris, C., & Appels, R. (2013, September). Catalogue of gene symbols for wheat. In *Proceedings of the 12th International Wheat Genetics Symposium* (pp. 8-13). Yokohama Japan.
- Mu, J., Liu, L., Liu, Y., Wang, M., See, D. R., Han, D., & Chen, X. (2020). Genome-wide association study and gene specific markers identified 51 genes or QTL for resistance to stripe rust in US winter wheat cultivars and breeding lines. *Frontiers in plant science*, 998.
- Murray, G. M., & Brennan, J. P. (2009). Estimating disease losses to the Australian wheat industry. *Australasian Plant Pathology*, 38(6), 558-570.
- Nandni, S., Dev, D., & Singh, K. P. (2020). Ug99: Wheat Stem Rust Race: Exploring the World. In *Diseases of Field Crops: Diagnosis and Management* (pp. 81-88). Apple Academic Press.
- Nelson, J. C., Singh, R. P., Autrique, J. E., & Sorrells, M. E. (1997). Mapping genes conferring and suppressing leaf rust resistance in wheat. *Crop science*, 37(6), 1928-1935.
- Neu, C., Stein, N., & Keller, B. (2002). Genetic mapping of the Lr20 Pm1 resistance locus reveals suppressed recombination on chromosome arm 7AL in hexaploid wheat. *Genome*, 45(4), 737-744.
- Olivera, P. D., Sikharulidze, Z., Dumbadze, R., Szabo, L. J., Newcomb, M., Natsarishvili, K., ... & Jin, Y. (2019). Presence of a sexual population of *Puccinia graminis* f. sp. *tritici* in Georgia provides a hotspot for genotypic and phenotypic diversity. *Phytopathology*, 109(12), 2152-2160.
- Pakeerathan, K., Bariana, H., Qureshi, N., Wong, D., Hayden, M., & Bansal, U. (2019). Identification of a new source of stripe rust resistance Yr82 in wheat. *Theoretical and Applied Genetics*, 132(11), 3169-3176.
- Periyannan, S., Moore, J., Ayliffe, M., Bansal, U., Wang, X., Huang, L., ... & Lagudah, E. (2013). The gene Sr33, an ortholog of barley Mla genes, encodes resistance to wheat stem rust race Ug99. *Science*, 341(6147), 786-788.

- Prabhu, K. V., Gupta, S. K., Charpe, A., & Koul, S. (2004). SCAR marker tagged to the alien leaf rust resistance gene Lr19 uniquely marking the *Agropyron elongatum*-derived gene Lr24 in wheat: a revision. *Plant Breeding*, 123(5), 417-420.
- Pretorius, Z. A., Singh, R. P., Wagoire, W. W., & Payne, T. S. (2000). Detection of virulence to wheat stem rust resistance gene Sr31 in *Puccinia graminis* f. sp. *tritici* in Uganda. *Plant Disease*, 84(2), 203-203.
- Randhawa, M. S., Bains, N. S., Sohu, V. S., Chhuneja, P., Trethowan, R. M., Bariana, H. S., & Bansal, U. (2019). Marker assisted transfer of stripe rust and stem rust resistance genes into four wheat cultivars. *Agronomy*, 9(9), 497.
- Randhawa, M. S., Bariana, H. S., Mago, R., & Bansal, U. K. (2015). Mapping of a new stripe rust resistance locus Yr57 on chromosome 3BS of wheat. *Molecular breeding*, 35(2), 1-8.
- Roelfs, A. P., Hughes, M. E., & Long, D. L. (2000). Rust resistance genes in wheat lines and cultivars. *US Dep. Agric. Res. Serv. Cereal Dis. Lab. Online, Publication CDL-EP*, 6.
- Saintenac, C., Zhang, W., Salcedo, A., Rouse, M. N., Trick, H. N., Akhunov, E., & Dubcovsky, J. (2013). Identification of wheat gene Sr35 that confers resistance to Ug99 stem rust race group. *Science*, 341(6147), 783-786.
- Savadi, S., Prasad, P., Kashyap, P. L., & Bhardwaj, S. C. (2018). Molecular breeding technologies and strategies for rust resistance in wheat (*Triticum aestivum*) for sustained food security. *Plant pathology*, 67(4), 771-791.
- Schachermayr, G., Siedler, H., Gale, M. D., Winzeler, H., Winzeler, M., & Keller, B. (1994). Identification and localization of molecular markers linked to the Lr9 leaf rust resistance gene of wheat. *Theoretical and applied genetics*, 88(1), 110-115.
- Shahin, S. I., & El-Orabey, W. M. (2015). Relationship between partial resistance and inheritance of adult plant resistance gene Lr 46 of leaf rust in six bread wheat varieties. *International Journal of Science and Research*, 4(1), 1511-1517.
- Todorovska, E., Christov, N., Slavov, S., Christova, P., & Vassilev, D. (2009). Biotic stress resistance in wheat—breeding and genomic selection implications. *Biotechnology & Biotechnological Equipment*, 23(4), 1417-1426.
- Tsilo, T. J., Jin, Y., & Anderson, J. A. (2007). Microsatellite markers linked to stem rust resistance allele Sr9a in wheat.
- Volkova, G. V., Alekseeva, T. P., Anpilogova, L. K., Dobryanskaya, M. V., Vaganova, O. F., & Kol'bin, D. A. (2009). Phytopathological characteristics of leaf rust resistance of new winter wheat varieties. *Russian Agricultural Sciences*, 35(3), 168-171.

- Wu, X. X., Li, T. Y., Chen, S., Wang, G. Q., Cao, Y. Y., Ma, S. L., & Li, M. J. (2014). Stem rust resistance evaluation and Ug99-resistance gene detection of 139 wheat cultivars. *Scientia Agricultura Sinica*, 47(23), 4618-4626.
- Yaniv, E., Raats, D., Ronin, Y., Korol, A. B., Grama, A., Bariana, H., ... & Fahima, T. (2015). Evaluation of marker-assisted selection for the stripe rust resistance gene Yr15, introgressed from wild emmer wheat. *Molecular breeding*, 35(1), 1-12.
- Yu, L. X., Barbier, H., Rouse, M. N., Singh, S., Singh, R. P., Bhavani, S., ... & Sorrells, M. E. (2014). A consensus map for Ug99 stem rust resistance loci in wheat. *Theoretical and Applied Genetics*, 127(7), 1561-1581.
- Yu, L. X., Barbier, H., Rouse, M. N., Singh, S., Singh, R. P., Bhavani, S., ... & Sorrells, M. E. (2014). A consensus map for Ug99 stem rust resistance loci in wheat. *Theoretical and Applied Genetics*, 127(7), 1561-1581.
- Zhang, Q., Klindworth, D. L., Friesen, T. L., Chao, S., Jin, Y., Cai, X., & Xu, S. S. (2012, January). Development and characterization of wheat lines with Sr37 for stem rust resistance derived from wild Timopheev's wheat. In *Meeting Abstract* (Vol. 316, No. 6).
- Zhang, T., Jin, Y., Zhao, J. H., Gao, F., Zhou, B. J., Fang, Y. Y., & Guo, H. S. (2016). Host-induced gene silencing of the target gene in fungal cells confers effective resistance to the cotton wilt disease pathogen *Verticillium dahliae*. *Molecular Plant*, 9(6), 939-942.