

Effect of Mulching and Sulphur on Growth and Yield of Yellow mustard (*Sinapis alba*)

ABSTRACT

A field experiment was carried out at Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, SHUATS, Prayagraj (U.P) in *Rabi*, 2021 to study the Effect of Biofertilizers and Sulphur on Growth and Yield of Yellow Mustard (*Sinapis alba*). It was consisting of combination of three types of Mulching No mulch, Paddy straw and Poly sheet mulch and three Sulphur levels (20kg S/ha, 40 kg S/ha and 60 kg S/ha). The experiment was laid out in Randomized Block Design with nine treatments each replicated thrice. The experiment results revealed that the growth parameters and yield parameters such as plant height (145.94 cm), dry weight (34.30 g), number of branches/plant (11.58), number of seeds/siliquae (37.41) and test weight (3.08 g) at harvest, significantly recorded in treatment T₉ with the application of Poly sheet mulch+60 kg S/ha. Moreover, seed yield (1.82 t/ha), stover yield (3.94 t/ha), gross return (95,762.64 INR/ha), net return (52,487.44 INR/ha) and B:C ratio (2.21) were also recorded significantly higher in the treatment which is Poly sheet mulch + Sulphur 60 kg/ha among all treatments.

Keywords: *Mulching, Sulphur, yellow mustard, Growth and Yield.*

INTRODUCTION

In present era oil seed crops form the main source of energy for major portion of Indian population due to less availability of animal fat and its ill effect on heart. Among the rapeseed and mustard group, yellow sarson which belongs to family Brassicaceae is an important crop in terms of its high seed oil and protein content. value. In India, about 27.5 million ha area is occupied by oilseeds which represent 14 per cent of the total cropped area with the production of 24.72 million tonnes accounting for 5 per cent of gross national product and 10 per cent value of all agricultural commodities (Anonymous, 2015).

India mustard is predominantly cultivated in the states of Rajasthan, Uttar Pradesh, Haryana, Madhya Pradesh and Gujarat. Rajasthan ranks first in area and production of rapeseed and mustard with 2.50 million ha area and 3.71 million tonnes production. Mustard oil is used as condiment in pickles, flavoring curries and vegetables, preparation of hair oils, medicines, soap making and in the tanning industry for softening of leather. The mustard cake is used mostly for cattle feed and manure (Potdar *et al.*, 2019).

One of the common practices to reduce evaporation loss from the soil and prolonging the availability of moisture to the crop is the use of mulches. Mulching increases soil moisture, regulates soil temperature, suppresses weed growth, minimized leaching loss of nutrients, checks excessive evaporation, reduces soil erosion, improves production and quality. Thus, mulching economises the use of irrigation or rainwater and boosts nutrient use efficiency by conserving more water in the crop rhizosphere. The effects of mulch on soil temperature, moisture regime and root growth as well as yield depend on the micro-environment, made of mulch application and quality and quantity of mulch materials (Manoj *et al.*, 2014). Mulch increased soil organic matter and soil moisture contents but decreased bulk density and soil strength compared to control. The effects of mulch on soil temperature, moisture regime and root growth as well as yield depend on the micro-environment, made of mulch application and quality and quantity of mulch materials. Keeping in view these circumstances, it was contemplated to work out the suitable tillage and mulching practices for taking the higher yield of mustard in rainfed condition.

The poly sheet can be used in moisture deficit areas. The polythene mulch is easily available in the market and its use in agriculture is eco-friendly as after use, poly sheets may be collected from the crop field and can be well mixed with coltar during construction of pucca roads. If black polyethylene is used for the purpose, it prevents entry of light to the soil surface thus restricting any possible growth of weed, which is a menace in rainfed agriculture.

Sulphur is the nutrient which plays a multitudinous role in providing nutrition to oilseed crops, particularly those belonging to cruciferae family. The importance of sulphur is obvious in oilseed production as it is required for the synthesis of sulphur containing amino acids methionine (21%), cysteine (26%) and cystine (27%), which are essential components of protein and oil as well as for vegetative growth of the plant, it is involved in the formation of chlorophyll, glucosides and glucosinolates (mustard oils), activation of enzymes and sulphhydryl (-SH) linkages that are the source of pungency in oils and increases the root growth and stimulate seed formation. Sulphur is a secondary plant nutrient which plays a significant role in increasing production specially in oil seed crops. Rapeseed-mustard crops are particularly sensitive to sulphur deficiency mainly due to the fact that S plays an important role in the chemical composition of seed and increases the percentage of oil content of seed. Sulphur increases the oil content and gives pungency to oil as it forms certain disulphide linkages. (Khan *et al.*, 2002). Oilseed crops require more Sulphur than cereals as their oil storage organs are mostly proteins, rich in S. Deficiency of Sulphur is known to hamper N metabolism in plants as well as synthesis of S-containing amino acids and thus exerts adverse effects on both seed and oil yield.

MATERIALS AND METHODS

The present examination was carried out during *Rabi* 2021-2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, UP, which is located at 25.28°N latitude, 81.54°E longitude and 98 m altitude above the mean sea level. The experiment laid out in Randomized Block Design which consisting of nine treatments with No Mulch + Sulphur 20 kg/ha, Paddy straw mulch + Sulphur 40 kg/ha, Poly sheet mulch + Sulphur 60 kg/ha, No Mulch + Sulphur 20 kg/ha, Paddy straw mulch + Sulphur 40 kg/ha, Poly sheet mulch + Sulphur 60 kg/ha, No mulch + Sulphur 20 kg/ha, Paddy straw mulch + Sulphur 40 kg/ha, Poly sheet mulch + Sulphur 60 kg/ha. The observations recorded on different growth parameters at harvest viz, plant height (cm), number of branches per plant, Plant dry weight, number of siliques per plant, number of grains per silique, test weight, grain yield and stover yield and harvest index were recorded and statistically analyzed using analysis of variance (ANOVA) as applicable to Randomized Block Design (Gomez and Gomez 1984).

RESULT AND DISCUSSION

Growth Attributes

Effect on the growth of yellow mustard.

As can be seen in Table.1, growth parameters are summarized statistically. At 100 DAS, significantly taller plant height (145.94 cm) was recorded with Poly sheet mulch + Sulphur 60 kg /ha. However, *Paddy straw mulch + Sulphur 60 kg/ ha, Poly sheet mulch + Sulphur 40 kg/ ha* was statistically at par with *Paddy straw mulch + Sulphur 60 kg/ ha*. The minimum plant height was recorded in the treatment combination of No mulch + Sulphur 20 kg/ha which is 128.52 cm. Significantly maximum Number of branches (11.69) was recorded with application of *Paddy straw mulch + Sulphur 60 kg/ ha*. However, *Poly sheet mulch + Sulphur 60 kg/ ha* was statistically at par with *Poly sheet mulch + Sulphur 60 kg/ ha*. The minimum number of branches was recorded in the treatment combination of No mulch + Sulphur 20 kg/ha which is 9.01 g. Significantly maximum plant dry weight (34.30 g) was recorded in the treatment combination *Poly sheet mulch + Sulphur 60 kg/ ha*. However, *Paddy straw mulch + Sulphur 60 kg/ ha, Poly sheet mulch + Sulphur 40 kg/ ha* was statistically at par with *Paddy straw mulch + Sulphur 60 kg/ ha*. Pachauri *et al.*, (2012) reported that various levels of sulphur significantly influenced the growth parameters viz., plant height and dry weight of plant. The plant height increased significantly with each increment in the dose up to 60 kg /ha. However, the difference in plant height due to further increase in the dose of sulphur was not significant. Application of 60 kg S /ha produced more dry weight of plant at 90 DAS as compared to control and 30 kg S/ ha. Better nutrition to plant resulted in more height and number of branches and other growth parameters, which resulted in higher dry weight of plant. These results are in conformity with those reported by Kumar and Yadav (2007) in mustard. The overall improvement in growth character of plant owing of sulphur application were known sulphur enhance cell multiplication, elongation and expansion, imparts a deep colour to leaves due to better chlorophyll synthesis resulting in greater amount of dry matter in comparison to sulphur deficient plant. This argument was also supported by Singh and Meena (2004) [14], Mishra (2001) [8], Nepalia and Jain (2000) [9]

Yield Attributes:

As can be seen in Table.2, yield parameters are summarized statistically. At the time of harvest, significantly length of silique (8.02 cm) was recorded with Paddy straw mulch + Sulphur 60 kg /ha. However, *Poly sheet mulch + Sulphur 40 kg/ ha*, *Poly sheet mulch + Sulphur 60 kg/ ha* was statistically at par with *Paddy straw mulch + Sulphur 60 kg/ ha*. The minimum length of silique was recorded in the treatment combination of No mulch + Sulphur 20 kg/ha which is (6.10). Significantly maximum Number of seeds per silique (37.41) was recorded with application of *Poly sheet mulch + Sulphur 60 kg/ ha*. However, *Paddy straw mulch + Sulphur 60 kg/ ha* was statistically at par with *Poly sheet mulch + Sulphur 60 kg/ ha*. The minimum number of seeds per silique was recorded in the treatment combination of No mulch + Sulphur 20 kg/ha which is (27.07). Significantly maximum test weight (3.08 g) was recorded in the treatment combination *Poly sheet mulch + Sulphur 60 kg/ ha*. However, *Paddy straw mulch + Sulphur 60 kg/ ha*, *Poly sheet mulch + Sulphur 40 kg/ ha* was statistically at par with *Paddy straw mulch + Sulphur 60 kg/ ha*. The minimum test weight was recorded in the treatment combination of No mulch + Sulphur 20 kg/ha which is (2.13 g). Significantly maximum grain yield (1.82 t) was recorded in the treatment combination *Poly sheet mulch + Sulphur 60 kg/ ha*. However, *Paddy straw mulch + Sulphur 60 kg/ ha*, *Poly sheet mulch + Sulphur 40 kg/ ha* was statistically at par with *Paddy straw mulch + Sulphur 60 kg/ ha*. The minimum grain yield was recorded in the treatment combination of No mulch + Sulphur 20 kg/ha which is (1.13 t). Significantly maximum stover yield (3.94 t) was recorded in the treatment combination *Poly sheet mulch + Sulphur 60 kg/ ha*. However, *Paddy straw mulch + Sulphur 60 kg/ ha*, *Poly sheet mulch + Sulphur 40 kg/ ha* was statistically at par with *Paddy straw mulch + Sulphur 60 kg/ ha*. The minimum stover yield was recorded in the treatment combination of No mulch + Sulphur 20 kg/ha which is (1.71 t). Significantly maximum harvest index % (37.68) was recorded in the treatment combination *Poly sheet mulch + Sulphur 60 kg/ ha*. However, *Paddy straw mulch + Sulphur 60 kg/ ha*, *Poly sheet mulch + Sulphur 40 kg/ ha* was statistically at par with *Paddy straw mulch + Sulphur 60 kg/ ha*. The minimum harvest index was recorded in the treatment combination of No mulch + Sulphur 20 kg/ha which is (32.22 %). Makeen et al., (2008) reported that number of siliqua plant, number of grains siliqua weight of 1000 seeds, seed yield and harvest index were significantly influenced by application of sulphur. Application of sulphur @ 60 kg/ha recorded the highest values with respect to these parameters. The seed yield increased to 25.5 q/ha at 60 kg S ha⁻¹ as compared to 11.1 q/ha at control. These results are in agreement with the findings of Raut et al., (2000), Sharma (1994) and Sarma and Dehnath (1999). Harvest index increased significantly and harvest index at 60 kg S/ ha was higher than other treatments.

Table 1. Effect of Mulching and Sulphur on growth attributes of Yellow Mustard

Treatment combination	At 100DAS		
	Plant Height (cm)	Number of Branches per plant	Plant dry weight (g/plant)
1-No mulch + sulphur 20 kg / ha	128.52	9.01	22.74
2-Paddy straw mulch+ sulphur 20kg ha	135.23	9.21	26.25
3-Polysheet mulch + sulphur 20 kg / ha	132.79	9.17	24.39
4- No mulch + sulphur 40 kg / ha	138.04	9.70	28.43
5-Paddy straw mulch +sulphur 40 kg/ ha	143.18	10.75	32.11
6-Poly sheet mulch + sulphur 40 kg/ ha	143.62	10.93	30.18
7- No mulch + sulphur 60 kg/ ha	141.50	9.88	31.56
8 Paddy straw mulch + sulphur 60kg/ha	144.38	11.69	33.12
9-Polysheet mulch + sulphur 60 kg/ ha	145.94	11.58	34.30
F-Test	S	S	S
Sem (±)	2.29	0.63	1.13
CD (5%)	6.88	1.89	3.40

Table 2. Effect of Mulching and sulphur on yield attributes of Yellow Mustard

Treatment combination	At Harvest					
	Length	No. of	Test	Grain	Stover	Harvest

	of silique	Seeds/silique	weight(g)	yield (t/ha)	yield (t/ha)	Index (%)
1	6.10	27.07	2.13	1.13	2.71	32.22
2	7.46	29.19	2.31	1.32	2.95	41.19
3	7.56	28.45	2.26	1.21	2.86	33.15
4	6.51	30.14	2.49	1.40	3.17	35.99
5	7.63	34.22	2.84	1.68	3.72	40.78
6	7.78	33.73	2.72	1.54	3.61	35.48
7	6.98	32.68	2.56	1.49	3.32	33.41
8	8.02	35.16	2.97	1.71	3.83	35.85
9	7.78	37.41	3.08	1.82	3.94	37.68
F-Test	S	S	NS	S	S	S
Sem (±)	0.35	1.32	0.27	0.06	0.04	1.94
CD (5%)	1.04	3.95	0.80	0.17	0.12	5.80

CONCLUSION

It is concluded that, treatment 9 with Polysheet mulch and sulphur 60 kg/ha had performed better in growth and yield parameters. As it was more productive it can be recommended to farmers after further trials.

REFERENCES

- A.O.A.C., Official methods of analysis, Association of official Analytical Chemists. 15th ed. Washington, D.C, USA, 1995.
- Bouyoucos, G.J. (1927). The Hydrometer as a new method of the mechanical Analysis of soils. *Soil science*, **23**: 343-353
- Chauhan, D.R., Ram, M. and Singh, I. 2002. Response of Indian mustard (*Brassica juncea*) to irrigation and fertilization with various sources and levels of Sulphur. *Indian Journal of Agronomy*, **47**(3):422-426.
- Das, A. and Biswas, P.K. (2019). Effect of sulphur and bio fertilizers on growth attributes of sesame (*Sesamum indicum* L.) and soil fertility in red and lateritic soils of West Bengal, India. *Indian Journal of Hill Farming*. **32**(2): 287-294.
- Fisher, R.A. and Yates, F. (1963). Statistical tables for biological, agricultural and medical research. **22**(2): 151-158.
- Jackson, M.L. (1967). Soil chemical Analysis. *Open Journal of Soil Science*. **5**(4): 498
- Jaiswal A.D., Singh S.K., Singh Y.K., Surendra Singh and Yadav S.N. (2014). Effect of sulphur and boron on yield and quality of mustard (*Brassica juncea* L.) grown on vindhyan red soil. *Journal of Indian Society of Soil Science* **63**(3): 362-364.
- Kapur L.T., Patel A.R. And Thakor R.F. (2010). Yield attributes and yield of mustard (*Brassica juncea* L. Czern and Coss) as affected by sulphur levels. *An Asian Journal of Soil Science* **5**(1): 216-217.

- Lakshman K., Vyas A. K., Shiva Kumar B. G. and Rana D.S. (2015). Effect of levels and time of Sulphur application on growth, yield and quality of soybean (*Glycine max*). *Indian Journal of Agronomy* **60**(1): 121-125.
- Mallick, R.B. and Raj, A. (2015). Influence of phosphorus, sulphur and boron on growth, yield,
- Manoj, K. and Premi, O.P. (2003). Effect of Poly Mulch on Soil Moisture and Yield of Rainfed Mustard. *Agric. Sci. Digest*. **23**(3): 183-186.
- Neha, Dashora, L. N., Kaushik, M. K., Upodhayay, B. (2014). Yield, nutrient content uptake and quality of Indian mustard as influenced by sulphur under southern Rajasthan conditions *Annals of Agro-Bio Research* **19**(1):81-84.
- Om Prakash and Singh, B.P. 2002. Effect of Sulphur fertilization on growth, yield and quality of Indian mustard (*Brassica juncea* L.) genotypes. *Annals of Agricultural Research*, **23**(2): 275-279.
- Parihar, S., Kameariya, P.R. and Choudary, R. (2014). Response of mustard (*Brassica juncea*) to varying levels of Sulphur and fortified vermicompost under loamy sand soil. *Agriculture Science Digest*. **34**(4):296-298.
- Rana, K.S. and Rana, D.S. (2003). Response of mustard (*Brassica juncea*) to nitrogen and Sulphur under dryland conditions. *Indian Journal of Agronomy*, **48**(3): 217-219.
- Sah, D., Sewak, R., Singh, A.K. and Swami, S. (2013). Growth, yield and profitability of Indian mustard (*Brassica juncea* (L.) Czern & Coss) with different weed control measures and sulphur levels. *Agricultural Science Digest* **33**(1): 15-20.
- Tiwari, R.C., Kumar, S. and Singh, D.P. (2003). Sulphur status of soil and crops *Fertilizer News*, **48**(8):35-38.
- Upadhyay P.K., Singh D.P., Singh M.P. and Ashutosh Srivastava (2018). Effect of phosphorus and sulphur level on plant growth and dry matter production of mustard (*Brassica juncea* L.). *International Journal of Pure and Applied Bioscience* **6**(6): 751-757.
- Virendra, S.A., Sangha, M.K. (2008). Effect of foliar application of Sulphur on yield quality and economics of Indian Mustard (*Brassica Juncea* (L.) Czer & Coss". *Research on Crops*, **9**(3):728-730.
- William, M.J. (1975). USDA triangle, A Basic System of Soil Classification for Making and Interpreting Soil Surveys. United States Department of Agriculture, *Agriculture Handbook*, 470-472.
- Yadav, S.S. Singh, B. Singh, S. and Tikoo, A. (2005). Effect of levels and sources of Phosphorus and Sulphur on growth and yield of Indian mustard (*Brassica juncea* L.) in light textured soils of Southern Haryana. *Haryana Journal of Agronomy*, **21** (2):136-137.

UNDER PEER REVIEW




Image 1 : Arrangement of Poly sheet mulch in experimental field at Crop Research Farm.



Image 2 : Arrangement of Paddy straw mulch in experimental field at Crop Research Farm.




Image 3 : Different types of Mulches which I have took for my Research during Rabi season, 2021-22



Image 4 : The Recommended Dose of Fertilizer (80:40:40 NPK kg/ha) will be applied, as Half dose of Nitrogen along with full dose of Phosphorus, Potassium and Sulphur as basal and remaining half dose of Nitrogen after 30 Days of Sowing (DAS).



Image 5 : Harvesting in Experimental field at Crop Research Farm during Rabi season, 2021-22

UNDER REVIEW