

Standardization of Aonla Candy (*Emblica officinalis L.*) Cv. Kanchan

ABSTRACT

An experiment was carried out at the Post Harvest Technology, Department of Horticulture, SHUATS, Prayagraj (Uttar Pradesh) during the year 2021 - 2022. The experiment consisted of 6 different treatments comprising the different syrups (sugar syrup, rose syrup, tulsi syrup, ginger syrup) with the addition of the standard recommended dosage of citric acid and sodium benzoate. This investigation was laid out in a completely randomized design with three replications. After preparation, the aonla candy samples were evaluated for Physico-chemical alterations, and sensory evaluation was done using a 9-point hedonic scale that was tested on a panel of 7 experts. These candies were stored for about 90 days at ambient temperature. From storage studies, it was revealed that T₂ (Aonla + 68% Sugar syrup + 10% Spices) is most suitable in terms of their Physico-chemical properties and organoleptic test of aonla candy. The effect of storage on Physico-chemical and organoleptic characteristics were also observed.

Keywords: - Aonla candy, citric acid, sodium benzoate, syrup.

INTRODUCTION

Aonla or Indian Gooseberry (*Emblica officinalis*) is an indigenous fruit to the Indian subcontinent. Aonla is thought to be a native of India, Sri Lanka, Malaysia, or China (Kalra, 1998). Aonla fruit is a rich source of vitamin 'C', the vitamin C content in aonla varies from 200-900 mg /100 g depending upon the variety and size of the fruit (Barthakur and Arnold, 1991). Aonla is highly valued among indigenous medicines. It has astringent, cooling, refrigerant, diuretic, and laxative properties. Dried fruits have been reported to be useful in haemorrhages, diarrhoea, dysentery, anaemia, jaundice, dyspepsia, and cough. Aonla is a fruit that is now underutilized but has huge potential in the global market. It has got great potential in processed forms, such as candy, pickles, preserve (murabba), sauce, jam, jelly, dried chips, tablets, etc. Candy is the most popular aonla product. A fruit, when impregnated with sugar free of syrup, drained and dried, is called candied fruit. It is plump, tender, and exceedingly sweet with a strong flavor and without sticking. Aonla candy can be improved by the addition of different syrups in the aonla candy like rose syrup, tulsi syrup, and ginger syrup. Rose or Queen of flowers (*Rosa* spp.) belongs to the family of Rosaceae. Rose is a highly nutritive flower with a high content of Vitamin C, carotenoids, phenolic components, and some

minerals and essential oil. Tulsi (*Ocimum sanctum*) is known as Holy Basil in English and Tulasi in Sanskrit. The extricates acquired from the plant are widely brought to use for relieving different illnesses, for example, the basic cold, irritation, intestinal sickness, coronary illness, migraines, stomach issues, kidney stones, heart issues, and curing malaria. Ginger (*Zingiber officinalis L.*) belongs to the family of Zingiberaceae. Ginger contains gingerol and oleoresin, (a combination of volatile oils and resin) that accounts for the characteristic aroma and therapeutic properties. Components of gingerol possess beneficial properties for the treatment of poor digestion, heartburn vomiting, and preventing motion sickness.

MATERIAL AND METHODS

Raw material

Healthy uniform size aonla, free from diseases, pests, and bruises were randomly selected and brought from our research field. Other ingredients like sugar, rose, tulsi, ginger, spices (jeera, fennel, black salt, asafoetida, lemon, black pepper, and chat masala), and citric acid were brought from the local shops in Prayagraj.


Preparation of Candy

Aonla candy was prepared with different syrup solutions i.e., sugar syrup, rose syrup, tulsi syrup, and ginger syrup. After preparation of different syrup solutions (68%).

Washed aonla fruits were soaked with 2% salt (24hrs) and after washing, soaked with 2% alum (24hrs). After washing, blanched aonla fruits till they become soft and then allowed to cool, remove seeds and separate blanched fruits into pieces. Steeped aonla pieces in different syrup. After completion of steeping time, the syrup was drained and aonla pieces were spread on trays.

Then aonla pieces in the tray were kept for drying in the dehydrator for 60⁰C (6-7hrs). After drying samples were collected and packed in LDPE bags for storage purposes. Candies were stored at room temperature for 3 months and evaluated at 30 days intervals for storage studies.

Chart 1 : FLOW CHART OF AONLA CANDY PREPARATION:



Treatment details

T₁ - Aonla + 68% Sugar Syrup

T₂ - Aonla + 68% Sugar Syrup + 10% Spices

T₃ - Aonla + 68% Rose Syrup

T₄ - Aonla + 68% Tulsi Syrup

T₅ - Aonla + 68% Ginger Syrup

T₆ - Aonla + 48% Sugar Syrup + 10% Rose Syrup + 10% Tulsi Syrup

Physico-chemical and organoleptic quality of aonla candy

Titration acidity (%)

Titration acidity was determined by titrating a known quantity of sample (10ml) of the homogenized sample was taken and made up to 100 ml volume in a volumetric flask. The contents were filtered through Whatman No. 1 filter paper. Titration against 0.1N NaOH was performed with a 10 ml aliquot using phenolphthalein as an indicator. The turn of the aliquot to light pink color which persists for 15 seconds was considered an endpoint. The titration acidity was estimated in terms of percent citric acid (Ranganna, 1986).

pH

The sample was soaked in distilled water till it softens and ground along with the little amount of distilled water. Then the pH was noted with the help of an electronic pH meter.

Total soluble solids (%)

The percentage of total soluble solids was determined by using a hand refractometer (Erma, Japan) by placing a drop of the filtered juice on the prism of the refractometer and observing the coincidence of the shadow of the sample with the reading on the scale and expressed as °Brix.

Ascorbic acid (mg/100g)

Determination of ascorbic acid was done by the 2, 6 – dichlorophenol indophenols dye method as described by Ranganna (1997). A known quantity of sample was blended with 3 percent metaphosphoric acid (HPO₃) to make the final volume of 100 ml and then filtered. A known quantity of aliquot was titrated against 0.025 percent 2, 6 - dichlorophenol indophenols dye to a pink color endpoint. The ascorbic acid content of the sample was calculated taking into consideration the dye factor and expressed as mg of Ascorbic acid per 100g.

Organoleptic evaluation

To assess consumer preference, the sensory evaluation of the experimental sample was conducted at different intervals by a panel of 7-8 judges. The sample was evaluated for color and appearance, taste, aroma, and overall acceptability.

RESULTS AND DISCUSSION

Titration Acidity (%)

During the storage term (90 days) at ambient room temperature, pectic substances have been reported to increase the titration acidity of aonla candy. The maximum value (0.68) was observed in the combined treatment of aonla candy with T₂ (Aonla + 68% Sugar syrup + 10% Spices) and the minimum value (0.53) in treatment T₁ (Aonla + 68% Sugar syrup). Similar results were found by Kumar *et al.*, (2020) in aonla candy and Mahato *et al.*, (2020) in unripe mango candy.

pH

The gradual decrease in pH during the storage period (90 days) was reported from aonla candy. The decrease in pH might be due to the overall increase in acidity of stored candy. The maximum value (3.54) was observed in the combined treatment of aonla candy with T₂ (Aonla + 68% Sugar Syrup + 10% Spices) and the minimum value (3.22) in the treatment of T₃ (Aonla + 68% Rose syrup). Similar results were found by Navitha *et al.*, (2018) in ber candy and Dinde *et al.*, (2021) in karonda candy.

Total Soluble Solids

This gradual increase in total soluble solids during the storage period (90 days) might be due to the conversion of polysaccharides into sugars during the hydrolysis process. The maximum value (71.12) was observed in the combined treatment of Aonla Candy with T₂ (Aonla + 68% Sugar Syrup + 10% Spices) and minimum value (70.03) in T₁ (Aonla + 68% Sugar syrup). Similar results were found by Dwivedi S. and Pandey A., (2017) studied aonla preserves and Balaji *et al.*, (2014) on honey coating of aonla candy.

Ascorbic Acid

This gradual reduction in ascorbic acid during the storage period (90 days) might be due to oxidation by trapped oxygen in the packaging container, which results in the formation of dehydroascorbic acid. The maximum value (108.33) was observed in the combined treatment of Aonla Candy with T₂ (Aonla + 68% Sugar Syrup + 10% Spices) and the minimum value (89.73) in T₁ (Aonla + 68% Sugar syrup). Similar observations were reported by Kumar and Singh., (2001) in aonla products and Deepika *et al.*, (2016) to maintain the quality of enriched fruit bars during storage.

Organoleptic evaluation

In the organoleptic evaluation such as color and appearance, taste, aroma, and overall acceptability. Sensory scores for treatment T₂ (Aonla + 68% Sugar Syrup + 10% Spices) were found to be highest in all parameters of organoleptic attributes. The highest overall acceptability score (8.13) indicated that it was well-received by the judges.

Shelf life of aonla candy

From the shelf life point of view during the storage period, the highest was recorded in T₂ (Aonla + 68% Sugar syrup + 10% Spices) at 89.23 and minimum (87.93) in treatment T₃ (Aonla + 68% Rose syrup) respectively.

Table 1: Changes in acidity, pH, and TSS in the aonla candy during storage.

| Treatments | Acidity (%) | | | | pH | | | | TSS (°Brix) | | | |
|----------------|-----------------------|-------|-------|-------|-----------------------|-------|-------|-------|-----------------------|-------|-------|-------|
| | Storage period (days) | | | | Storage period (days) | | | | Storage period (days) | | | |
| | 0 | 30 | 60 | 90 | 0 | 30 | 60 | 90 | 0 | 30 | 60 | 90 |
| T ₁ | 0.40 | 0.45 | 0.49 | 0.51 | 3.63 | 3.56 | 3.52 | 3.47 | 69.47 | 69.62 | 69.83 | 70.03 |
| T ₂ | 0.62 | 0.64 | 0.66 | 0.68 | 3.69 | 3.64 | 3.58 | 3.54 | 70.23 | 70.47 | 70.68 | 71.12 |
| T ₃ | 0.43 | 0.47 | 0.51 | 0.55 | 3.36 | 3.32 | 3.27 | 3.22 | 69.75 | 69.98 | 70.22 | 70.42 |
| T ₄ | 0.51 | 0.53 | 0.56 | 0.58 | 3.58 | 3.54 | 3.49 | 3.44 | 69.58 | 69.85 | 70.04 | 70.23 |
| T ₅ | 0.53 | 0.56 | 0.60 | 0.62 | 3.56 | 3.52 | 3.48 | 3.43 | 69.96 | 70.26 | 70.49 | 70.67 |
| T ₆ | 0.56 | 0.60 | 0.62 | 0.64 | 3.45 | 3.41 | 3.38 | 3.34 | 70.05 | 70.31 | 70.66 | 71.01 |
| F-test | S | S | S | S | S | S | S | S | S | S | S | S |
| S.Ed(±) | 0.050 | 0.045 | 0.035 | 0.042 | 0.041 | 0.038 | 0.034 | 0.041 | 0.026 | 0.039 | 0.042 | 0.052 |
| CD @ 5% | 0.110 | 0.098 | 0.077 | 0.093 | 0.091 | 0.084 | 0.076 | 0.090 | 0.056 | 0.085 | 0.093 | 0.114 |

Table2: Changes in ascorbic acid, color and appearance, and taste in the aonla candy during storage.

| Treatments | Ascorbic acid (mg/100g) | | | | color and appearance | | | | taste | | | |
|----------------------|-------------------------|----------|----------|----------|-----------------------|----------|----------|----------|-----------------------|----------|----------|----------|
| | Storage period (days) | | | | Storage period (days) | | | | Storage period (days) | | | |
| | 0 | 30 | 60 | 90 | 0 | 30 | 60 | 90 | 0 | 30 | 60 | 90 |
| T₁ | 98.77 | 97.53 | 94.00 | 89.73 | 6.37 | 6.23 | 6.12 | 6.02 | 6.46 | 6.37 | 6.28 | 6.18 |
| T₂ | 125.87 | 120.06 | 114.76 | 108.33 | 8.66 | 8.55 | 8.46 | 8.36 | 8.14 | 8.09 | 8.01 | 7.96 |
| T₃ | 116.47 | 113.19 | 108.46 | 102.67 | 7.52 | 7.41 | 7.32 | 7.19 | 6.04 | 6.01 | 5.96 | 5.92 |
| T₄ | 102.67 | 98.77 | 95.15 | 90.72 | 7.81 | 7.72 | 7.63 | 7.54 | 7.39 | 7.35 | 7.26 | 7.16 |
| T₅ | 108.33 | 105.44 | 104.35 | 97.53 | 7.96 | 7.84 | 7.74 | 7.61 | 7.44 | 7.38 | 7.33 | 7.25 |
| T₆ | 109.90 | 107.01 | 102.78 | 98.86 | 6.57 | 6.43 | 6.33 | 6.19 | 6.29 | 6.22 | 6.17 | 6.03 |
| F-test | S | S | S | S | S | S | S | S | S | S | S | S |
| S.Ed(±) | 2.078 | 2.934 | 3.605 | 2.177 | 0.190 | 0.194 | 0.196 | 0.191 | 0.110 | 0.103 | 0.107 | 0.109 |
| CD @ 5% | 4.578 | 6.463 | 7.941 | 4.796 | 0.418 | 0.427 | 0.432 | 0.422 | 0.242 | 0.226 | 0.235 | 0.239 |

Table3: Changes in aroma, overall acceptability, and shelf life in the aonla candy during storage.

| Treatments | aroma | | | | overall acceptability | | | | Shelf life |
|----------------------|-----------------------|----------|----------|----------|-----------------------|----------|----------|----------|-----------------------|
| | Storage period (days) | | | | Storage period (days) | | | | Storage period (days) |
| | 0 | 30 | 60 | 90 | 0 | 30 | 60 | 90 | 90 |
| T₁ | 6.34 | 6.27 | 6.16 | 5.98 | 6.62 | 6.54 | 6.47 | 6.38 | 88.63 |
| T₂ | 8.22 | 8.12 | 8.06 | 7.97 | 8.38 | 8.30 | 8.24 | 8.13 | 89.23 |
| T₃ | 6.81 | 6.77 | 6.7 | 6.63 | 6.39 | 6.33 | 6.27 | 6.17 | 87.93 |
| T₄ | 6.98 | 6.91 | 6.82 | 6.74 | 7.27 | 7.19 | 7.10 | 6.97 | 88.10 |
| T₅ | 7.54 | 7.45 | 7.35 | 7.26 | 7.54 | 7.46 | 7.40 | 7.33 | 89.09 |
| T₆ | 6.68 | 6.60 | 6.50 | 6.42 | 6.44 | 6.38 | 6.30 | 6.19 | 88.41 |
| F-test | S | S | S | S | S | S | S | S | S |
| S.Ed(±) | 0.266 | 0.269 | 0.270 | 0.276 | 0.110 | 0.112 | 0.115 | 0.141 | 0.178 |
| CD @ 5% | 0.586 | 0.592 | 0.595 | 0.608 | 0.242 | 0.247 | 0.254 | 0.310 | 0.393 |

CONCLUSION

In this investigation, the treatment T₂ (Aonla + 68% Sugar syrup + 10% Spices) was found most suitable treatment in terms of Physico-chemical properties i.e., total soluble solids (71.12), titrable acidity (0.68), pH (3.54) and ascorbic acid (108.33) and organoleptic test i.e., color and appearance (8.36), taste (7.96), aroma (7.96) and overall acceptability (8.13). The maximum shelf life (89.23) was recorded in T₂ (Aonla + 68% Sugar syrup + 10% Spices). Similarly, the treatment T₂ (Aonla + 68% Sugar syrup + 10% Spices) showed the highest BC ratio (1:2.02).

REFERENCES

1. Balaji, V., Prasad, V.M. and Saroj, P.L., 2014. Comparative study of varieties, honey coating, and storage durations on aonla candy. *Indian J. Hort.* 71 (1): 104-108.
2. Barthakur, N.N. and Arnold, N.P. (2001). Chemical analysis of the emblic (*Phyllanthus Emblica* L.) and its potential as a food source. *Scientia Horticulturae.*, 47: 99- 105.
3. Deepika, Panja, P., Marak, D.S. and Thakur, P.K. (2016). Effect of packaging on quality of enriched fruit bars from aonla (*Embllica officinalis* G.) during storage. *International Journal of Agriculture, Environment and Biotechnology.*, 9(3): 411-419.
4. Dinde A.B., Joshi, P.S., Dalal, S.R. and Tayade S.A. (2021). Effect of Different Recipes on Storage of Karonda Candy. *International Journal of Agriculture Sciences*, ISSN: 0975-3710 & E-ISSN: 0975-9107, Volume 13, Issue 2, pp.- 10677-10679.
5. Dwivedi, S.K., and Pandey, A. (2017). Development of a protocol for preparation and preservation of ginger-flavored aonla candy for nutritional and biochemical evaluation. *The Pharma Innovation Journal.*, 6(11): 78-83.
6. Kalra, S.K., Tandon, D.H. and Singh, B.P. (1998). Simplified processing of aonla fruits for preserve making. Central Institute for Subtropical Horticulture, Lucknow. *Beverage Food World.*, 25(3): 35-36.
7. Kumar, R., Pathak, S. (2020). Studies on Preparation and Storage of Aonla Candy Enriched with Different Natural Oil/Extract (*Embllica officinalis* L.) cv. NA-7. *Int.J.Curr.Microbiol.App.Sci.*, 9(10):3285-3299.
8. Kumar, S. and Singh, I.S, (2001). Storage studies of aonla fruit products at ambient temperature. *Progressive Horticulturae*, 33(2): 169-173.

9. Mahato, A., Chakraborty, and Baidya, B.K. (2020). Preparation and evaluation of fruit candy from unripe mango. *International Journal of Chemical Studies.*, 8(1): 2727-2731.
10. Navitha, D. and Mishra, S. (2018). Standardization of a recipe for the preparation of candy from ber. *Research Journal of Chemical and Environmental Sciences. Res J. Chem. Environ. Sci.* Vol 6 (4):15-18.
11. Ranganna, S. (1986). *Handbook of Analysis and Quality Control for Fruit and Vegetable Products.* McGraw-Hill Tata Pub. Co. Ltd., New Delhi. pp. 7-12 and 109.

UNDER PEER REVIEW