

AN OVERVIEW: INTERACTION BETWEEN MICROBIAL ENDOPHYTES AND ROOT EXUDATES

Abstract:

Rhizosphere bacterial endophytes reside in a vast number of plant tissues as a microbiome and from their they positively induce plant growth. Endophytic community structure within the plant cells is dynamic and is influenced by various abiotic and biotic factors such as soil conditions, biogeography, plant species, microbe–microbe interactions and plant–microbe interactions, both at local and larger scales. Root exudates are a complex mixture of soluble organic substances, which constitute sugars, amino acids, organic acids, enzymes, and other substances. The root surroundings also contain cell lysates, which are released when root cortex cells are ruptured by external mechanical damage, microbial graze, microbial infection, or the natural pores formed in the process of lateral roots emergence that break through the root epidermis. Since root exudates have abundance roles in plant growth promotion at different stages of plant life cycle. In view of this hypothesis, the present review has attempted to describe the various functions of root exudates and their interactions with microbial community present with in the plant tissues and their positive effects on plant health.

Keywords: Root exudates, plant growth promotion, endophytes, biocontrol.

Introduction:

The interaction between plants and microorganisms in the soil is well recognized. Hiltner [1]; Hartmann et al. [2] initial observations made evidenced that microorganisms (both beneficial & pathogenic) are more abundant in the rhizospheric soil surrounding the plant roots than in soil that are away from the root area. Plant roots exude various organic compounds that promote microbial growth conditions which shows major impact on the

composition of the rhizosphere microbiome [3,4,5]. In recent times, research is more focused on the composition of the rhizosphere microbiome, exploring the impact of them on plant growth and health [6,7,8]. The microbial endophyte life cycle is inextricably linked with that of the host plant and is influenced by the plant species, plant life stage and plant health. Some characteristics of plant-endophytic colonization are well programmed, for example, endophytic microorganism take advantage of plant wounds for entry into tissues, but plants also control which endophytes are favored before and after tissue penetration [9,10]. This results in plants of different species grown on the same soil harboring distinct groups of endophytes [11,12]. For evaluation of microbial population, Dong et al. [13] stated that monocots have 100-fold higher number of *Klebsiella pneumonia* cells than dicots. The reason for differential populations is unknown, but in fact, the alterations root architecture, host-bacteria interactions, precise selection/recruitment, or the types of chemicals in root exudates attracting specific endophytes may be factors. On other hand, root exudates may alter microbial populations in the rhizosphere, determining their abundance and diversity of microbial species that colonize the host plant [8]. The nature of chemicals in root exudates released by plants can act as signals for beneficial bacteria identify suitable hosts [14,9,15].

Plants produce various secondary metabolites to protect themselves against pathogens and to attract beneficial microbes. *For example*, the symbiotic associations of *Rhizobium* and mycorrhizal (VAM) interactions in which roots secrete compounds that act as messengers to attract *Rhizobium* and arbuscular mycorrhizal fungi [16,17]. Regardless of progression in understanding the symbiotic plant-microbe interactions, little progress has been made in genetic and biochemical changes responsible for the attraction of non-symbiotic microbes towards plants. Plant associations with plant growth-promoting rhizobacteria (PGPR) provide protection from soil-borne plant pathogens by antagonistic mechanisms [18,19]. Plant and endophytic colonization are a complex process that requires the capacity to compete with other bacteria in the rhizosphere soil and communicate, interact with plant roots. Root exudates are chemically diverse attracting microorganisms to colonize roots, or to be able to colonize the internal plant tissues for endophytes. A whole genome sequence approach was studied by Shidore et al. [20] to know the endophyte genes that help in root colonization. Furthermore, they also exposed the bacterial endophyte *Azoarcus* spp. strain BH72 to exudates of rice plants (*Oryza sativa* cv. *Nipponbare*) and analyzed the endophyte's gene expression changes. As expected, many genes known to be involved in rhizosphere competence were found to be upregulated; Several other genes were altered in substantiate their role for endophytic colonization, such as gene for minor pilin PilX, signal transduction

proteins and a serine–threonine kinase as a putative component of the type IV secretion system (T6SS). Based on the results of Shidore et al. [20] it was suggested that rhizosphere competence, as well as endophytic colonization was induced by root exudates. Although evidence exists for intra-plant communication, but, as such proven reports were not existed demonstrating whether plants exude specific chemical signals through their roots to attract beneficial bacteria in the rhizosphere. Furthermore, it is very hypothetical to explain whether shoot infection by pathogenic bacteria will induces recruitment of beneficial rhizobacteria to the root surface. The microbe-microbe signaling under distress conditions, especially between aboveground and belowground tissues, are poorly documented. Such signaling, although potentially complex due to the involvement of significant physical distances, may be an important and effective strategy in plant defense that has thus far been overlooked [15]. In this scenario, the present review focused on the interaction between the microbial endophytes and plant root exudates and their mechanisms for plant health.

Interaction between endophytes and root exudates:

The microorganisms with colonizing ability in colonization around or in the plant are collectively called colonization properties. In the interactive colonization processes, communication between the plant and bacterium (and vice versa) has a key role [9]. Bacterial root colonization often starts with the recognition of specific compounds in the root exudates [21]. These compounds probably also have major roles in below-ground community interactions [18]. At same time plants communicate with almost all the groups of commensalistic, mutualistic, symbiotic and pathogenic microorganisms *via* chemicals exuded by roots [22]. Chemotaxis is important factor in colonization around roots with the help of motility directed by chemotactic response to root exudates [23]. Chemotaxis is not common in all endophytic species, and other multiple parallel paths have evolved during different plant–microbe interactions. Root-exuded organic acids are major chemo-attractants for *P. fluorescens* and has close association with tomato plant [21], whereas carbohydrates and amino acids attract *Corynebacterium flavescens* and *Bacillus pumilus* to rice [24]. The apparent specificity in these interactions probably relates to bacterial nutritional requirements and, in each of these cases, chemotaxis towards specific resources probably determines the specificity of the interaction.

Among plant materials, seeds carry a range of endophytes [25,26]. Seedborne endophytes confirm their presence in new germination. Unlikely, in vegetative propagation (such as potatoes or sugarcane) transmission of their endophytes to the next generation and

would not require the infection process. In the rhizosphere there is a selection of the microorganisms that can survive in the root exudates and compete with others. Rosenblueth and Martínez-Romero [27] found that some strains can act as both epiphytes and endophytes for colonizing the rhizosphere surface and inside tissues. Beneficial microbes in the rhizosphere is important not only as the first step in biocontrol, but also is crucial in plant growth. It has been assumed that many microorganisms are attracted by carbon-containing compounds exuded by plant roots. Indeed, this phenomenon was first described by Hiltner [1], who observed increased numbers and activity of microorganisms near plant roots. Evidence continues to accumulate which suggests a major role for root exudates in determining the fate of microbial species in the rhizosphere (saprophytes/endophytes).

The most complex interactions in the rhizosphere surrounding were differentiated by chemical, physical and biological properties among land plants. For the last one decade, the biological importance of root exudates by plants is gained as one of important trait in the rhizosphere in mediating interactions among plants and microbes [28,29,30,31]. Root exudates classified as low molecular weight compounds such as amino acids, organic acids, sugars, phenolics, and other secondary metabolites and high molecular weight compounds include mucilage and proteins. Root exudation represents a major carbon cost to the plant [32]. Plant roots even responds to various environmental factors through the secretion of a wide range of compounds, depending on nutritional status and soil conditions [33,34]. This action interferes with the plant-bacteria interaction and an important factor contributing to the efficiency of the inoculant [33,35,34]. Apart from low and high molecular weight compounds, root exudation includes the secretion of ions, free oxygen species, water, enzymes, mucilage and a diverse array of carbon containing metabolites [22]. In plants, roots alone excretes 10-44% of photosynthetically fixed carbon, which acts as primary energy source, signaling molecules or antimicrobials for soil microorganisms [36]. The release of exudates purely depends on plant age, genotype and microorganisms response and interaction with different host plants [37,38,39]. Thus, artificial microbial inoculants are generally guided to the one specific plant from which the bacterium was isolated.

Root exudates in plant growth promotion

Plant–microbe interactions can positively influence plant growth through a variety of mechanisms includes nitrogen fixation by alpha and betaproteobacteria [40], biotic and abiotic stress tolerance communicated by the presence of endophytic microbes [41], and

direct and indirect applications by PGPR [42]. Several proved interactions prove the importance of root exudates in positive plant–microbe interactions, where they act either directly or indirectly for healthy plant growth promotion [43]. The classical example for positive plant-microbe interaction is legume roots and *Rhizobia* bacteria interactions, which can form dinitrogen-fixing nodules in the roots of legumes. Similarly, vascular arbuscular mycorrhizal (VAM) fungi symbiotically interacts with plant roots more than 80% of terrestrial plants. Mycorrhizal fungus and bacterial rhizobial associations are thought to derived from a common-ancestral and plant–microbe interaction likely of fungal origin and demonstrated that some host proteins regulates both mycorrhizal and rhizobial interactions with roots [44]. Root exudates play a role in the recognition of mycorrhizal fungi with compatible host plant [45,46]. Although root exudates have been suspected to play a communicative role in mycorrhizal associations, the identification of specific molecule interactions from AMF and host remains elusive. Akiyama *et al.* [47] stated that, a sesquiterpene called strigolactone 5-deoxystrigol was identified in the root exudates of *Lotus japonicus* which is responsible for inducing VAM fungi hyphal branching in germinating spores. In the presence of VAM symbiosis, plants trade carbon with phosphate from their fungal partners [48,49,50]. Recently, a leucine-rich-repeat receptor kinase (SymRK) was identified as a common genetic basis for plant root endosymbiosis with AMF, *Rhizobia*, and *Frankia* bacteria [51].

Root exudates in biocontrol activity

Certain strains of root-colonizing fluorescent pseudomonads can provide efficient protection of crop plants against a variety of soilborne phytopathogenic fungi, notably by the secretion of extracellular antimicrobial secondary metabolites into the rhizosphere [52]. In another study *in vitro* screened *Bacillus subtilis* and *Bacillus licheniformis* MSEB (Maize seed endophytic bacteria) showed strong antifungal property against three broad fungal pathogens *Rhizoctonia solani*, *Sclerotia rolfsii* and *Macrophomina phaseolina* and these were seed treated with tagged rifampicin mutant endophytic bacteria (*Bacillus* spp.) before sowing showed potent root and stem colonizers [53]. Werra *et al.* [54] investigated the impacts of 63 different low-molecular-weight compounds, most of them plant derived, on the *in vitro* expression of two antifungal biosynthetic genes by the plant-protecting rhizobacterium *Pseudomonas fluorescens* CHA0. Most of the compounds tested, affected the expression of one or both antifungal genes. This suggests that biocontrol activity in plant-beneficial pseudomonads is modulated by plant-bacterium signaling. It was well documented that

Trichoderma spp. as effective biological control agents of plant diseases caused by soilborne fungi [55]. Seeds treated with *Trichoderma lignorum* plants had roots with lower levels of *Rhizoctonia solani* in their rhizosphere than roots of untreated seeds. *Trichoderma* had little effect on the survival of *Rhizoctonia solani* in non-rhizosphere soil. However, application of *Trichoderma lignorum* as a wheat-bran preparation, conidial suspension or seed coating reduced the pathogen counts in the rhizosphere soil of beans [56].

Compounds secreted from plant roots often alter the gene expression of benefitted microorganisms by acting as signals that either induce or repel the interaction with beneficial and or harmful strains, respectively. The prominent unknown fact is whether two distinct groups of beneficial bacteria, such as soil microorganisms (not plant associated) and plant endophytic strains (plant colonizers) strains, respond uniformly or variably to the root exudates. In a study, *Bacillus mycoides*, a potential biocontrol and plant growth promoting bacterium strain, was isolated from the endosphere of potatoes and also from soil of the same geographical region. To characterize their phenotypic differences, the whole genome of the two strains were sequenced and the transcriptome response to potato root exudates was compared. The transcriptome profiles showed that endophytic strains responded more distinct than the soil-derived isolate and a higher number of significant differentially expressed genes were detected [57]. Biocontrol inoculants often show inconsistency in their efficacy at field scale and the reason for this remains often unclear. A high rhizosphere competence of inoculant strains is assumed to be a key factor for successful biocontrol effects as the biocontrol strain must compete with the indigenous microbial community in the rhizosphere. It is known that many factors, among them plant species and soil type shape the rhizosphere microbial community composition. However, microbial community composition in the rhizosphere can also be influenced by the presence of a pathogen [58]. The biological control process mediated by microbes relies on multiple interactions among plants, pathogens and biocontrol agents (BCAs). One such efficient BCA is *Bacillus cereus* AR156, a bacterial strain that controls a broad spectrum of plant diseases and potentially works as a microbe elicitor of plant immune reactions. The strain *B. cereus* AR156 has induced some specific components in plant root exudates was probably involved in further biocontrol processes [59].

Root exudates and their mechanisms to establish endophytic colonization:

Plants often use transport mechanisms to distribute and produce compounds into the rhizosphere surroundings [60,61]. In general, root exudates can be released by plant roots *via*

either passive (diffusates) or active (secretions) mechanisms by different transport methods. Most of them were low molecular weight organic compounds through a passive process. Small polar and uncharged molecules are transported by direct passive diffusion, a process that depends on (i) membrane permeability, (ii) the polarity of the exuded compounds and (iii) cytosolic pH [60]. Plant root cells secrete other compounds, such as secondary metabolites, polysaccharides and proteins, with the help of different membrane-bound proteins [61]. Such as, ATP-binding cassette (ABC) transporters [62,63,64,65], MATE family (multidrug and toxic compound extrusion) [66], the major facilitator superfamily [67] and the aluminum-activated malate transporter family [61]. Although, there are not many reports on the details of these membrane-bound transport protein functions, they have been associated with the transport of a wide range of compounds into the rhizosphere.

For the last two decades, interactions in rhizosphere mediated by root exudates were widely studied [68,69,70,71,72,73]. Plant root associated phytochemicals can facilitate several interactions, such as plant–plant, plant–microbe, and plant–faunal. These interactions can vary from neutral to beneficial or deleterious [74,75]. In some cases, microbes can transition from pathogenic to symbiotic depending upon the environmental conditions [76]. For example, *rhizobia*, symbiotic nitrogen (N)-fixing bacteria, range from a symbiotic to a neutral interaction with plants depending on soil N levels [77,78]. Furthermore, under N-limiting conditions, legumes secrete more flavones and flavonols to attract and initiate legume–rhizobia symbiosis [79,80].

Table 1: Potential functional role of root exudate components identified from different plant rhizospheres, for example tomato, Brassica napus etc., adapted from [81,69,82].

Exudates component	Functions	Specific compounds identified in root exudates
Organic acids	Nutrient source	Citric, glutaric, oxalic, malonic
	Chemoattractant signals to microbes	Malic, aldonic, fumaric, erythronic
	Chelators of poorly soluble mineral	Succinic, ferulic, acetic, butanoic

	<p>nutrients</p> <p>Acidifiers of soil</p> <p>Detoxifiers of Al</p> <p>nod gene inducers</p>	<p>Butyric, syringic, valeric, rosmarinic, lactic, glycolic</p> <p>trans-cinnamic, piscidic, formic</p> <p>aconitic, pyruvic</p> <p>vanillic, tetronic</p>
Amino acids	<p>Nutrient source</p> <p>Chelators of poorly soluble mineral nutrients</p> <p>Chemoattractant signals to microbes</p>	<p>a- and b-alanine proline</p> <p>asparagine, valine, threonine, aspartate, tryptophan cystein,</p> <p>ornithine, cystine, histidine, glutamate, arginine, glycine,</p> <p>homoserine, isoleucine, phenylalanine, leucine, - Aminobutyric acid,</p> <p>lysine a-Aminoadipic acid, methionine, serine, homoserine</p>
Sugars & Vitamin	<p>Promoters of plant and microbial growth nutrient source</p>	<p>Glucose, desoxyribose, oligosaccharides galactose, biotin,</p> <p>maltose, thiamin, ribose, niacin, xylose, raffinose pantothenate,</p> <p>rhamnose, riboflavin, arabinose, fructose</p>
Proteins and enzymes	<p>Catalysts for P release from organic molecules</p> <p>Biocatalysts for organic matter transformations</p> <p>Plant defense</p>	<p>Acid/alkaline, phosphatase</p> <p>amylase, invertase, protlease</p> <p>PR proteins, lipases, b-1,3-glucanases</p>
Inorganic ions and gases	<p>Chemoattractant signals to microbes</p>	<p>HCO₃⁻ OH⁻ H₂ CO₂ H₂</p>
Phenolics	<p>Nutrient source</p> <p>Chemoattractant signals to microbes</p> <p>Microbial growth promoters</p> <p>nod gene inducers and inhibitors in rhizobia</p> <p>Resistance inducers against</p>	<p>Liquiritigenin, luteolin</p> <p>Daidzein, 40,7-dihydroxyflavanone</p> <p>Genistein, 40,7-dihydroxyflavone</p> <p>Coumetrol, 4,40-dihydroxy-2'-methoxychalcone</p> <p>Eriodictyol, 40-7-dihydroxyflavone</p> <p>3,5,7,30-tetrahydroxy- 40methoxyflavone</p>

	phytoalexins Chelators of poorly soluble mineral nutrients Detoxifiers of Al Phytoalexins against soil pathogens	naringenin isoliquiritigenin, 7,30-dihydroxy-40-methoxyflavone umbelliferone, (p)- and (")- catechin
Root border cells	Produce signals that control mitosis Produce signals controlling gene expression Stimulate microbial growth Release chemoattractant Synthesize defense molecules for the rhizosphere Act as decoys that keep root cap infection-free Release mucilage and proteins	

Transport mechanisms of root exudates:

During the last one decade many researchers substantially gained progress in isolation and characterization of root exudates and their important interactions in rhizosphere soil. However, the mechanism of root exudates secretion is poorly understood. The production and release of root-derived compounds are commonly constitutive but, may be induced by biotic or abiotic stress as described previously in this article. The secretion of root exudates by plant roots is initially thought to be a passive process which dependent on three separate pathways: diffusion, ion channels and vesicle transport [83].

Diffusion process:

During diffusion, small polar molecules and uncharged molecules are transported through permeability nature of lipid membranes [84]. This passive diffusion process depends on membrane permeability [85] and cytosolic pH [86]. Other compounds like sugars, amino acids and carboxylate anions were transported across membranes by the aid of proteins, and their direction of movement is dependent on their electrochemical gradient that allows them to pass from the cytoplasm of intact root cells to the soil. Samuel et al. [87] showed that the

large cytosolic K^+ diffusion potential and the extrusion of protons through ATPase generate a positively charged gradient that promotes the release of carboxylate anions. Factors that affect membrane integrity could also promote the release of organic acids [88]. In such cases, mediate the controlled release of these compounds mediated by anion channels in the roots [89,90,91,92].

Ion-exchange method:

Sugars, amino acids and metals were specifically transported by the compounds that in turn involved in the secretion of other compounds from root cells [93,94,95,96,97,98,99,100]. Plants cells possess metal homeostasis to avoid excess concentrations of free metal ions inside the cells (e.g. Fe, Zn, Mn and Cu); these mechanisms involve coordination of metal ion transporters for uptake, translocation and compartmentalization [101]. For e.g. graminaceous plants secrete mugineic acid, a metal-binding ligand secreted into the rhizosphere, and form Fe(III)-MA ligand to reduce the Fe (iron) toxicity and then enter into the root cells *via* a specific transporter YSL identified in maize [102,103]. Recent evidence demonstrated that the silicon efflux transporter from rice is involved in efflux of silicon from root cells [104,105]. Similarly, it has been demonstrated that malate transporter (AtALMT1) plays a critical role in releasing malate from root cells under aluminium toxicity in *Arabidopsis* [106]. There are other transporters like monosaccharide transporters that are involved in transporting hexoses, pentoses including ribose and polyols such as myo-inositol and glycerol [107,108].

Vesicular transport system:

Vesicular transport mechanisms involves in secretion of high-molecular weight compounds by roots [109]. Knowledge of the vesicle-mediated trafficking of proteins is well understood [110], but the mechanism of vesicle-mediated transport of phytochemicals is not fully characterized [111,112]. There are reports demonstrating that plant defense responses are accompanied by trafficking of antimicrobial compounds to the site of pathogen infection. For e.g. pigmented vesicles accumulate on sorghum leaves at sites of attempted fungal infection they often termed as antimicrobial pigments [113,114]. Similar reports by [115], Yazaki et al. [116,117] suggested the pigmented antimicrobial naphthoquinones are secreted in response to fungal elicitation. into the apoplast of the boraginaceous plant, *Lithospermum erythrorhizon* by a vesicle-mediated mechanism.

Membrane bound transport proteins:

Membrane-bound proteins offer an alternative transport mechanism for the release of metabolites from cells and different proteins have been involved in the transport of secondary compounds across plant membranes. Among these, some localize to the plasma membrane which directly export compounds from the cell. Others localize to inner membranes where they can help sequester compounds into subcellular compartments or perhaps load vesicles ready for exocytosis. These proteins include the ABC family, the MATE family, the major facilitator superfamily (MFS), and the ALMT (aluminium-activate malate transporter) family of transport proteins [118].

ABC proteins mediated transport system:

ABC proteins comprise a large and ancient family of proteins found in all phyla. They drive the transport of various substrates (metabolic products, ions, lipids and xenobiotics) using the energy of ATP hydrolysis. Hence, these are primarily active transporters that have the potential to transport substrates against their electrochemical gradient. In eukaryotic cells, ABCs export substrates from the cytosol to the apoplast through the plasma membrane or from the cytosol to other internal organelles such as the vacuole. There are >100 putative ABCs in the rice and *Arabidopsis* genomes, some of which are involved in the transport of glutathione conjugates [119], chlorophyll catabolites [120], auxins [121,122]. For example, ABCs are thought to export the antifungal diterpene sclareol from the leaves of *Nicotiana glauca* [123] and the isoflavone genistein from soybean roots [124]. Genistein is a signaling molecule that is released from the roots of many legume species to stimulate nodulation. It may also function as a phytoalexin due to its slight antimicrobial activity [125]. Another gene required for powdery mildew resistance in *Arabidopsis* encodes an ABC transporter called PEN3 and it localizes to the plasma membrane, and its recruitment to sites of pathogen attack on the epidermal cells and root hairs is triggered by pathogen-associated molecular patterns such as flagellin and chitin [126]. PEN3 could be releasing antimicrobial compounds, including glucosinolate derivatives into the apoplast adjacent to the appressoria to restrict pathogen invasion. Interestingly, PEN3 (=AtPDR8) also prevents heavy metal toxicity by exporting cadmium ions or cadmium conjugates from root cells [127], suggesting that the same transporter serves different functions in different tissues. Another example related to leaf pathogens is Lr34, one of the most important disease resistance genes in wheat. Lr34 is a full-sized ABC transporter expressed in leaves that is necessary for durable rust resistance in wheat around the world [128]. The function of Lr34

remains unclear, but one model proposes that Lr34 protects infected cells by exporting antimicrobial compounds into the apoplast in response to pathogen attack. Badri et al. [62] identified 25 ABC transporter genes highly expressed in *Arabidopsis* root cells that are likely to be involved in secretion processes. Plants with knockout mutations in these genes were obtained, and the exudates collected from wild-type plants and mutant lines were compared. They found that exudate composition differed between the wild-type and mutant plants and that more than one ABC transporter could be involved in the secretion of a specific phytochemical.

MATE proteins

The MATE genes encode transporters that also export a variety of substrates, including secondary metabolites. The family is found in eukaryotes and prokaryotes [129,130], and certain members in bacteria and mammals are responsible for multidrug resistance. Details of their function are not well understood, MATs appear to function as secondary active transporters using the electrochemical gradient of other ions (sodium or protons) to drive substrate motion. Arabidopsis gene called AtDTX1 encodes a plasma membrane-located protein that facilitates the efflux of plant alkaloids, antibiotics and other toxic compounds from the roots [131]. Other MATE genes in sorghum (SbMATE1), barley (HvAACT1), and Arabidopsis (AtMATE1) confer resistance to aluminum by facilitating aluminum-activated efflux of citrate anions from root tips [132,133]. Citrate released from the roots is thought to chelate the toxic Al³⁺ cations in the apoplast and prevent damage occurring to the rapidly dividing and elongating cells in the apices [134,135]. More recently a MATE protein in rice, named PEZ1, was shown to export phenolic compounds into the xylem [136] and it was speculated that similar proteins might also be facilitating the transport of phenolic compounds into the rhizosphere.

MFS proteins

Release of a special class of secondary metabolites, the phytosiderophores, is an important iron nutrient in grasses [137]. These phytosiderophores chelate iron(III) in the rhizosphere and the entire complex is then transported to root cells by proton-coupled transporters belonging to the oligopeptide transporter family [138]. However, the initial export of these compounds from roots involves the MFS proteins, one of the largest and most diverse. TOM1 expression is induced when iron supply is restricted, and transgenic plants

overexpressing TOM1 show increased release of deoxymugic acid and increased tolerance to reduced iron supply.

Future aspects

The quantities of organic compounds exuding from roots is not large, about only 0.4% of the total carbon photosynthesized, they do exert a very strong influence on the soil microorganisms attraction and may be significant in affecting plant nutrient availability. There is evidence that exudates from the roots of some plants are toxic to roots of neighboring plants and to some germinating seeds. The present review compilation is based data generated from the experiments carried on media based solution-grown plants so that there are still some important questions to be answered about root exudates from soil-grown plants: (i) How much organic material is exuded from roots into soil ? (ii) How far from roots do the compounds diffused from diffusion site ? (iii) What concentration gradients exist ? (iv) What effects do soil properties have upon exudation ? (v) What are the main sites of exudation ? The application of C¹⁴-labeling techniques offers a powerful tool which answer these questions. Use of radioisotopes to investigate the sites of exudation along roots and the role of lateral roots in the exudation process should enable us to determine what part of the soil will be most affected by root exudates.

Conclusion

Based on the current status, rhizosphere microbial community in the soil is closely linked to plant growth, and it can be changed by alteration of root exudate secretion pattern. Studies proved that changing the root exudate patterns by modifying ABC transporter could affect the rhizosphere microbiome levels. However, benefits of PGPR to plants in terms of growth and yield are well-documented and majority of the studies have only emphasized on plant parameters aboveground. But, in fact the interactions of PGPR with the plant starts in the rhizosphere through root exudation followed by attraction, movement and colonization on the rhizoplane, and/or in the endorhizosphere. In rhizosphere, most of the biological activities guided by the root exudates. Chemotaxis is a vital motility property that allows bacteria to move towards the rhizosphere and this process of chemotaxis towards root exudates has been suggested to be the first step of bacterial colonization on root surface. It was concluded that root exudates transported through the various mechanisms for the movement of secondary plant metabolites in roots are probably similar to those used elsewhere in the plant, but much larger quantities of these compounds are released from roots than from leaves. A variety of

transport mechanisms are employed to move bioactive secondary products around root cells and into the rhizosphere, and these fluxes are often highly dependent on the local soil environment and root health. This review has highlighted the possible pathways for the safe transport of secondary metabolites and their interactions with microbial communities especially endophytes in inducing plant growth.

References:

1. Hiltner L. Über neuere Erfahrungen und Probleme auf dem Gebiet der Bodenbakteriologie und unter besonderer Berücksichtigung der Grundung und Brache. Arb Dtsch Landwirsch Ges, 1904;98:59.
2. Hartmann A, Rothballer M, Schmid M, Lorenz H. A pioneer in rhizosphere microbial ecology and soil bacteriology research. Plant Soil, 2008;312:7–14. <https://doi.org/10.1007/s11104-007-9514->
3. Lemanceau P, Corberand T, Gardan L, Latour X, Laguerre G. Effect of two plant species, flax (*Linum usitatissimum* L.) and tomato (*Lycopersicon esculantum* Mill.), on the diversity of soilborne populations of fluorescent Pseudomonads. Applied and Environmental Microbiology, 1995;61:1004–1012.
4. Grayston S, Wang JS, Campbell CD, Edwards AC. Selective influence of plant species on microbial diversity in the rhizosphere. Soil Biology & Biochemistry, 1998;30:369–378.
5. Miethling R, Wieland G, Backhaus H, Tebbe C. Variation of microbial rhizosphere communities in response to crop species, soil origin, and inoculation with *Sinorhizobium meliloti* L33. Microbial Ecology, 2000;41:43–56.
6. Berg G, Smalla K. Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. FEMS Microbiology Ecology, 2009;68:1–13.
7. Mendes R, Kruijt M, De Bruijn I, Dekkers E, Van Der Voort M, Schneider JHM, Piceno YM. Deciphering the rhizosphere microbiome for disease-suppressive bacteria. Science, 2011;332:1097–1100.
8. Berendsen RL, Pieterse CMJ, Bakker PAHM. 2012. The rhizosphere microbiome and plant health. Trends in Plant Science, 2012;17:478–486.

9. Rosenblueth M, Martínez-Romero E. Bacterial endophytes and their interactions with hosts. *Molecular Plant-Microbe Interactions*, 2006;19:827–837.
10. Hardoim PR, van Overbeek LS, van Elsas JD. Properties of bacterial endophytes and their proposed role in plant growth. *Trends in Microbiology*, 2008;16:463–471.
11. Olivares F, Baldani V, Reis V. Occurrence of the endophytic diazotrophs *Herbaspirillum* spp. in roots, stems, and leaves, predominantly of Gramineae. *Biology and Fertility of Soils*, 1996;21:197–200.
12. Weber O, Baldani V, Teixeira K. Isolation and characterization of diazotrophic bacteria from banana and pineapple plants. *Plant and Soil*, 1999;210:103–113.
13. Dong Y, Iniguez AL, Triplett EW. Quantitative assessments of the host range and strain specificity of endophytic colonization by *Klebsiella pneumoniae* 342. *Plant and Soil*, 2003;257:49–59.
14. Mathesius U, Mulders S. Extensive and specific responses of a eukaryote to bacterial quorum-sensing signals. *Proceedings of the National Academy of Sciences, USA*, 2003;100:1444–1449.
15. Rudrappa T, Czymmek K, Paré PW, Bais HP. Root-secreted malic acid recruits beneficial soil bacteria. *Plant Physiology*, 2008;148:1547–1556.
16. Kent Peters N, Long SR. Alfalfa root exudates and compounds which promote or inhibit induction of *Rhizobium meliloti* nodulation genes. *Plant Physiol*, 1998;88:396–400.
17. Besserer A, Puech-Pagès V, Kiefer P, Gomez-Roldan V, Jauneau A, Roy S, Portais JC, Roux C, Bécard G, Séjalon-Delmas N. Strigolactones stimulate arbuscular mycorrhizal fungi by activating mitochondria. *PLoS Biol*, 2006;4:e226.
18. Bais HP, Fall R, Vivanco JM. Biocontrol of *Bacillus subtilis* against infection of *Arabidopsis* roots by *Pseudomonas syringae* is facilitated by biofilm formation and surfactin production. *Plant Physiol*, 2004;134:307–319.
19. Cavaglieri L, Orlando J, Rodriguez MI, Chulze S, Etcheverry M. Biocontrol of *Bacillus subtilis* against *Fusarium verticillioides* in vitro and at the maize root level. *Res Microbiol*, 2005;156:748–754.
20. Shidore T, Dinse T, Öhrlein T, Becker A, Reinhold-Hurek B. Transcriptomic analysis of responses to exudates reveal genes required for rhizosphere competence of the endophyte *Azoarcus* sp. strain BH72. *Environ. Microbiol*, 2012;14:2775–2787.
21. Sandra de W, Hans V, Ine, HMM, Irene K, Nico H, Guido VB, Jos V, René De M, Ben JLL. Flagella-Driven Chemotaxis Towards Exudate Components Is an Important Trait

- for Tomato Root Colonization by *Pseudomonas fluorescens*. *Mol Plant Microbe Interact*, 2002;15(11):1173-1180.
22. Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM. *The role of root exudates in rhizosphere interactions with plants and other organisms. Annual review of Plant Biology*, 2006;57:233-266.
 23. Lugtenberg BJ, Dekkers L, Bloemberg GV. Molecular determinants of rhizosphere colonization by *Pseudomonas*. *Annu Rev Phytopathol*, 2001;39:461-90.
 24. **Bacilio-Jiménez M, Aguilar-Flores S, Ventura-Zapata E, Pérez-Campos ES, Bouquelet S, Zenteno E.** Chemical characterization of root exudates from rice (*Oryza sativa*) and their effects on the chemotactic response of endophytic bacteria. *Plant Soil*, 2003;249:271-277.
 25. Coombs JT, Franco CMM. Visualization of an endophytic *Streptomyces* species in wheat seed. *Appl Environ Microbiol*, 2003b;69:4260-4262.
 26. Hallmann J, Quadt-Hallmann A, Mahaffee WF, Kloepper JW. Bacterial endophytes in agricultural crops. *Can. J. Microbiol*, 1997;43:895-914.
 27. Rosenblueth M, Martinez Romero E. *Rhizobium etli* maize populations and their competitiveness for root colonization. *Arch. Microbiol*, 2004;181:337-344.
 28. Bais HP, Fall R, Vivanco JM. Biocontrol of *Bacillus subtilis* against infection of *Arabidopsis* roots by *Pseudomonas syringae* is facilitated by biofilm formation and surfactin production. *Plant Physiol*, 2004a;134:307.
 29. Bais HP, Park SW, Weir TL, Callaway RM, Vivanco JM. How plants communicate using the underground information superhighway. *Trends Plant Sci*, 2004b;9:26.
 30. Walker TS, Bais HP, Grotewold E, Vivanco JM. Root exudation and rhizosphere biology. *Plant Physiol*, 2003;1320:44.
 31. Weir TL, Park SW, Vivanco JM. Biochemical and physiological mechanisms mediated by allelochemicals. *Curr Opin Plant Biol*, 2004;7:472.
 32. Uren NC. 2000. Types, amounts and possible functions of compounds released into the rhizosphere by soil grown plants. In: Pinton R, Varanini Z, Nannipieri P (eds) *The rhizosphere: biochemistry and organic substances at the soil interface*. Dekker, New York, pp 19–40.
 33. Cai Z, Kastell A, Knorr D, Smetanska I. Exudation: An expanding technique for continuous production and release of secondary metabolites from plant cell suspension and hairy root cultures. *Plant cell reports*, 2012;3:461-477.

34. Carvalhais LC, Dennis PG, Fan B, Fedoseyenko D, Kierul K, Becker A, Von Wiren N, Borriss R. Linking plant nutritional status to plant-microbe interactions. *PLoS One*, 2013;8:e68555.
35. Cai T, Cai W, Zhang J, Zheng H, Tsou AM, Xiao L, Zhong Z, Zhu J. Host legume-exuded antimetabolites optimize the symbiotic rhizosphere. *Mol Microbiol*, 2009;73:507-517.
36. Guttman D, McHardy AC, Schulze-Lefert P. Microbial genome-enabled insights into plant-microorganism interactions. *Nat Rev Genet*, 2014;15:797-813.
37. Bergsma-Vlami M, Prins ME, Raaijmakers JM. Influence of plant species on population dynamics, genotypic diversity and antibiotic production in the rhizosphere by indigenous *Pseudomonas* spp. *FEMS Microbiol Ecol*, 2005;52:59-69.
38. Aira M, Gómez-Brandón M, Lazcano C, Baath E, Domínguez J. Plant genotype strongly modifies the structure and growth of maize rhizosphere microbial communities. *Soil Biol Biochem*, 2010;42:2276-2281.
39. Ramachandran VK, East AK, Karunakaran R, Downie JA, Poole PS. Adaptation of *Rhizobium leguminosarum* to pea, alfalfa and sugar beet rhizospheres investigated by comparative transcriptomics. *Genome Biol*, 2011;12:R106.
40. Moulin L, Munive A, Dreyfus B, Boivin-Masson C. Nodulation of legumes by members of the beta-subclass of Proteobacteria. *Nature*, 2001;411:948.
41. Schardl CL, Leuchtman A, Spiering MJ. Symbioses of grasses with seedborne fungal endophytes. *Annu Rev Plant Biol*, 2004;55:315.
42. Gray EJ, Smith DL. Intracellular and extracellular PGPR: commonalities and distinctions in the plant-bacterium signaling processes. *Soil Biol Biochem*, 2005;37:395.
43. Bais HP, Broeckling CD, Vivanco JM. 2008. Root Exudates Modulate Plant—Microbe Interactions in the Rhizosphere. In: Karlovsky P. (eds) *Secondary Metabolites in Soil Ecology. Soil Biology*, 14. Springer, Berlin, Heidelberg, pp. 241-252.
44. Le´vy J, Bres C, Geurts R, Chalhoub B, Kulikova O, Duc G, Journet EP, Ane´ JM, Lauber E, Bisseling T, De´narie´ J, Rosenberg C, Debelle´ F. A putative Ca²⁺ and calmodulin dependent protein kinase required for bacterial and fungal symbioses. *Science*, 2004;303:1361–1364.
45. Nagahashi G, Douds DD. Rapid and sensitive bioassay to study signals between root exudates and arbuscular mycorrhizal fungi. *Biotechnol Tech*, 1999;13:893–897.

46. Tamasloukht MB, Sejalon-Delmas N, Kluever A, Jauneau A, Roux C, Becard G, Franken P. Root factors induce mitochondrial-related gene expression and fungal respiration during the developmental switch from asymbiosis to pre-symbiosis in the arbuscular mycorrhizal fungus *Gigaspora rosea*. *Plant Physiol*, 2003;131:1468–1478.
47. Akiyama K, Ki M, Hayashi H. Plant sesquiterpenes induce hyphal branching in arbuscular mycorrhizal fungi. *Nature*, 2005;435:824–827.
48. Harrison MJ. Signaling in the arbuscular mycorrhizal symbiosis. *Annu Rev Microbiol*, 2005;59:19–42.
49. Karandashov V, Bucher M. Symbiotic phosphate transport in arbuscular mycorrhizas. *Trends Plant Sci*, 2005;10:22–29.
50. Paszkowski U. Mutualism and parasitism: the yin and yang of plant symbioses. *Curr Opin Plant Biol*, 2006;9:364–370.
51. Gherbi H, Markmann K, Svistoonoff S, Estevan J, Autran D, Giczey G, Auguy F, Pe´ret B, Laplaze L, Franche C, Parniske M, Bogusz D. SymRK defines a common genetic basis for plant root endosymbioses with arbuscular mycorrhiza fungi, rhizobia, and Frankia bacteria. *Proc Natl Acad Sci USA*, 2008;105:4928–4932.
52. Lugtenberg B, Kamilova F. Plant-growth-promoting rhizobacteria. *Annu. Rev. Microbiol*, 2009;63:541–556.
53. Bodhankar S, Grover M, Hemanth S, Reddy G, Rasul S, Yadav SK, Desi S, Mallappa M, Mandapaka M, Srinivasarao C. Maize seed endophytic bacteria: dominance of antagonistic, lytic enzyme-producing *Bacillus* spp. *3 Biotech*, 2017;7(4):232.
54. Werra P, Huser A, Tabacchi R, Keel C, Maurhofer M. Plant- and Microbe-Derived Compounds Affect the Expression of Genes Encoding Antifungal Compounds in a Pseudomonad with Biocontrol Activity. *Applied and Environmental Microbiology*, 2011;77(8):2807–2812.
55. Cazorla FM, Mercado-Blanco J. Biological control of tree and woody plant diseases: an impossible task? *Bio Control*, 2016;61(3):233-242.
56. Aziz NH, El-Fouly El, El-Essawy AA, Khalaf MA. Influence of bean seedling root exudates on the rhizosphere colonization by *Trichoderma lignorum* for the control of *Rhizoctonia solani*. *Bot. Bull. Acad. Sin.*, 1997;38: 33-39.
57. Yi Y, Jong A, Elrike Frenzel E, Kuipers OP. Comparative transcriptomics of *Bacillus mycooides* Strains in Response to Potato-Root Exudates Reveals Different Genetic Adaptation of Endophytic and Soil Isolates. *Front. Microbiol*, 2017;8:1487.

58. Schreiter S, Babin D, Kornelia S, Rita G. Rhizosphere Competence and Biocontrol Effect of *Pseudomonas* sp. RU47 Independent from Plant Species and Soil Type at the Field Scale. *Front. Microbiol*, 2018;9:97.
59. Wang N, Wang L, Zhu K, Hou S, Chen L, Mi D, Gui Y, Qi Y, Jiang C, Guo J. Plant Root Exudates Are Involved in *Bacillus cereus* AR156 Mediated Biocontrol Against *Ralstonia solanacearum*. *Front. Microbiol*, 2019;10:98.
60. Badri DV, Vivanco JM. Regulation and function of root exudates. *Plant Cell Environ*, 2009;32:666–681.
61. Weston LA, Ryan PR, Watt M. Mechanisms for cellular transport and release of allelochemicals from plant roots into the rhizosphere. *J. Exp. Bot.*, 2012;63:3445–3454. doi:10.1093/jxb/ers054. PMID:22378954.
62. Badri DV, Loyola-Vargas VM, Broeckling CD, De-la-Peña C, Jasinski M, Santelia D, Martinoia E, Sumner LW, Banta LM, Stermitz F, Vivanco JM. Altered profile of secondary metabolites in the root exudates of *Arabidopsis* ATP-binding cassette transporter mutants. *Plant Physiol*, 2008;146(2):762–771. doi:10.1104/pp.107.109587. PMID:18065561.
63. Badri DV, Quintana N, El Kassis EG, Kim HK, Choi YH, Sugiyama A, Verpoorte R, Martinoia E, Manter DK, Vivanco JM. An ABC transporter mutation alters root exudation of phytochemicals that provoke an overhaul of natural soil microbiota. *Plant Physiol*, 2009a;151(4):2006–2017. doi: 10.1104/pp.109.147462. PMID:19854857.
64. Loyola-Vargas V, Broeckling C, Badri D, Vivanco J. Effect of transporters on the secretion of phytochemicals by the roots of *Arabidopsis thaliana*. *Planta*, 2007;225(2): 301–310. doi:10.1007/s00425-006-0349-2. PMID:16868775
65. Sugiyama A, Shitan N, Yazaki K. Signaling from soybean roots to *Rhizobium*: an ATP-binding cassette-type transporter mediates genistein secretion. *Plant Signal. Behav*, 2008;3(1): 38–40. doi:10.4161/psb.3.1.4819. PMID:19704765
66. Yazaki K. Transporters of secondary metabolites. *Curr. Opin. Plant Biol*, 2005;8(3):301–307. doi:10.1016/j.pbi.2005.03.011. PMID:15860427.
67. Reddy VS, Shlykov MA, Castillo R, Sun EI, Saier MH. The major facilitator superfamily (MFS) revisited. *FEBS J*, 2012;279(11):2022–2035. doi:10.1111/j.1742-4658.2012.08588.x. PMID:22458847.

68. Broeckling CD, Broz AK, Bergelson J, Manter DK, Vivanco JM. Root exudates regulate soil fungal community composition and diversity. *Appl. Environ. Microbiol*, 2008;74(3):738–744. doi:10.1128/AEM.02188-07.
69. Doornbos R, Loon L, Bakker PHM. Impact of root exudates and plant defense signaling on bacterial communities in the rhizosphere. A review. *Agron. Sust. Dev*, 2012;32(1):227–243. doi:10.1007/s13593-011-0028-y.
70. Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM. Root exudation of phytochemicals in *Arabidopsis* follows specific patterns that are developmentally programmed and correlate with soil microbial functions. *PloS ONE*, 2013a;8(2):e55731. doi:10.1371/journal.pone.0055731. PMID:23383346.
71. Chaparro JM, Badri DV, Vivanco JM. Rhizosphere microbiome assemblage is affected by plant development. *ISME J*, 2013b;8(4):790–803. doi:10.1038/ismej.2013.196.
72. Micallef SA, Channer S, Shiaris MP, Colón-Carmona A. Plant age and genotype impact the progression of bacterial community succession in the *Arabidopsis* rhizosphere. *Plant Signal. Behav*, 2009a;4(8):777–780. doi:10.4161/psb.4.8.9229. PMID:19820328.
73. Micallef SA, Shiaris MP, Colón-Carmona A. Influence of *Arabidopsis thaliana* accessions on rhizobacterial communities and natural variation in root exudates. *J. Exp. Bot*, 2009b;60(6):1729–1742. doi:10.1093/jxb/erp053. PMID:19342429.
74. Mercado-Blanco J, Bakker P. Interactions between plants and beneficial *Pseudomonas* spp.: exploiting bacterial traits for crop protection. *Antonie van Leeuwenhoek*, 2007;92(4):367–389. doi:10.1007/s10482-007-9167-1.
75. Raaijmakers J, Paulitz T, Steinberg C, Alabouvette C, Moënne-Loccoz Y. The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. *Plant Soil*, 2009;321(1–2): 341–361. doi:10.1007/s11104-008-9568-6.
76. Newton AC, Fitt BDL, Atkins SD, Walters DR, Daniell TJ. Pathogenesis, parasitism and mutualism in the trophic space of microbe– plant interactions. *Trends Microbiol*, 2010;18(8):365–373. doi:10.1016/j.tim.2010.06.002. PMID:20598545.
77. Davidson IA, Robson MJ. Effect of contrasting patterns of nitrate application on the nitrate uptake, N₂-fixation, nodulation and growth of white clover. *Ann. Bot*, 1986;57(3):331–338.

78. Zahran HH. Rhizobium–legume symbiosis and nitrogen fixation under severe conditions and in an arid climate. *Microbiol. Mol. Biol. Rev.* 1999;63(4):968–989. PMID:10585971.
79. Coronado C, Zuanazzi J, Sallaud C, Quirion JC, Esnault R, Husson HP, Kondorosi A, Ratet P. Alfalfa root flavonoid production is nitrogen regulated. *Plant Physiol*, 1995;108(2): 533–542. doi:10.1104/pp.108.2.533. PMID:12228491.
80. Zhang J, Subramanian S, Stacey G, Yu O. Flavones and flavonols play distinct critical roles during nodulation of *Medicago truncatula* by *Sinorhizobium meliloti*. *Plant J*, 2009;57(1):171–183. doi:10.1111/j.1365-313X.2008.03676.x. PMID:18786000.
81. Jones DL, Hodge A, Kuzyakov Y. Plant and mycorrhizal regulation of rhizodeposition. *New Phytol*, 2004;163:459e480.
82. Vranova V, Rejsek K, Formanek P. Aliphatic, cyclic, and aromatic organic acids, vitamins, and carbohydrates in soil. *Sci. World J*, 2013;15:524239, <http://dx.doi.org/10.1155/2013/524239>.
83. Neumann G, Romheld V. 2000. The release of root exudates as affected by the plant's physiological status. In *The Rhizosphere, Biochemistry and Organic Substances at the Soil–Plant Interface* (eds. R.Pinton, Z.Varanini & P.Nannipieri), pp. 41–93. Marcel Dekker, New York, NY, USA.
84. Sanders D, Bethke P. 2000. Membrane transport. In *Biochemistry and Molecular Biology of Plants* (eds. B.B. Buchanan, W. Gruisham & R.L. Jones), pp. 110–158. ASPP, Rockville, MD, USA.
85. Guern J, Renaudin JP, Brown SC. 1987. The compartmentation of secondary metabolites in plant cell cultures. In *Cell Culture and Somatic Cell Genetics of Plants* (eds. F.Constabel & I.K.Vasil), pp. 43–76. Academic Press, San Diego, CA, USA.
86. Marschner H. 1995. *Mineral Nutrition of Higher Plants*. Academic Press, London, UK.
87. Samuel AL, Fernando M, Glass ADM. Immunofluorescent localization of plasma membrane H⁺-ATPase in barley roots and effects of K nutrition. *Plant Physiology*, 1992;99:1509-1514.
88. Jones DL, Darrah PR. Influx and efflux of organic acids across the soil–root interface of *Zea mays* L. and its implications in rhizosphere C flow. *Plant and Soil*, 1995;173:103–109.

89. Ryan PR, Delhaize E, Randall PJ. Characterization of Al-stimulated efflux of malate from the apices of Al-tolerant wheat roots. *Planta*, 1995;196:103–110.
90. Zheng SJ, Ma JF, Matsumoto H. High aluminum resistance in buckwheat. Aluminum-induced specific secretion of oxalic acid from root tips. *Plant Physiology*, 1998;117:745–751.
91. Neumann G, Massonneau A, Martinoia E, Romheld V. Physiological adaptation to phosphorus deficiency during proteoid root development in white lupin. *Planta*, 1999;208:373–382.
92. Sakaguchi T, Nishizawa NK, Nakanishi H, Yoshimura E. The role of potassium in the secretion of meginic acids family phytosiderophores from iron-deficient barley roots. *Plant and Soil*, 1999;215:221–227.
93. Williams LE, Pittman JK, Hall JL. Emerging mechanisms for heavy metal transport in plants. *Biochimica et Biophysica Acta*, 2000;1465:104–126.
94. Hussain D, Haydon MJ, Wang Y, Wong E, Sherson SM, Young J, Camakaris J, Harper JF, Cobbett CS. P-type ATPase heavy metal transporters with roles in essential zinc homeostasis in *Arabidopsis*. *The Plant Cell*, 2004;16:1327–1339.
95. Colangelo EP, Guerinot ML. Put the metal to the petal: metal uptake and transport throughout plants. *Current Opinion in Plant Biology*, 2006;9:322–330.
96. Hirner A, Ladwig F, Stransky H, Okumoto S, Keinath M, Harms A, Frommer WB, Koch W. *Arabidopsis* LHT1 is a high-affinity transporter for cellular amino acid uptake in both root epidermis and leaf mesophyll. *The Plant Cell*, 2006;18:1931–1946.
97. Hoekenga OA, Maron LG, Piñeros MA, Cançado GMA, Shaff J, Kobayashi Y, Ryan PR, Dong B, Delhaize E, Sasaki T, Matsumoto H, Yamamoto Y, Koyama H, Kochian LV. AtALMT1, which encodes a malate transporter, is identified as one of several genes critical for aluminum tolerance in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America*, 2006;103:9749–9750.
98. Grabov A. Plant KT/KUP/HAK potassium transporters: a single family – multiple functions. *Annals of Botany*, 2007;99:1035–1041.
99. Lee YH, Foster J, Chen J, Voll LM, Weber AP, Tegeder M. AAP1 transports uncharged amino acids into roots of *Arabidopsis*. *The Plant Journal*, 2007;50:305–319.

100. Svennerstam H, Ganeteg U, Bellini C, Nasholm T. Comprehensive screening of *Arabidopsis* mutants suggests the lysine histidine transporter 1 to be involved in plant uptake of amino acids. *Plant Physiology*, 2007;143:1853–1860.
101. Hayden MJ, Cobbett CS. Transporters of ligands for essential metal ions in plants. *New Phytologist*, 2007;174:499–506.
102. Curie C, Panaviene Z, Loulergue C, Dellaporte SL, Briat JF, Walker EL. Maize yellow stripe1 encodes a membrane protein directly involved in Fe(III) uptake. *Nature*, 2001;409:346–349.
103. Curie C, Briat JF. Iron transport and signaling in plants. *Annual Review of Plant Biology*, 2003;54:183–206.
104. Ma JF, Yamaji N, Mitani N, Tamai K, Konishi S, Fujiwara T, Katsuhara M, Yano M. An efflux transporter of silicon in rice. *Nature*, 2007;448:209–212.
105. Ma JF, Yamaji N. Functions and transport of silicon in plants. *Cellular and Molecular Life Sciences*, 2008;65:3049–3057.
106. Kobayashi Y, Hoekenga OA, Itoh H, Nakashima M, Saito S, Shaff JE, Maron LG, Pineros MA, Kochian LV, Koyama H. Characterization of AtALMT1 expression in aluminum-inducible malate release and its role for rhizotoxic stress tolerance in *Arabidopsis*. *Plant Physiology*, 2007;145:843–852.
107. Klepek YS, Geiger D, Stadler R, Klebl F, Landour-Arsivaud L, Lemoine R, Hedrich R, Sauer N. *Arabidopsis* POLYOL TRANSPORTERS, a new member of the monosaccharide transporter-like superfamily, mediates H⁺-symport of numerous substrates including myo-inositol, glycerol and ribose. *The Plant Cell*, 2005;17:204–218.
108. Buttner M. The monosaccharide transporter (-like) gene family in *Arabidopsis*. *FEBS Letters*, 2007;581:2318–2324.
109. Battey NH, Blackbourn HD. The control of exocytosis in plant cells. *New Phytologist*, 1993;125:307–308.
110. Field B, Jordan F, Osbourn A. First encounters – deployment of defence-related natural products by plants. *New Phytologist*, 2006;172:193–207.
111. Lin Y, Irani NG, Grotewold E. Sub-cellular trafficking of phytochemicals explored using auto-fluorescent compounds in maize cells. *BMC Plant Biology*, 2003;3:e10.
112. Grotewold E. The challenges of moving chemicals within and out of cells: insights into the transport of plant natural products. *Planta*, 2004;219:906–909.

113. Snyder BA, Nicholson RL. Synthesis of phytoalexins in sorghum as a site-specific response to fungal ingress. *Science*, 1990;248:1637–1639.
114. Snyder BA, Leite B, Hipskind J, Butler LG, Nicholson RL. Accumulation of sorghum phytoalexins induced by *Colletotrichum graminicola* at the infection site. *Physiological and Molecular Plant Pathology*, 1991;39:463–470.
115. Tabata M. The mechanism of shikonin biosynthesis in cell *Lithospermum* cultures. *Plant Tissue Culture Letters*, 1996;13:117–125.
116. Yazaki K, Kuniyama M, Fujisaki T, Sato F. Geranyl diphosphate: 4-hydroxybenzoate geranyl transferase from *Lithospermum erythrorhizon*. *Journal of Biological Chemistry*, 2002;277:6240–6246.
117. Yazaki K, Matsuoka H, Shimomura K, Bechthold A, Sato F. A novel dark-inducible protein LeDI-2 and its involvement in root-specific secondary metabolism in *Lithospermum erythrorhizon*. *Plant Physiology*, 2001;125:1831–1841.
118. Leslie WA, Peter R, Ryan Michelle W. Mechanisms for cellular transport and release of allelochemicals from plant roots into the rhizosphere. *Journal of Experimental Botany*, 2012;63(9):3445–3454, <https://doi.org/10.1093/jxb/ers054>
119. Martinoia E, Klein M, Geisler M, Bovet L, Forestier C, Kolukisaoglu U, Muller R, Schulz B. Multifunctionality of plant ABC transporters – more than just detoxifiers. *Planta*, 2002;214:345-355.
120. Tommasini R, Vogt E, Fromenteau M, Hörtensteiner S, Matile P, Amrhein N, Martinoia E. An ABC-transporter of *Arabidopsis thaliana* has both glutathione-conjugate and chlorophyll catabolite transport activity. *The Plant Journal*, 1998;13:773-780.
121. Noh B, Murphy AS, Spalding EP. Multidrug resistance-like genes of *Arabidopsis* required for auxin transport and auxin-mediated development. *Plant Cell*, 2001;13:2441-2454.
122. Geisler M, Murphy A. The ABC of auxin transport: the role of P-glycoproteins in plant development. *FEBS Letters*, 2006;580:1094-1102.
123. Jasinski M, Stukkens Y, Degand H, Purnelle B, Marchand Brynaert J, Boutry MA. Plant plasma membrane ATP binding cassette-type transporter is involved in antifungal terpenoid secretion. *Plant Cell*, 2001;13:1095-1107.

124. Sugiyama A, Shitan N, Yazaki K. Involvement of a soybean ATP-binding cassette-type transporter in the secretion of genistein, a signal flavonoid in legume–Rhizobium symbiosis. *Plant Physiology*, 2007;144:2000-2008.
125. Geibel M. Sensitivity of the fungus *Cytospora personii* to the flavonoids of *Prunus cerasus*. *Phytochemistry*, 1994;38:599-601.
126. Stein M, Dittgen J, Sanchez-Rodriguez C, Hou BH, Molina A, Schulze-Lefert P, Lipka V, Shauna S. Arabidopsis PEN3/PDR8, an ATP binding cassette transporter, contributes to nonhost resistance to inappropriate pathogens that enter by direct penetration. *Plant Cell*, 2006;18:731-746.
127. Kim DY, Bovet L, Maeshima M, Martinoia E, Lee Y. The ABC transporter AtPDR8 is a cadmium extrusion pump conferring heavy metal resistance. *The Plant Journal*, 2007;50:207-218.
128. Krattinger SG, Lagudah ES, Spielmeier W, Singh RP, Huerta Espino J, McFadden H, Bossolini E, Selter LL, Keller B. Putative ABC transporter confers durable resistance to multiple fungal pathogens in wheat. *Science*, 2009;323:1360-1363.
129. Hvorup RN, Winnen B, Chang AB, Jiang Y, Zhou XF, Saier MH. The multidrug/oligosaccharidyl-lipid/polysaccharide (MOP) exporter superfamily. *European Journal of Biochemistry*, 2003;270:799-813.
130. Magalhaes JV. How a microbial drug transporter became essential for crop cultivation on acid soils: aluminium tolerance conferred by the multidrug and toxic compound extrusion (MATE) family. *Annals of Botany*, 2010;106:199-203.
131. Li L, He Z, Pandey GK, Tsuchiya T, Luan S. Functional cloning and characterization of a plant efflux carrier for multidrug and heavy metal detoxification. *Journal of Biological Chemistry*, 2002;277:5360-5368.
132. Furukawa J, Yamaji N, Wang H, Mitani N, Murata Y, Sato K, Katsuhara M, Takeda K, Ma JF. An aluminum-activated citrate transporter in barley. *Plant and Cell Physiology*, 2007;48:1081-1091.
133. Liu JP, Magalhaes JV, Shaff J, Kochian LV. Aluminum-activated citrate and malate transporters from the MATE and ALMT families function independently to confer Arabidopsis aluminum tolerance. *The Plant Journal*, 2009;57:389-399.
134. Ma JF, Ryan PR, Delhaize E. Aluminium tolerance in plants and the complexing role of organic acids. *Trends in Plant Science*, 2001;6:273-278.

135. Delhaize E, Gruber BD, Ryan PR. The roles of organic anion permeases in aluminium tolerance and mineral nutrition. *FEBS Letters*, 2007;581:2255-2262.
136. Ishimaru Y, Kakei Y, Shimo H, Bashir K, Sato Y, Sato Y, Uozumi N, Nakanishi H, Nishizawa NK. A rice phenolic efflux transporter is essential for solubilizing precipitated apoplasmic iron in the plant stele. *Journal of Biological Chemistry*, 2011;286:24649-24655.
137. Marschner H, Romheld V, Kissel M. Localization of phyto siderophore release and of iron uptake along intact barley roots. *Physiology Plant*, 1987;71:157-162.
138. Kim SA, Guerinot ML. Mining iron: Iron uptake and transport in plants. *FEBS Letters*, 2007;581:2273-2280.

UNDER PEER REVIEW