

Original Research Article

PHENOLOGY OF PLANT SPECIES FOUND IN UGWUEME HYDROCARBON SEEP SITE, ENUGU STATE, NIGERIA

ABSTRACT

This study assessed the phenological events of plant species for flowering, fruiting, shedding of leaves, regeneration/germination, yellowing of leaves and death, every month for three consecutive years from 2018 till 2020, at Ugwueme oil seep site vegetation. Phenological information from the phenological studies of plants can be used to detect patterns associated with climate change, address environmental issues on the timing of plants life cycle, and to predict changes in the habitat and ecosystem at large. The stratified random sampling design was adopted for the study. A sampling zone of 130×60 m² was mapped out for the study. The zone was thereafter split into 60 grid plots, each measuring 13 x 10 m² and 60% of these were selected at random. Data obtained from the study was represented in a phenological diagram. Vegetation phenology showed that flowering and fruiting occur through all the months of the year, although with variations. In March-April, most herbs and woody herbs germinated or regenerated in September/October, fruited in October/November towards the end of the rainy season and became senescent, die or lie dormant in February/March. Given that, the Savanna vegetation of Ugwueme showed variations in the timing of phenological events, however, the diversity of available flora species can be protected to sustain its ecological integrity, especially in this era of emerging climate change disasters and food insecurity, through a longer periods of study like 10 years to understand the relationship between changes of human impact and climate change.

Key words: Plant phenology, Vegetation, Diversity, Food Security, Climate Change, Ecosystem

INTRODUCTION

Phenology is the study of the relationship between seasons (climate), the timing of ecological events such as germination, growth, flowering, fruiting, yellowing of leaves and death, as well as

the causes of their occurrence with respect to biotic and abiotic forces (Schwartz, 2013). It may indicate the manifestation of well-marked seasonal periodicity in the appearance of the vegetation. It is another functional physiognomic method which may be used to supplement information on life-forms (Hamann, 2004). In Nigeria, the Southern and Northern Guinea savanna zones have very well marked seasonal periodicity in the appearance of vegetation. In the dry season, between January and April, after the vegetation has been burnt and when the rainy season is about to begin, the trees generally produce new leaves and flowers.

Phenological studies may be tedious and requires a long period for observation. However, it can provide a useful basis for more detailed investigation (Nwadinigwe, 2013). Data from phenological studies has increasingly been used to detect, explain and predict various patterns related to other forms of global change (Schwartz, 2013), as well as to address environmental issues (Morollato *et al.*, 2016). Jenerette *et al.* (2010) assessed the relationships between the phenology of vegetation and seasonal rainfall in summer and winter in a semi-arid region in southern Arizona, USA. Wessels *et al.* (2011) used remotely sensed vegetation phenology to describe and predict the biomes of South Africa. They found that the phenology and productivity showed a clear relationship with the seasonality of rainfall and mean annual precipitation, respectively.

Phenological studies, monitored year after year will help in assessing the effects of pollution, man-made activities, global warming and climate change on the phenology and productivity of plant species in the ecosystem (Chuine and Régnière, 2017). It helps in monitoring changes in the ecosystem and complexities in their vegetation over time and space for proper rangeland management and land use. There is need to know the time for flowering, fruiting, leaf production and regeneration, for the feeding and pollination by the animals to correspond with the planning and management strategies of savanna rangelands. Plant phenology has been used for assessing the impact of climate change on the seasonality of vegetation (Reed, 2006), and to improve our understanding of the drivers of vegetation growth (Archibald and Scholes, 2007). Phenology invariably affects the health of individual plants, as well as affects the health condition of organisms that rely on plants. However, shifts in plant phenology can adversely affect demography; result in agricultural failures and cause unstable ecosystem (Stucky *et al.*, 2018). At different altitudes and growing seasons, vegetation of any area shows different physiognomic

conditions and each species shows different growth forms such as leafing, flowering, fruiting and death. However, different phenological events are triggered by rainfall, water availability, altitude, temperature and photoperiod (Bhat and Murali, 2001; Hamann, 2004). The phenology of almost all the species studied showed nearly a similar pattern of response to the local climatic conditions. The climate of the study sites has heavy rainfall and high humidity, which is conducive for the growth and development of plants (Okeke and Enoh, 2016). So far, the present work focused on the phenological observations of plant species in Ugwueme oil seep site.

MATERIALS AND METHODS

Description of Study Area

The study area is a crude oil seep site at Ugwueme in Awgu Local Government Area of Enugu State, Nigeria. Ugwueme is situated at longitude 7°25'00" E and 7° 35'00" E and latitude, 6°0'00" N and 6°15' 00" N of geographic co-ordinates. It consists of roughly elevated topography (hills) with steep slopes and could attain an altitude of about 850- 400 meters above the sea level. The natural vegetation of Ugwueme is Guinea Savanna. The area often experiences heavy rainfall during the rainy season with a rainfall record of 1,800 mm (Okeke and Enoh, 2016) annually and high temperature during the dry season with an annual temperature average of 26.6°C. During the first quarter of the year, the temperature normally rises up to 37.7°C and reaches its maximum towards the end of the dry season (Okeke and Enoh, 2016). According to Nigerian Meteorological Agency (NIMET) (2018), the study area experiences a minimum temperature of 25°C and a maximum temperature of 34°C. The vegetation is made up of a mosaic of arable farmlands, tree crop plantations and patches of natural vegetation. Presently, agricultural activities, animal poaching and grazing go on at the site.

The major economic activity of Ugwueme's indigenes is farming. They practice mixed farming where food crops (*Manihot esculenta*, *Dioscorea* sp, *Zea mays*, *Capsicum annuum*, *Solanum lycopersicum* and cash crops (*Musa paradisiaca*, *Elaeis guineensis*, *Anacardium occidentale*) are produced. They also keep local livestock such as goats and sheep for meat and for export.




Fig. 1: Map of Awgu showing Ugwueme

Sampling Design and Techniques

A stratified random sampling field technique by Osuji *et al.*, 2004 was adopted for the study. Sampling zones were erected based on grid system with the aid of measuring tape, ropes and pegs. A sampling zone of 130 x 60 m² was erected around the epicenter; the epicenter is the point of oil seepage (figure 5). The zone was thereafter split into 60 grid plots, each measuring 13 x 10m² and 60% of these (i.e. 36 grid plots) were selected at random. To sample each chosen plot, quadrats were established as follows (a) 4mx 4m for shrubs (b) 1 m x 1m for herbs and grasses. During sampling, plant species inside the quadrat as well as those at the quadrat edge, but with about 75% of their branches inside the quadrat were sampled.



Figure 2 : Morphology of plant species



Figure 3 Habitat and quadrat study of plant

Method of studying phenology

The quadrats were laid at the site of the study. The plant species within the quadrat were identified, marked or tagged. Each plant was monitored every month for flowering, fruiting, shedding of leaves, regeneration of leaves, yellowing of leaves and death. Each phenological process observed or not observed was marked accordingly (Jenerette *et al.*, 2010).

Result

The plant was investigated for the period of three years in the study area for germination, flowering, fruiting, yellowing of leaves, and death. Thereafter, the observed information were represented in phenological diagram (Figure 4).

Germination/leafing: The seedlings and new shoots of annuals and perennials started emerging from ground at the onset of rainy season in the month of March and the number steadily increased up to June. The peak period of germination in herbaceous plants was April (Table 1). The perennial grasses such as *Cymbopogon* and *Paspalum* sprouted from the rhizome/root-stocks. There is an integration of the phenological activities in most herbaceous plant species especially flowering and fruiting which occur in the same month. The herbs that germinated or regenerated with the first rains in March and April were *Aspilia africana*, *Amaranthus hybridus*, *Cynodon dactylon*, *Panicum maximum*, *Chloris pilosa*, *Andropogon tectorum* and *Andropogon gayanus* (Fig. 4).

Flowering: Most herbaceous plants started flowering with increase in temperature during May-June and that continued up to August, and another set of species initiated flowering when the ambient temperature started falling in the later part of September to October (Fig. 4). Furthermore, the trees and shrubs have two peaks of flowering in May and December.

Fruiting: Flowering and fruiting in herbaceous species are simultaneous, since the majority of them need to complete their life cycle within a short span of time. At Ugwueme, most herbs mostly fruited in October towards the end of the rainy season while woody plants mostly fruited in September/October (Fig. 4). Almost all tree species had phenological pattern that synchronized flowering and fruiting in dry months.

Death/Rest: Majority of the annual herbs characteristically complete their life cycles with no resting period. Death or resting phase of most herbs started during Nov to Feb, while the peak was observed in the month of January (Table 1). In the study area, dry season is the most

difficult season for the normal survival of herbs as a result of the severe climatic conditions. On the other hand, perennials develop continuously but most of them undergo a period of rest in which active growth and flowering reduced even though the environment continues to be favourable. Hence, senescence and death are negligible among trees and shrubs when compared with the herbs.

It was noted that plants of the same species growing side by side may not flower and fruit at the same time. One may flower and fruit while the other may not flower and fruit even for the year. Again, in one plant, some branches may not flower and fruit along with other branches. These characteristics were observed in *Daniellia oliveri* and *Mangifera indica*.

The summary of phenology of these species separated into herbs, woody herbs, shrubs and trees is presented in Table 1.

	Plant species	Jan.	Feb.	Mar.	Apr.	Ma y	Jun.	Jul.	Aug.	Set.	Oct.	Nov.	Dec
1	<i>Alchornea cordifolia</i>	Green	Green	Green	Green	Green	Blue	Blue	White	White	White	Green	Green
2	<i>Allophyllus africanum</i>	Green	Green	Green	Green	Blue	Blue	Purple	Purple	Purple	Green	Green	Green
3	<i>Amaranthus hybridus</i>	Red	Red	Green	Green	Green	Blue	Blue	White	White	White	Red	Red
4	<i>Andropogon gayanus</i>	Red	Red	Green	Green	Green	Blue	Blue	Blue	Blue	White	Yellow	Red
5	<i>Andropogon tectorum</i>	Red	Red	Green	Green	Green	Blue	Blue	Blue	Blue	White	White	Yellow
6	<i>Artocarpus altilis</i>	Green	Green	Green	Green	Blue	Blue	White	White	White	White	Green	Green
7	<i>Aspilia Africana</i>	Red	Red	Green	Green	Green	Blue	Blue	Blue	White	White	White	Yellow
8	<i>Axonopus compressus</i>	Red	Red	Green	Green	Green	Blue	Blue	Blue	Blue	White	White	Yellow
9	<i>Capsicum annuum</i>	Yellow	Yellow	Green	Green	Green	Blue	Blue	Blue	White	White	White	Yellow
10	<i>Chamaecrista mimosoides</i>	Red	Red	Red	Green	Green	Blue	Blue	Blue	White	White	White	Yellow
11	<i>Chloria pilosa</i>	Red	Red	Green	Green	Green	Blue	Blue	Blue	Blue	White	White	Yellow
12	<i>Chromolaena odorata</i>	Red	Red	Green	Green	Green	Blue	Blue	Green	Blue	Blue	Purple	Purple
13	<i>Colocasia antiquorum</i>	Red	Red	Red	Green	Green	Blue	Blue	White	White	White	Yellow	Red
14	<i>Conoclinium coelestinum</i>	Red	Red	Red	Green	Green	Blue	Blue	Purple	White	White	White	Yellow
15	<i>Costus afer</i>	Red	Red	Green	Green	Green	Blue	Blue	White	White	White	Yellow	Yellow
16	<i>Cymbopogon citrates</i>	Red	Red	Green	Green	Green	Blue	Blue	Blue	Blue	White	White	Yellow
17	<i>Cynodon dactylon</i>	Red	Red	Green	Green	Green	Blue	Blue	Blue	Blue	White	White	Yellow
18	<i>Daniellia oliveri</i>	Blue	Blue	White	White	White	Green	Green	Green	Green	Green	Green	Blue
19	<i>Dialium guineense</i>	Purple	Purple	Purple	Green	Green	Green	Green	Green	Green	Blue	Blue	Purple
20	<i>Dioscorea alata</i>	Red	Red	Green	Green	Green	Blue	Blue	Purple	Purple	Purple	Purple	Yellow
21	<i>Elaeis guineense</i>	Purple	Purple	Purple	Purple	Green	Green	Green	Green	Green	Blue	Blue	Blue
22	<i>Erythrophleum suaveolens</i>	Green	Green	Green	Blue	Blue	Blue	White	White	White	White	Green	Green

		Green	Green	Green	Blue	Blue	Blue					Green	Green
23	<i>Euphobia thymifolla</i>	Blue					Green	Green	Green	Green	Green	Green	Blue
24	<i>Ficus sur</i>	Green	Green	Green	Blue	Blue	Blue				Green	Green	Green
25	<i>Holarrhena floribunda</i>	Blue	Blue	Purple	Purple	Purple	Green	Green	Green	Green	Green	Blue	Blue
26	<i>Hyparrhenia barteri</i>	Red	Red	Green	Green	Green	Blue	Blue					Yellow
27	<i>Irvingia gabonensis</i>	Green	Green	Blue	Blue			Green	Green	Blue			Green
28	<i>Lecaniodiscus cupanioides</i>	Green	Green	Green	Green	Green	Blue	Blue				Green	Green
29	<i>Lonchocarpus cyanescens</i>	Green	Green	Green	Blue	Blue	Purple	Purple	Purple				Green
30	<i>Mangifera indica</i>	Blue	Blue	Purple	Purple	Purple	Green	Green	Green	Green	Green	Blue	Blue
31	<i>Manihot esculenta</i>	Yellow	Yellow	Green	Green	Green							Yellow
32	<i>Margaritaria citrates</i>	Green	Green	Green	Green	Green	Blue	Blue				Green	Green
33	<i>Millettia aboensis</i>	Blue	Purple	Purple			Green	Green	Green	Green	Green	Blue	Blue
34	<i>Murraya koenigii</i>	Green	Green	Green	Green	Green	Blue	Blue	Purple	Purple	Purple	Green	Green
35	<i>Sarcocephalus latifolius</i>	Green	Green	Green	Green	Green	Blue	Blue				Green	Green
36	<i>Ocimum basilicum</i>	Green	Green	Green	Green	Green	Blue	Blue	Blue				Green
37	<i>Olax subscorpioidea</i>	Green	Green	Green	Green	Green	Blue	Blue	Blue				Green
38	<i>Panicum maximum</i>	Red	Red	Green	Green	Green	Green	Green	Blue	Blue			Yellow
39	<i>Parkia biglobosa</i>	Blue	Blue	Purple	Purple			Green	Green	Green	Green	Green	Blue
40	<i>Paspalum veginatum</i>	Red	Red	Red	Green	Green	Green	Green	Green	Blue			Red
41	<i>Phyllanthus amarus</i>	Red	Red	Red	Green	Green						Yellow	Yellow
42	<i>Ricinus communis</i>	Green	Green	Green	Green	Green	Blue	Blue	Blue	Purple	Purple	Purple	Green
43	<i>Sida acuta</i>	Red	Red	Green	Green	Green	Green	Green	Blue	Blue			Yellow
44	<i>Sida cordifolia</i>	Red	Red	Green	Green	Green	Green	Green	Blue	Blue	Blue		Yellow
45	<i>Talinum fruticosum</i>	Red	Red	Green	Green	Green	Green	Green	Purple	Purple			
46	<i>Tridax procumbens</i>	Red	Red	Green	Green	Green	Green	Green	Blue	Blue	Purple	Purple	Yellow
47	<i>Vernonia amygdalina</i>	Yellow	Yellow	Green	Green	Green							Yellow
48	<i>Vitex doniana</i>	Green	Green	Green	Green	Green			Blue	Blue			Green
49	<i>Zea mays</i>	Red	Green	Green	Green	Green					Red	Red	Red
	Months	Jan.	Feb.	Mar.	Apr	Ma y	Jun.	Jul.	Aug.	Sept	Oct.	Nov.	Dec.

Figure 4: Phenological diagram of plant species found at Ugueme sites.

Key:

	Leave emergence
	Flowering
	Fruiting

	Yellowing of leaves
	Death
	Flowering/fruiting

Table 2: Summary of the phenology of species at Ugwueme vegetation (no of species)

Phenology of herbs and woody herbs	Jan	Feb	mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec.
Germination	0	0	1	21	20	16	0	0	0	0	0	0
Flowering	0	0	0	1	1	3	4	5	7	17	2	1
Fruiting	1	0	0	1	2	1	2	10	5	16	7	4
Death	20	20	19	1	0	0	0	0	0	0	5	12
Phenology of trees and shrubs												
Flowering	7	5	6	5	9	7	5	0	4	4	3	10
Fruiting	2	0	2	7	6	7	9	8	10	14	5	3
Death	1	1	1	0	0	0	0	0	0	0	0	0

Discussion

Phenology is a periodic phenomenon in plants. The climate of the study sites has heavy rainfall and high humidity, which is conducive for the growth and development of plants (Okeke and Enoh, 2016).

The results of the present investigation showed that flowering and fruiting occur throughout all the months of the year, although with variations. This is in agreement with the observation of Marqucus *et al.* (2004) who noted that floral activities in the subtropics occur throughout the year; in contrast to the temperate areas where low temperatures make it impossible for floral activities to take place all the year round. In March-April, most herbs and woody herbs germinated or regenerated; fruited in October/November towards the end of the rainy season and became senescent, die or lie dormant in January/February. These are mostly annual species and

perennial with an annual ephemeral phase. A typical example of the latter is *Chromolaena odorata*, whose meristem is seasonally activated to produce an annual growth in the rainy season, only to have its aerial parts die off in the dry season. These herbaceous species have a short cycle (5-7 months) from germination to flowering period. Their roots are normally shallow. They utilize the shallow soils and their competitive ability is quite low (Nwadinigwe, 2002). Moreover, the results of the study showed that the majority of the herbaceous species germinate in April when the rainfall is moderate at the beginning of the rainy season. This agrees with Baudena *et al.* (2015) who observed that germination and growth in the savanna regions occur in the period of early rains immediately after fruiting and dispersal and after the fires might swept the herbage.

In most of the trees and shrubs, leafing and regeneration are more uniformly distributed throughout the year than the herbaceous species. Even in the dry season (December to March) trees like *Daniellia oliveri*, *Nauclea latifolia*, *Lacaniodiscus cupanioides*, and *Olax subscorpioidea* produce leaves when most herbaceous species are turning brown and dying off.

In the present investigation, the results showed that trees and shrubs have peak of flowering in December, fruited mostly in September and October. Almost all trees had phenological pattern which synchronized flowering and fruiting in dry season. This agrees with the observation of Nwadinigwe (2002) who reported that the dry season is the time for flowering and fruiting for many savanna species so that fruits are ready for dispersal after the fires and before the rains. In woody tree species, the ripening of fruits began in late part of rainy season and continued up to end of cool and dry period which is due to differences in time taken for fruit maturation.

Senescence and death are negligible among trees and shrubs when compared to the herbs. These trees and shrubs are mostly perennial species with seasonal or continuous growth and they possess deep roots utilizing the deeper soils. These investigations are in agreement with the work of Nwadinigwe (2002), who reported that perennials develop continuously but most of them undergo a period of rest in which active growth and flowering are reduced even though the environment continues to be favourable.

However, from a close observation the crude oil seepage made remarkable differences in vegetation phenology of the study sites. The vegetation at the oil-seep site had few species number, delay in regeneration/germination, flowering and fruiting while the control site had the

highest species diversity. Tane and Albert (2015) reported reduction of soil nutrient content in oil polluted site and Johnson *et al.* (2019) who said that oil contamination declines plant performance. In conclusion, plants perform various vegetative and reproductive functions throughout the year in order to survive in the ecosystem. This work provided the phenological timing of different plant species and this will serve as a guide to planning for specific land uses.

REFERENCES

- Archibald, S. and Scholes, R.J. (2007). Leaf green-up in a semi-arid African savanna-separating tree and grass responses to environment cues. *Journal of vegetation science*, 18: 583-594
- Baudena, M., Decker, S. C., van-Bodegom, P. M., Cuesta, B., Higgins, S. I., Lehsten, V., Reich, C. H., Rietkerk, M., Scheiter, S., Yin, Z., Zavala, M. A. , and Brovkin, V. (2015). Forests, savannas, and grasslands: bridging the knowledge gap between ecology and Dynamic Global Vegetation Models, *Biogeosciences*, 12: 1833–1848,
- Bhat, D. M. and Murali, K. S. (2001). Phenology of understory species of tropical moist forest of Western Ghats region of Uttara Kannada district in South India, *Current Science*, 81(7): 799–805.
- Bisht, S., Pandey, P., Sood, A., Sharma, S, and Bisht, N. S. (2010). Biodegradation of naphthalene and anthracene by Chemo-tactically active *Rhizobacteria* of *Populus deltoids*. *Brazilian Journal of Microbiology*, 4 (4): 922-930.
- Chaine, I., and Régnière, J. (2017). Process-based models of phenology for plants and animals. *Annual Review of Ecology, Evolution, and Systematics*, 48, 159–182.
- Denning, A. S., Nicholls, M., Prihodko, L., Baker, I., Vidale, P. L., Davis, K. and Bakwin, P. (2003). Stimulated variations in atmospheric CO₂ over a Wisconsin forest using a coupled ecosystem-atmosphere model. *Global Change Biology*, 9:1241-1250.

- Devatha, C.P., Vishnu, A. and Purna, C.R. (2019). Investigation of physical and chemical characteristics on soil due to crude oil contamination and its remediation. *Journal of Applied Water Science*, 9:89-100
- Hamann, A. (2004). Flowering and fruiting phenology of a Philippine submontane rain forest: climatic factors as proximate and ultimate causes. *Journal of Ecology*, 92:24–31.
- Jenerette, G. D., Scott, R. L. and Huete, A. R. (2010). Functional differences between summer and winter season rain assessed with MODIS phenology in a semi-arid region. *Journal of Vegetation Science*, 21 (1): 16-30.
- Johnson, O., Ronnie, L. and Ruben, S. (2019). Understanding the impacts of crude oil and its induced stresses on agrifood production: A review. *Horticulture*, 5(2): 47 – 58.
- Marqucus, M.C.M., Roper, J.J. and Salvaggio, A.P.B. (2004). Phenological patterns among plants life-form in a subtropical forest in southern Brazil. *Journal of Plant Ecology*, 172 (2): 203-213.
- Morellato, L.P.C., Alberton, B., Alvarado, S.T., Borges, B., Buisson, E., Camargo, M.G.G., Cancian, L.F., Carstensen, D.W., Escobar, D.F.E., Leite, P.T.P., Mendoza, I., Rocha, N.M.W.B., Soares, N.C., Silva, T.S.F., Staggemeier, V.G., Streher, A.S., Vargas, B.C., Peres, C.A. (2016). Linking plant phenology to conservation biology. *Biological Conservation* 195:60–72.
- Nigerian Meteorological Agency (NIMET) (2018). Weather report, <http://nimet.gov.ng>.
- Nwadinigwe, A. O. (2002). Ecological studies of Obollo Afor savannah woodland. Unpublished *Ph.D. Thesis*, University of Nigeria Nsukka.
- Nwadinigwe, A.O. (2013). *Studies on methods and analysis of vegetation*. Welfare and Industrial Promotions International Publishers, Nigeria. 216 pp.
- Okeke, F. I. and Enoch, M.A. (2016). Analyzing the effect of hydrocarbon seepage on vegetation in Ugwueme town, Awgu Local Government Area of Enugu State using normalized differencing vegetation index (NDVI) threshold classification method. *International*

- Journal of Multidisciplinary Research and Modern Education (IJMRME)*, 2 (2): 2454 – 6119.
- Osuji, L., Adesiyon, S. O. and Obute, G. C. (2004). Post impact assessment of oil pollution in Agbada west plain of Niger Delta, Nigeria: field reconnaissance and total extractable hydrocarbon content. *Chemistry and Biodiversity* 1: 1569-1578.
- Reed, B. C. (2006). Trend analysis of time-series phenology of North - America derived from satellite data, *Geoscience and Remote Sensing*, 43: 1-15.
- Saheed, R., Sattar, A., Iqbal, Z., Imran, M. and Nadeem, R. (2012). Environmental impact assessment (EM): an overlooked instrument for sustainable development in Pakistan. *Environmental Monitoring and Assessment*. 184(4): 1909-1919.
- Schwartz, M. D (2013). *Phenology: An Integrative Environmental Science*, Édition: 2nd ed. 2013. Springer, Dordrecht.
- Stucky, B. J., Guralnick, R., Deck, J., Denny, E.G., Bolmgren, K. and Walls, R. (2018). The Plant Phenology Ontology: A New Informatics Resource for Large-Scale Integration of Plant Phenology Data. *Frontiers in Plant Science*, 9: 517.
- Tanee, F.B.G. and Albert, E. (2011). Post-remediation assessment of crude oil polluted site at Kegbara-Dere Community, Gokana L.G.A. of Rivers State, Nigeria. *Journal of Bioremediation & Biodegradation*, 2: 122-131.
- Wessels, K., Steenkamp, k., Maltitz G. V. and Archibald S. (2011). Remotely sensed vegetation phenology for describing and predicting and biomes of South Africa. *Applied Vegetation Science*. 14(1): 49-66.