

Effect of foliar application of Nano-urea on yield attributes and yield of Pearl millet (*Pennisetum glaucum* L.)

ABSTRACT

To evaluate the effect of different doses of Nano-urea on yield attributes and yield of pearl millet a field experiment was conducted at Eastern Block Farm of Tamil Nadu Agricultural University, Coimbatore during Summer, 2022. This field experiment was laid out in Randomized Block Design (RBD) with ten treatments and three replications. In this experiment different doses of Nano-urea were applied along with the Recommended Dose of Nitrogen (RDN) as per the Crop Production Guide, 2020 of TNAU, Coimbatore. The results of study revealed that yield of pearl millet was significantly differed with treatments imposed. Among these treatments, application of 50% RDN as basal + 0.5% Nano-urea foliar spray @ panicle initiation and booting stage had recorded significantly higher number of productive tillers plant⁻¹ (3.79), earhead length (26.82 cm), earhead girth (7.9 cm), number of grains earhead⁻¹ (2,462), test weight (15.64 g), grain yield (3,314 kg ha⁻¹) and stover yield (5,136 kg ha⁻¹). Which was at par with treatment 50% RDN as basal + 25% RDN @ panicle initiation + 25% RDN @ booting stage (T₁₀).

Key words: Pearl millet, Nano-urea, foliar nutrition, yield attributes.

1. INTRODUCTION

Pearl millet is the world's fifth most significant crop after rice, wheat, maize and sorghum, respectively. It is frequently farmed as a rainfed crop in Africa and Southern Asia's arid and semi-arid regions and can be grown in areas where rainfall is insufficient for maize and sorghum growth (Reddy *et al.*, 2013). Pearl millet ranks fourth among cereal crops in India, behind rice, wheat and sorghum. Pearl millet may be an alternative crop with superior physiological properties to other cereals, such as drought resistance, low fertility, high salinity tolerance and high temperature tolerance. Rajasthan is the most productive state in India, producing 4.68 million tonnes from 4.28 million hectares with a productivity of 1,093 kg ha⁻¹. It has covered 0.67 L ha in Tamil Nadu, with a production of 1.85 L t and a productivity of 2,743 kg ha⁻¹ (INDIASTAT, 2019-2020).

In pearl millet, plant nutrition is critical for increased development and productivity. Nitrogen is one of the important macronutrients that plants require for their growth, development and yield (Tremblay *et al.*, 2011). The national average yield and the potential achievable yield in pearl millet are substantially different. One of the reasons for such a large discrepancy in pearl millet yields is a lack of key minerals (Khairwal *et al.*, 2007). Urea is the most widely used commercial nitrogen fertilizer in the world for increasing crop productivity. By virtue of its hydrolysis, urea elevates soil pH, resulting in massive ammonia volatilization losses (Fan and Meckenzie, 1993 and Hamid *et al.*, 1998).

To address the unbalanced and excessive usage of conventional prilled urea, various fertilizer businesses in India created nanotechnology-based Nano Urea (Liquid) fertilizer. Nano-urea comprises nitrogen particles (18-30 nm) with a surface area (10,000 times that of a 1 mm urea prill) and a greater number of particles (55,000 nitrogen particles over 1 mm urea prill). At important crop growth phases, spraying Nano-urea at a rate of 2 to 4 ml per litre of water initiates crop response, meets nutritional requirements and enhances nutrient availability in the rhizosphere. Two applications of Nano-urea to crops in a season enough to meet the plant's nitrogen requirements in above-ground tissues (Yogendra Kumar *et al.*, 2021). According to Jakhar *et al.* (2011) 50% Recommended Dose of Nitrogen (RDN) as basal + 25% Recommended Dose of Nitrogen at panicle initiation stage + 25% Recommended Dose of Nitrogen at booting stage was recorded maximum grain yield in pearl millet crop. Hence, present investigation was designed based on this schedule of fertilizer recommendation in which different doses of Nano-urea was applied at panicle initiation and booting stage to analyse the optimum dose of Nano-urea which gives higher productivity.

2. MATERIALS AND METHODS

A field experiment was conducted at Eastern Block Farm of Tamil Nadu Agricultural University, Coimbatore during Summer, 2022 to learn the effect of foliar nutrition of Nano-urea on yield attributes and yield of pearl millet crop. The experiment field fall under Western Agro Climatic Zone of Tamil Nadu. The texture of experiment soil is sandy clay loam. The soil was found to be slightly alkaline with pH of 7.9 and with low soluble salts of 0.59 dS m⁻¹. The nutrient status of soil was found to be low in available nitrogen (265 kg ha⁻¹), medium range of available phosphorus (17.5 kg ha⁻¹) and high in available potassium (766 kg ha⁻¹). Randomized Block Design with three replication and ten treatments were followed in this experiment. The treatments detailed are as follows.

T₁ - 50% RDN as basal + 0.3% Nano-urea FS @ P.I + 25% RDN @ booting stage

T₂ - 50% RDN as basal + 25% RDN @ P.I + 0.3% Nano-urea FS @ booting stage

T₃ - 50% RDN as basal + 0.3% Nano-urea FS @ P.I and booting stage

T₄ - 50% RDN as basal + 0.4% Nano-urea FS @ P.I + 25% RDN @ booting stage

T₅ - 50% RDN as basal + 25% RDN @ P.I + 0.4% Nano-urea FS @ booting stage

T₆ - 50% RDN as basal + 0.4% Nano-urea FS @ P.I and booting stage

T₇ - 50% RDN as basal + 0.5% Nano-urea FS @ P.I + 25% RDN @ booting stage

T₈ - 50% RDN as basal + 25% RDN @ P.I + 0.5% Nano-urea FS @ booting stage

T₉ - 50% RDN as basal + 0.5% Nano-urea FS @ P.I and booting stage

T₁₀ - Control (50% RDN as basal + 25% RDN @ P.I + 25% RDN @ booting stage)

Pearl millet variety CO 10 was sown at a seed rate of 5 kg ha⁻¹ with a spacing of 45 cm × 15 cm to study the effect of Nano-urea. For this experiment as per the blanket recommendation of TNAU, 70:35:35 N, P₂O₅, K₂O kg ha⁻¹ (CPG, 2020) were supplied through urea, single super phosphate and muriate of potash, respectively. Full recommended dosage of P and K and 50% dose of N were applied as basal. Remaining 50% dose of N was applied on panicle initiation stage and booting stage in the form of Nano-urea foliar spray at different doses and prilled urea as per the treatment plan. For taking the yield attributes viz., number of productive tillers plant⁻¹, ear head length, ear head girth, number of grains earhead⁻¹ and test weight (1000 grain weight), five plants from each treatment plots were selected randomly and tagged. Whereas, grain and stover yield were taken from net plot area and converted to per hectare. Data collected and analysed statistically with procedure described by Gomez and Gomez (1984). Critical differences were worked at 5% probability level and significant differences among treatments were determined.

3. RESULT AND DISCUSSION

3.1 Effect on yield attributes and yield

Yield attributing characters in pearl millet were found to be significantly influenced by application of different doses of Nano-urea (Table 1).

3.1.1 Number of productive tillers plant⁻¹

The number of productive tillers plant⁻¹ were significantly influenced by the different dose of nano-fertilizer. Among all the treatment combinations, application of 50% RDN as basal + 0.5% Nano-urea FS @ P.I and booting stage (T₉) resulted in significantly higher number of productive tillers plant⁻¹

(4.09). It was statistically on par with T₁₀ - Control (50% RDN as basal + 25% RDN @ P.I + 25% RDN @ booting stage) showed a mean value of 3.65. while the lowest number of productive tillers plant⁻¹ (3.17) was found with (T₂) -50% RDN as basal + 25% RDN @ P.I + 0.3% Nano-urea FS @ booting stage (Table 1). When Nano-urea was applied twice to the leaves, it increases the nutrient uptake through the stomatal openings and nutrient translocation within the plant. This increased nutrient uptake resulted in increased cell division, meristematic activity, and stimulation of cell elongation, which produced more productive tillers plant⁻¹. Similar results were revealed by Meena *et al.* (2017) and Jassim *et al.* (2019) in rice.

Table 1. Effect of foliar application of Nano-urea on yield attributes of pearl millet

Treatments		Productive tillers plant ⁻¹	Earhead length (cm)	Earhead girth (cm)	No. of grains earhead ⁻¹
T ₁	50% RDN as basal + 0.3% Nano-urea FS @ P.I + 25% RDN @ booting stage	3.24	19.19	6.20	1790
T ₂	50% RDN as basal + 25% RDN @ P.I + 0.3% Nano-urea FS @ booting stage	3.17	18.70	5.90	1775
T ₃	50% RDN as basal + 0.3% Nano-urea FS @ P.I and booting stage	3.24	19.64	6.41	1795
T ₄	50% RDN as basal + 0.4% Nano-urea FS @ P.I + 25% RDN @ booting stage	3.33	21.43	6.92	1862
T ₅	50% RDN as basal + 25% RDN @ P.I + 0.4% Nano-urea FS @ booting stage	3.25	20.79	6.90	1797
T ₆	50% RDN as basal + 0.4% Nano-urea FS @ P.I and	3.35	22.43	7.06	1893

	booting stage				
T ₇	50% RDN as basal + 0.5% Nano-urea FS @ P.I + 25% RDN @ booting stage	3.48	24.55	7.21	1992
T ₈	50% RDN as basal + 25% RDN @ P.I + 0.5% Nano-urea FS @ booting stage	3.41	23.60	7.20	1958
T ₉	50% RDN as basal + 0.5% Nano-urea FS @ P.I and booting stage	4.09	26.82	7.90	2472
T ₁₀	Control (50% RDN as basal + 25% RDN @ P.I + 25% RDN @ booting stage)	3.65	25.23	7.41	2120
	S.Ed	0.24	1.34	0.40	193
	CD (p=0.05)	0.52	2.82	0.85	407

3.1.2 Earhead length and girth (cm)

The earhead length and girth is influenced by the different dose of Nano-urea foliar spray. Among these treatments, T₉ (50% RDN as basal + 0.5% Nano-urea FS @ P.I and booting stage) resulted in significantly higher earhead length (26.82 cm) and earhead girth (7.9 cm). This treatment is on par with T₁₀ (50% RDN as basal + 25% RDN @ P.I + 25% RDN @ booting stage). Which resulted in an earhead length and girth of 25.23 cm and 7.41 cm, respectively. While the lowest earhead length (18.70 cm) and earhead girth (5.9 cm) were found with (T₂) -50% RDN as basal + 25% RDN @ P.I + 0.3% Nano-urea FS @ booting stage (Table 1). Because of the faster growth rate and photosynthetic assimilation rate brought on by the improved availability and translocation of nutrients, which in turn encouraged a greater partitioning of photosynthates, as a result there has been a considerable rise in earhead length and girth. Similar findings were reported by Choudhary *et al.* (2019) in corn.

3.1.3 No. of grains earhead⁻¹

Data revealed that significantly higher No. of grains earhead⁻¹ (2472) were reported in T₉ (50% RDN as basal + 0.5% Nano-urea FS @ P.I and booting stage). Whereas significantly lowest no. of grains earhead⁻¹ (1775) were found in (T₂) -50% RDN as basal + 25% RDN @ P.I + 0.3% Nano-urea FS @ booting stage (Table 1). The considerable increase in grain earhead-1 counts may be attributable to the increase in overall plant growth brought on by the increased availability of nutrients as a result of

fertiliser foliar spraying. In addition, timely nitrogen delivery increases the start of grain formation by increasing photosynthate uptake and translocation from source to sink. These results were in relation with findings of Algym *et al.* (2020) in corn.

3.1.4 Test weight

In this field trial, test weight (1000 grain weight) was significantly influenced by different dose of Nano-urea (Table 2). Application of 50% RDN as basal + 0.5% Nano-urea FS @ P.I and booting stage (T_9) had produced maximum test weight of 15.71 g, which is on par with T_{10} (50% RDN as basal + 25% RDN @ P.I + 25% RDN @ booting stage) (14.20 g). The lower test weight of 12.26 g was recorded with in (T_2) -50% RDN as basal + 25% RDN @ P.I + 0.3% Nano-urea FS @ booting stage. This is a result of the enhanced level of Nano-urea foliar spray, which may have guaranteed nitrogen supply at later phases of grain filling. Foliar application of nutrients may promote assimilate build-up in the grain and increasing grain weight as reported by Manikandan and Subramanian (2016) in wheat and Naveenaa *et al.* (2018) in maize crop.

3.1.5 Grain and stover yield

In this field investigation, the grain yield and stover yield were also following the similar trends as yield attributes. The yield of pearl millet was significantly influenced by different dose of Nano-urea as presented in Table 2.

Significantly higher yield of 3413 kg ha⁻¹ was observed with the treatment 50% RDN as basal + 0.5% Nano-urea FS @ P.I and booting stage (T_9). Which is statistically on par with T_{10} (50% RDN as basal + 25% RDN @ P.I + 25% RDN @ booting stage) (3377 kg ha⁻¹). The lowest grain yield was recorded with (T_2) -50% RDN as basal + 25% RDN @ P.I + 0.3% Nano-urea FS @ booting stage (2552 kg ha⁻¹). Due to the improvement in yield attributes, a considerable increase in grain yield was seen with greater levels of foliar application of nano-nutrients. Also, it led to efficient photosynthetic translocation. These findings were reported by Kumar *et al.* (2020).

Data noted on stover yield was significantly influenced by the application of Nano-urea. Higher stover yield was observed in 50% RDN as basal + 0.5% Nano-urea FS @ P.I and booting stage (T_9) (5662 kg ha⁻¹). Which is statistically on par with the treatment T_{10} (50% RDN as basal + 25% RDN @ P.I + 25% RDN @ booting stage) (5481 kg ha⁻¹). Treatment T_2 (50% RDN as basal + 25% RDN @ P.I + 0.3% Nano-urea FS @ booting stage) recorded the lowest stover yield of 4160 kg ha⁻¹. The increase in the stover yield with the foliar spray of higher dose of Nano-urea might be due to their rapid uptake by plants and ease of translocation at a faster pace, that aided in higher rate of photosynthesis and more dry matter accumulation which resulted in higher stover yield. These findings were in agreement with the reports of Khalil *et al.* (2019) in maize and Sahu *et al.* (2022) in rice.

Table 2. Effect foliar application of Nano-urea on Test weight, Grain yield and Stover yield of Pearl millet

Treatments		Test weight (g)	Grain yield (kg ha ⁻¹)	Stover yield (kg ha ⁻¹)
T ₁	50% RDN as basal + 0.3% Nano-urea FS @ P.I + 25% RDN @ booting stage	12.31	2568	4169
T ₂	50% RDN as basal + 25% RDN @ P.I + 0.3% Nano-urea FS @ booting stage	12.26	2552	4160
T ₃	50% RDN as basal + 0.3% Nano-urea FS @ P.I and booting stage	12.68	2665	4333
T ₄	50% RDN as basal + 0.4% Nano-urea FS @ P.I + 25% RDN @ booting stage	13.64	2826	4597
T ₅	50% RDN as basal + 25% RDN @ P.I + 0.4% Nano-urea FS @ booting stage	13.26	2825	4552
T ₆	50% RDN as basal + 0.4% Nano-urea FS @ P.I and booting stage	14.14	2844	4581
T ₇	50% RDN as basal + 0.5% Nano-urea FS @ P.I + 25% RDN @ booting stage	14.19	3016	4902
T ₈	50% RDN as basal + 25% RDN @ P.I + 0.5% Nano-urea FS @ booting stage	14.18	2922	4597
T ₉	50% RDN as basal + 0.5% Nano-urea FS @ P.I and booting stage	15.71	3413	5662
T ₁₀	Control (50% RDN as basal + 25% RDN @ P.I + 25% RDN @ booting stage)	14.20	3377	5481
S.Ed		0.95	270.05	425.34
CD (p=0.05)		2.01	567.36	893.63

4. CONCLUSION

From the field experiment, it was concluded that significantly higher yield attributes and yield character of pearl millet were observed with application of 50% RDN as basal + 0.5% Nano-urea foliar

spray @ panicle initiation and booting stage (T₉). Which was at par with treatment 50% RDN as basal + 25% RDN @ panicle initiation + 25% RDN @ booting stage (T₁₀).

REFERENCES

1. Algym AJK, Alasady MHS. Effect of the method and level of adding NPK nanoparticles and mineral fertilizers on the growth and yield of yellow corn and the content of mineral nutrient of some plant parts. *Plant Archives* 2020;20(1):38-43.
2. Choudhary RC, Kumaraswamy RV, Kumari S, Sharma SS, Pal A, Raliya R, Biswas P, Saharan V. Zinc encapsulated chitosan nanoparticle to promote maize crop yield. *International Journal of Biological Macromolecules* 2019;127(1):126-135.
3. CPG. Crop Production Guide, Department of Agriculture, Government of Tamil Nadu, Chennai and Tamil Nadu Agricultural University, Coimbatore 2020.
4. Fan MX, AF Mackenzie. Urea and phosphate interaction in fertilizer microsites: ammonia volatilization and pH changes. *Soil Science Society of America Journal* 1993;57(1):839-844.
5. Gomez Kwanchai A, Arturo A Gomez. *Statistical procedures for agricultural research*. John Wiley and Sons 1984.
6. Hamid A, M Ahmad, M Yaqub. Behaviour of nitrogen fertilizers in alkaline calcareous soils. *Pakistan Journal of Soil Science* 1998;15(1):56-61.
7. Indiastat. Area, production and yield of principal crops, Directorate of Economics and Statistics, Department of Agriculture and Cooperation report, Government of India, New Delhi 2020, available on www.indiastat.com.
8. Jakhar GR, Golada SI, Sadhu AC. Influence of levels and time of application of nitrogen on growth, yield and nitrogen uptake by pearl millet during summer. *Madras Agriculture Journal* 2011;98(10/12):347-349.
9. Jassim RAH, Kadhem HN, Nooni GB. Impact of levels and time of foliar application of nano fertilizer (super micro plus) on some components of growth and yield of rice (*Oryza sativa* L.). *Plant Archives* 2019;19(1):1279-1283.
10. Khairwal IS, KN Rai, B Diwakar, YK Sharma, BS Rajpurohit, B Nirwan, R Bhattacharjee. *Pearl Millet: Crop Management and Seed Production Manual*. ICRISAT 2007.
11. Khalil MH, Abou AAF, Abdrabou RTH, Abdalhalim SH, Abdelmaaboud MSH. Response of two maize cultivars (*Zea mays* L.) to organic manure and mineral nano nitrogen fertilizer under siwa oasis conditions. *AUJAS, Ain Shams Univ., Cairo, Egypt*. 2019;27(1):299-312.
12. Kumar Y, Tiwari KN, Singh T, Sain NK, Laxmi S, Verma R, et al. Nano fertilizers for enhancing nutrient use efficiency, crop productivity and economic returns in winter season crops of Rajasthan. *Annals of Plant and Soil Research*. 2020;22(4):324-335.
13. Manikandan A, KS Subramanian. Evaluation of zeolite based nitrogen nano-fertilizers on maize growth, yield and quality on inceptisols and alfisols. *International Journal of Plant and Soil Science* 2016;9(4):1-9.

14. Meena DS, Gautham C, Patidar O, Meena PHM, Prakasha G, Viswa J. Nanofertilizers are a new way to increase nutrient use efficiency in crop production. *International Journal of Agriculture sciences* 2017;1(1):975-3710.
15. Naveenaa M, A Amarerouda, MK Meena, TC Suma, PH Kunchanur. Influence of foliar nutrition at different vegetative stages on growth and yield performance of maize (*Zea mays* L.). *Journal of Pharmacognosy and Phytochemistry* 2018;7(5):249-255.
16. Reddy AA, PP Rao, OP Yadav, IP Singh, NJ Ardeshna, KK Kundu. Prospects for kharif (Rainy Season) and summer Pearl Millet in Western India. Working Paper Series no. 36, ICRISAT. Patancheru 502 324, Andra Pradesh, India 2013, 24.
17. Sahu TK, Manish Kumar, Narendra Kumar, T Chandrakar, DP Singh. Effect of nano urea application on growth and productivity of rice (*Oryza sativa* L.) under midland situation of Bastar region. *The Pharma Innovation* 2022;11(6):185-187.
18. Tremblay N, Fallon E, Ziadi N. Sensing of crop nitrogen status: opportunities, tools, limitation and supporting information requirements. *Hort Technology* 2011;21(1):274-281.
19. Yogendra Kumar KN, Tiwari Tarunendu Singh, Ramesh Raliya. Nanofertilizers and their role in sustainable agriculture. *Annals of Plant and Soil Research* 2021;23(3):238-255.