

## Original Research Article

# Silicon Sources and Bacterial Inoculants on Growth Parameters, Leaf Yield, Quality of Coriander (*Coriandrum sativum* L) and Soil Adsorbed Silicon in Sandy Loamy Soil

### ABSTRACT

**Aims:** The objective of the present study was to investigate the effect of different silicon sources with and without microbial inoculants on growth parameters, leaf yield and quality of Coriander.

**Study design:** Factorial Completely Randomized Block Design (FCRD) was used in the present study with thirty-six treatments and three replications.

**Place and Duration of Study:** The pot culture experiment was carried out in the Department of Soil Science and Agricultural Chemistry, Tamil Nadu Agricultural University, Coimbatore, India during January to March 2022.

**Methodology:** A pot culture experiment was conducted with different silicon sources namely  $\text{CaSiO}_3$ ,  $\text{MgSiO}_3$  and Rice Husk Ash (RHA) at three levels (0, 50 and  $100 \text{ mg Si kg}^{-1}$ ) with and without two silicate solubilizing bacteria (SSB) and a zinc solubilizing bacterium (ZSB) at  $2 \text{ kg ha}^{-1}$  with coriander CO (CR) 4. The influence of the treatments on growth parameters, leaf Si content, leaf yield, total soluble sugar content of leaf and soil adsorbed Si were investigated at 25 and 50 DAS.

**Results:** The results revealed significant influence of Si sources with bacterial inoculants. The plant height and SPAD index values were the highest by the application of RHA @  $100 \text{ mg Si kg}^{-1}$  + SSB4 followed by  $\text{CaSiO}_3$  @  $100 \text{ mg Si kg}^{-1}$  + SSB4. All the sources increased the leaf Si as well as soil adsorbed Si and the effect was highly significant when Si sources were combined with SSB4 and other inoculants. Coriander leaf yield ranged from 4.321 to  $8.237 \text{ g plant}^{-1}$ . The maximum leaf yield was recorded with the application of RHA @  $100 \text{ mg Si kg}^{-1}$  with SSB4 ( $8.237 \text{ g plant}^{-1}$ ) followed by RHA @  $50 \text{ mg Si kg}^{-1}$  with SSB4 ( $7.669 \text{ g plant}^{-1}$ ) and  $\text{CaSiO}_3$  @  $100 \text{ mg Si kg}^{-1}$  with SSB4 ( $7.563 \text{ g plant}^{-1}$ ). The effect of Si sources with bacterial inoculants on TSS content of leaf was significant with a maximum value of ( $38.227 \text{ mg g}^{-1} \text{ DW}$ ) by RHA @  $100 \text{ mg Si kg}^{-1}$  + SSB4 followed by  $\text{CaSiO}_3$  @  $100 \text{ mg Si kg}^{-1}$  + SSB4 ( $36.103 \text{ mg g}^{-1} \text{ DW}$ ).

**Conclusion:** The current study revealed that the combined application of Si sources and microbial inoculants significantly enhanced the growth attributes, leaf Si, leaf yield, leaf quality of coriander and soil adsorbed Si. The leaf yield increase over control for RHA @  $100 \text{ mg Si kg}^{-1}$  with SSB4 was 88.92 per cent followed by RHA @  $50 \text{ mg Si kg}^{-1}$  with SSB4 (76.58 %) and  $\text{CaSiO}_3$  @  $100 \text{ mg Si kg}^{-1}$  with SSB4 (73.82%). The research work needs further validation at the field level.

**Keywords:** Silicon sources; leaf yield; Si content; coriander; *Bacillus altitudinis*; *Pseudomonas gessardii*; *Bacillus aryabhatai*.

## 1. INTRODUCTION

The second most abundant element in the Earth's crust after oxygen is Silicon (Si). The uptake of other nutrients by plants is positively impacted by Si. Due to its strong affinities for oxygen, silicon (Si) typically appears in nature as silica ( $\text{SiO}_2$ ) [1]. Depending on the pH of the soil, it can also be found as silicic acid [ $\text{H}_4\text{SiO}_4$ ] and silicate [2]. However, the capacity of different plant species to accumulate silicon varies substantially, with values ranging from 0.1 percent to 10 percent silicon on a dry weight basis [2,3]. Si moves from the root to the shoot area after being absorbed by the soil, where it can trigger a variety of physiological responses, such as crop improvement and growth [4,5], enzymatic activities [6,7] and expression of genes [8,9]. Si is able to regulate the availability of several elements in soils by competing with silicic acid for binding on soil particles [10].

Si increases the rate of photosynthesis, which is affected by the ultrastructure of the leaf, the level of chlorophyll, and the activity of the enzyme ribulose biphosphate carboxylase [11]. In addition to making plant cell walls thicker and more durable, silicon also expands the vascular system [12]. The increased vascular system allows for greater water and nutrient uptake, strengthening the plant in all ways. Si spraying dramatically increased relative water content (RWC) by promoting plant development, enhancing the capacity of roots to absorb more water from the growth medium, and enhancing water flow from roots to leaves. It also modulates plant water potential, stomatal conductance, and transpiration rate [13]. By causing a number of transcriptional changes, Si plays a significant part in the creation of elicitor-accelerated secondary metabolites [14]. The silicate minerals that are formed during weathering and are dissolved in the soil solution are affected by diffusion and leaching [15].

Coriander (*Coriandrum sativum* L), is one of the India's major spice crops [16], known as Dhania, Cilantro, Chinese parsley, Mexican parsley, and Japanese parsley [17], a member of the family *Apiaceae* or *Umbelliferae*. It is widely used in the food, beverage, pharmaceutical, cosmetic, and sanitary industries [18,19]. It is a culinary and medicinal plant, that originated in southern Europe and the Mediterranean region. Coriander is grown all year long in India for both tender leaves and grains. The best time to grow coriander is during the rabi season [20]. Although any type of soil, including light, well-drained, moist, loamy, and light to heavy black soil, can support its growth, dry climates are best for it [21]. In India, this crop is cultivated in an area of 2,94,542 ha with an estimated yield of 11,01,920 metric tonnes in 2021 [22]. On a bigger scale, it is grown in Rajasthan, Madhya Pradesh, Tamil Nadu, Gujarat, Karnataka, Maharashtra, Orissa, and Andhra Pradesh. The aroma of the coriander fruit is significantly different from that of the herb, which is due to changes in essential oil and its chemical composition throughout ontogenesis [23,24]. Its green leaf, which is rich in fiber, carbohydrates, proteins, vitamins, and minerals (including calcium, phosphorus, and iron), is used as a vegetable and in salads as well as in cooking as an ingredient in many foods [25]. Indigestion, dyspepsia, flatulence, and piles can all be treated with the seeds and leaves of this plant [26].

Numerous studies have demonstrated that using bio-fertilizers instead of synthetic fertilizers can increase crop yields and plant growth while using less dangerous agrochemicals. Based on specific strains of the genus *Bacillus* bacteria that have been proven to have natural benefits, Silicate Solubilizing Bacteria (SSB) are called bio-fertilizers. These bacteria provide the ability to plants to withstand biotic and abiotic stress by solubilizing silicon. In addition to the insoluble forms of silicon, silicate-solubilizing bacteria (SSB) may also solubilize potassium and phosphate which helps in increasing soil fertility and crop yield [27]. Zinc Solubilizing Bacteria (ZSB) also act as plant growth-promoting factors [28,29] which produce Indole acetic acid (IAA) and due to this it may have an impact on the growth of various plant species [30]. According to Narayanaswamy and Prakash [31], paddy plants cultivated on

Inceptisol soils eliminated a total of 205 to 611 kg/ha of Si. Si may become a yield-limiting component for rice cultivation; hence the use of exogenous Si fertilizer may be required for a system of rice production that is both profitable and sustainable [32]. The research on Silicon sources with SSB on Coriander in respect of leaf yield and quality is very meagre. Hence this study was carried out to investigate the effect of different Silicon sources with and without microbial inoculants on growth parameters, leaf yield, and quality of Coriander.

## 2. MATERIAL AND METHODS

### 2.1 Pot Culture Experiment

A pot culture experiment was conducted at the Department of Soil Science and Agricultural Chemistry, Tamil Nadu Agricultural University, Coimbatore during January to March 2022. The bulk soil sample low in adsorbed silicon collected from Madukkarai, Coimbatore was processed and each pot was filled with ten kg of soil. The soil was alkaline in soil reaction with an EC of 0.27 dS m<sup>-1</sup>, low organic carbon (0.32 %) and low organic matter (0.55 %). The available nitrogen (183.86 kg ha<sup>-1</sup>) was low whereas available phosphorus (15.51 kg ha<sup>-1</sup>) and available potassium (222 kg ha<sup>-1</sup>) were medium in status. The adsorbed silicon (0.5 M acetic acid method) of experimental soil was low (11.489 mg Kg<sup>-1</sup>). Soils were treated with three different Silicon sources viz., Calcium silicate (CaSiO<sub>3</sub>), Magnesium Silicate (MgSiO<sub>3</sub>), and Rice Husk Ash (RHA) with three levels (0, 50 and 100 mg Si kg<sup>-1</sup>). Two Silicate Solubilizing Bacteria (*Bacillus altitudinis* SSB4 & *Pseudomonas gessardii* SSB7) and one Zinc Solubilizing Bacteria (ZSB) (*Bacillus aryabhatai* KSBN<sub>2</sub>K<sub>7</sub>) were applied to the soil at 2 kg ha<sup>-1</sup>.

We used Coriander CO (CR) 4 seeds that were collected from the Department of Spices and Plantation Crops, Tamil Nadu Agricultural University, Coimbatore. Seeds were split in half, immersed in water overnight, and then treated with *Trichoderma viridi* before sowing. Seeds were sown at 1 cm depth and sufficient irrigation is applied to pots till the last stage. Throughout the growing season, standard management practices were followed as per Crop Production Guide, 2020, TNAU. After two weeks of germination, they were reduced to ten plants per pot. The plant samples were collected at two stages viz., 25 DAS and 50 DAS (Days After Sowing) and analyzed for various parameters by adopting standard procedures. The study was laid out in a Factorial Completely Randomized Block Design (FCRD) consisting of thirty-six treatments and replicated thrice.

### 2.2 Microbial culture

SSB4 (*Bacillus subtilis*), SSB7 (*Pseudomonas gessardii*) and ZSB (KSBN<sub>2</sub>K<sub>7</sub>, *Bacillus aryabhatai*) were obtained from the Department of Agricultural Microbiology, Tamil Nadu Agricultural University, Coimbatore.

### 2.2 Plant Height (cm)

Plant height was measured using a scale from the base of the plant to the tip at two stages viz., 25 DAS & 50 DAS (Days After Sowing). The plant height was calculated using the average of five plants in every pot and expressed in centimeters (cm).

### 2.3 Leaf SPAD Index Value

Leaf SPAD (Soil and Plant Analysis Development; an indication of leaf greenness) values were measured in the second expanded leaves from the top at two cropping stages (25 DAS & 50 DAS) using a portable SPAD-502 Chlorophyll Meter (Minolta Camera Co. Ltd., Tokyo, Japan). The average leaf SPAD value was calculated from 10 readings of plant leaves per pot.

## 2.4 Coriander Leaf Yield (g/plant)

The plants were chosen randomly from each treatment, and the fresh weight of the herbage was measured by removing any adherent soil from the roots and expressed as grams per plant.

## 2.5 Estimation of Leaf Silicon

To analyze plant Si, dry powdered coriander leaf samples (0.1 g) were digested in a mixture of 70% HNO<sub>3</sub> (7 ml), 30% H<sub>2</sub>O<sub>2</sub> (2 ml), and 40% Hydrofluoric acid (HF) (1 ml) with the help of a microwave digestion system (Anton Paar- Multiwave PRO) with following steps: 1200 watts for 20 minutes with a ramping rate of 7 °C per minute and 1200 watts for 30 min at holding temperature of 170 °C and venting for 20 min. The digested samples were diluted with 4% boric acid and made up the volume of 50 ml [31]. The Si content of digested samples was determined by the standard Spectrophotometry method as described by Ma *et al.*, 2001 [33].

## 2.6 Estimation of Adsorbed Silicon (Ad-Si) in Soils

Adsorbed silicon (Ad-Si) in soils was estimated using the 0.5 M acetic acid method. 5 gm soil was added into a 50 ml plastic centrifuge tube then 12.5 ml of 0.5 M acetic acid was added [34]. After shaking in an end-over-end shaker continuously for one hour, centrifuged the samples at 3000 rpm for 3 minutes. The supernatant was filtered and used to measure the amount of adsorbed silicon [31]. 0.25 ml of filtrate was taken as an aliquot, and 10.5 ml of distilled water and 0.25 ml of 1:1 hydrochloric acid were added to the centrifuge tube. After allowing it to stand for 2 minutes, 0.5 ml of 10 percent ammonium molybdate solution (pH = 7.5) was added. After 5 minutes, 0.5 ml of 20 percent tartaric acid solution was added. Then two additional minutes, a 0.5 ml reducing agent (1-amino-2-naphthol-4-sulfonic acid, ANSA) was added. After 5 minutes, but no later than 30 minutes of adding the reducing agent, absorbance was measured at 630 nm using a UV/VIS spectrophotometer (LABINDIA Analytical, UV-3200 series). Si standards (0, 0.2, 0.4, 0.8, 1.2, and 1.6 mg Si L<sup>-1</sup>) were prepared simultaneously and measured in a UV/VIS spectrophotometer.

Calculation:

Ad-Si (ppm or mg kg<sup>-1</sup>) = (ppm obtained from standard graph x Volume of extractant x Volume made) / (Weight of soil x Aliquot taken)

## 2.7 Total soluble sugar (TSS)

TSS was analyzed using the anthrone method as described [35] with glucose serving as the standard. 100mg of dried coriander leaves were put into a centrifuge tube and 1.5 mL of 80% ethanol was added for homogenization. After that kept it for 30 min for incubation in a water bath at 100°C, then the mixture was cooled rapidly and centrifuged at 6000 rpm for 10 min at 4°C. The supernatant was collected and measured at 620 nm using UV/VIS spectrophotometer.

## 2.8 Statistical analysis

The data for the various parameters obtained from the experiment were statistically analyzed by AGRES software version 7.01. To detect differences between treatments that were statistically significant via analysis of variance (ANOVA), Duncan's multiple range test (DMRT) were used at the 5% level of significance ( $P= 0.05$ ) [36].

## 3. RESULTS AND DISCUSSION

The different silicon sources and microbial inoculants significantly influenced the plant growth parameters, leaf yield and quality of Coriander.

### 3.1 Plant Height

The plant height of coriander ranged from 8.31 to 15.98 cm at 25 DAS and 18.18 to 38.48cm at 50 DAS due to the influence of Si sources and bio-inoculants (Table 1). The treatment of rice husk ash @ 100 mg Si kg<sup>-1</sup> + SSB4 recorded the maximum plant height among the prescribed treatments, whereas the control displayed the minimum plant height at 25 DAS. At 50 DAS, the maximum plant height of 38.48 cm was recorded by rice husk ash @ 100 mg Si kg<sup>-1</sup> + SSB4 application while the control treatment showed a minimum value. RHA recorded the highest value of plant height among the various silicon sources, followed by calcium silicate and magnesium silicate. With a rise in silicon fertilizer levels, the plant height grew significantly. The plant height was increased by the bacterial strains in the sequence of without microbes < ZSB < SSB7 < SSB4. The interaction effects were found to be significant at both the stages *i.e.*, 25 & 50 DAS as shown in Table 1. Ahmad *et al.*, 2013 [37] and Pati *et al.*, 2016 [38] showed a similar increase in plant height with the addition of Si fertilizers over control and claimed that the applied silicon promotes water status and increases plant photosynthetic rate, making the crop long and firm. According to [39], the erectness of the leaves and stems, is a result of Si deposition in the cell wall, which can result in an increase in rice plant height [38]. According to [40,41], the stem diameter was linearly related to the rate of Si applied, whereas the height of the sugarcane crop was quadratically related. Goteti *et al.*, 2013 [28] reported that ZSB inoculation improves the maize shoot height.

**Table 1. Silicon Sources and bacterial Inoculants on plant height of Coriander**

Plant Height (cm) at 25 DAS																
Si Sources Si Levels (mg Si kg <sup>-1</sup> )	No Microbes				SSB4				SSB7				ZSB			
	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean
0	8.31	8.47	8.56	8.45	11.68	11.51	11.57	11.59	10.30	9.78	10.31	10.13	9.55	9.51	9.53	9.53
50	9.91	9.36	10.58	9.95	14.89	12.29	14.68	13.95	11.24	10.36	12.23	11.27	10.44	9.82	10.69	10.32
100	11.03	10.56	12.29	11.29	15.22	13.46	15.98	14.89	13.02	11.14	13.57	12.57	11.89	10.71	12.89	11.83
Mean	9.75	9.46	10.48	9.90	13.93	12.42	14.08	13.48	11.52	10.43	12.04	11.33	10.63	10.01	11.04	10.56
Factors	M		S		L		M x S		M x L		S x L		M x S x L			
SEd	0.075		0.065		0.065		0.130		0.130		0.112		0.225			
LSD (P=0.05)	0.149		0.129		0.129		0.259		0.259		0.224		0.448			
Plant Height (cm) at 50 DAS																
Si Sources Si Levels (mg Si kg <sup>-1</sup> )	No Microbes				SSB4				SSB7				ZSB			
	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean
0	18.93	18.18	18.89	18.67	28.90	29.06	29.57	29.17	25.78	25.83	26.10	25.90	21.93	21.18	21.89	21.67
50	23.23	21.94	24.81	23.33	32.59	32.33	33.97	32.96	31.00	30.02	33.33	31.45	27.44	26.00	29.00	27.48
100	25.20	23.57	29.55	26.11	34.72	34.01	38.48	35.74	34.01	32.61	35.42	34.01	32.75	29.25	34.71	32.24
Mean	22.46	21.23	24.42	22.70	32.07	31.80	34.01	32.62	30.27	29.49	31.62	30.46	27.38	25.48	28.54	27.13
Factors	M		S		L		M x S		M x L		S x L		M x S x L			
SEd	0.17		0.147		0.147		0.295		0.295		0.255		0.511			
LSD (P=0.05)	0.339		0.294		0.294		0.588		0.588		0.509		1.018			

DAS – Days After Sowing; RHA – Rice Husk Ash; SSB – Silicate Solubilizing Bacteria; ZSB – Zinc Solubilizing Bacteria;  
M – Microbes; S – Sources; L – Levels; SEd – Standard Error of difference; LSD – Least Significant Difference;  
SSB4 - *Bacillus altitudinis*; SSB7 - *Pseudomonas gessardii*; ZSB - *Bacillus aryabhatai*

### 3.2 SPAD Index values

SPAD Index values increased from 21.10 to 40.31 at 25 DAS while at 50 DAS it ranged from 19.15 to 35.68 as shown in Table 2. Among the treatments, application of rice husk ash @ 100 mg Si kg<sup>-1</sup> with SSB4 showed the maximum SPAD values at both stages due to silicon fertilization which might have increased the chloroplasts size and number of grana in leaves [42] whereas the control treatment displayed a minimum value at 25 and 50 DAS. SPAD Index values increased linearly with increased levels of silicon sources. The mean SPAD value at 25 DAS increased from 24.02 to 29.62, 25.78 to 33.24, and 28.18 to 35.06 in the case of MgSiO<sub>3</sub>, CaSiO<sub>3</sub>, and RHA, respectively while at 50 DAS it was 20.67 to 26.91, 22.18 to 29.92 and 24.26 to 31.52. The SPAD index values at 50 DAS were lower compared to the values at 25 DAS irrespective of sources, levels of Si and bio inoculants. SSB4 showed higher SPAD Index values compared to SSB7 and ZSB, while treatments without microbes displayed lower values. Si causes an increase in growth that is accompanied by an improved rate of photosynthesis in various crops [43,11] which is ultimately influenced by leaf ultrastructure, chlorophyll concentration, and ribulose biphosphate carboxylase activity. According to Soundararajan *et al.*, 2014 [44], Si changes the leaf posture and increases total leaf area, increasing the amount of light that plants can absorb, potentially leading to increased chlorophyll synthesis.

### 3.3 Coriander Leaf Si


The effect of silicon sources with bacterial inoculants on leaf silicon percentage was presented in Table 3. At 25 DAS, the leaf silicon ranged from 1.352 to 3.384 per cent, where in the applied rice husk ash @ 100 mg Si kg<sup>-1</sup> + SSB4 and control displayed maximum ( 3.38 %) and minimum (1.352 %) leaf silicon percentage, respectively. The leaf silicon ranged from 2.01 per cent (control) to 4.11 per cent (Rice Husk Ash @ 100 mg Si kg<sup>-1</sup> + SSB4) at 50 DAS. The mean leaf silicon at 25 DAS increased from 1.497 to 1.994, 1.594 to 2.203, and 1.756 to 3.384 per cent in the case of MgSiO<sub>3</sub>, CaSiO<sub>3</sub>, and RHA, respectively while at 50 DAS it was 2.187 to 2.913, 2.347 to 3.219 and 2.566 to 3.447 per cent. RHA increased plant silica content, which suggested that when it decomposes over time, more silica will be exposed because phytoliths will be freed from the plant matrix and dissolved [45]. Si content in coriander leaves might have increased because CaSiO<sub>3</sub> was more reactive and effective at supplying Si to the soil and plants than MgSiO<sub>3</sub> [46]. The leaf silicon continuously increased as silicon fertilizer levels increased. The leaf silicon is increased significantly by the bacterial strains in the sequence of without bacteria < ZSB < SSB7 < SSB4 at both stages. The results are in accordance with [47,48,49] who suggested the improvements of Si content in rice seedlings through the application of rice husk ash (RHA) and other soil organic matters [50,51,52]. Peera SP *et al.*, 2016 [53] found that the application of SSB with FYM significantly increased the silicon content in rice. According to Etesami *et al.*, 2022 [54], inoculating Japonica rice plants with the SSB strain led to an increase in the plants' Si content.

### 3.4 Coriander Leaf Yield

The effect of silicon sources with bacterial inoculants on the leaf yield of coriander was significant and furnished in Table 4. Different silicon sources increased the leaf yield of coriander which was in the sequence of MgSiO<sub>3</sub> < CaSiO<sub>3</sub> < RHA. The maximum coriander leaf yield was recorded with the application of rice husk ash @ 100 mg Si kg<sup>-1</sup> with SSB4 (8.237 g plant<sup>-1</sup>) followed by Rice Husk Ash @ 50 mg Si kg<sup>-1</sup> with SSB4 (7.669 g plant<sup>-1</sup>) and CaSiO<sub>3</sub> @ 100 mg Si kg<sup>-1</sup> with SSB4 (7.563 g plant<sup>-1</sup>). The effect of RHA was found to be superior compared to CaSiO<sub>3</sub> and MgSiO<sub>3</sub>. However, the influence of RHA @ 50 mg Si kg<sup>-1</sup>

and  $\text{CaSiO}_3$  @  $100 \text{ mg Si kg}^{-1}$  were found to be comparable with each other, when the sources were combined with SSB4. With SSB7 the higher doses of RHA @ 50 and  $100 \text{ mg Si kg}^{-1}$  were at a par with each other with values of  $7.392$  and  $7.129 \text{ g plant}^{-1}$ , respectively. Among the inoculants, the mean values of leaf yield were the highest for SSB4 ( $6.587 \text{ g plant}^{-1}$ ) compared to SSB7 ( $5.855 \text{ g plant}^{-1}$ ) and ZSB ( $5.239 \text{ g plant}^{-1}$ ) irrespective of Si sources. Whereas, the mean leaf yield was  $5.082 \text{ g plant}^{-1}$  only without microbial inoculants. The results obtained by Peera SP *et al.*, 2016 [53] was in corroboration with the current findings. The control without silicon sources and without microbial inoculants showed a minimum leaf yield. As the amount of silicon fertilizer improved, the leaf yield increased with time. Both Deren CW I *et al.*, 1994 [55] and Pati *et al.*, 2016 [38] noted the same result in rice crops with rising Si levels. When compared to SSB7 and ZSB, SSB4 revealed greater leaf yield per plant, whereas treatments lacking microorganisms showed lower values. Due to SSB ability to dissolve insoluble types of silicates, soil fertility is increased, and plant defense mechanisms are strengthened [56]. Afshari M *et al.*, 2021 and Hassanein RA *et al.*, 2021 [57,58] observed that the application of silicon increased the leaf yield of coriander. Effect of silicon sources with and without microbes on per cent leaf yield of coriander in shown in Fig.1.

Many studies reported that Si fertilization can increase the number of grains [59,60] as well as the number of tillers per hill [61,62,38], by improving carbohydrates in a panicle. According to numerous studies [63,64,65,66], exogenous Si delivery to various horticultural crops, including cucumber, squash, bean, tomato, and roses, may lead to improved growth and productivity. Singh KK *et al.*, 2006 and Ma *et al.*, 1989 [67,68], suggested that the application of Si improved spikelets number per panicle, mature grain percentage, dry matter, and yield of rice. Application of calcium silicate as a Si source significantly increases rice [69] and sugarcane [70] yield over the control. Peera SP *et al.*, 2016 [53] found that the application of SSB with FYM significantly increases plant growth. Rosas SB *et al.*, 2009 [71] found that addition of ZSB to sandy loam soil in Argentina boosts wheat production by 36 per cent.



**Fig. 1. Silicon sources with and without *Bacillus altitudinis* (SSB4) on per cent leaf yield increase in Coriander**

UNDER PEER REVIEW

**Table 2. Silicon Sources and bacterial Inoculants on SPAD Index Values of Coriander**

SPAD Index Values at 25 DAS																
Si Sources Si Levels (mg Si kg <sup>-1</sup> )	No Microbes				SSB4				SSB7				ZSB			
	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean
0	22.25	22.10	22.30	22.22	26.46	26.56	26.97	26.66	24.77	24.53	24.64	24.65	22.48	22.30	22.45	22.41
50	27.12	24.42	30.28	27.27	35.67	29.75	37.91	34.44	32.27	25.88	35.23	31.12	27.82	25.02	30.79	27.88
100	27.96	25.55	31.98	28.49	37.59	32.56	40.31	36.82	34.01	27.75	36.19	32.65	31.05	26.06	33.11	30.07
Mean	25.78	24.02	28.18	25.99	33.24	29.62	35.06	32.64	30.35	26.05	32.02	29.47	27.12	24.46	28.78	26.79
Factors	M		S		L		M x S		M x L		S x L		M x S x L			
SEd	0.210		0.182		0.182		0.363		0.363		0.314		0.629			
LSD (P=0.05)	0.418		0.362		0.362		0.724		0.724		0.627		1.254			
SPAD Index Values at 50 DAS																
Si Sources Si Levels (mg Si kg <sup>-1</sup> )	No Microbes				SSB4				SSB7				ZSB			
	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean
0	19.15	19.02	19.19	19.12	24.50	24.65	24.76	24.64	21.32	21.11	21.21	21.21	19.34	19.19	19.32	19.29
50	23.34	21.02	26.06	23.47	30.70	25.60	34.12	30.14	27.77	22.27	32.39	27.48	23.95	21.53	26.49	23.99
100	24.06	21.99	27.52	24.52	34.56	30.47	35.68	33.57	33.19	23.01	34.78	30.33	26.72	22.42	28.49	25.88
Mean	22.18	20.67	24.26	22.37	29.92	26.91	31.52	29.45	27.43	22.13	29.46	26.34	23.34	21.05	24.77	23.05
Factors	M		S		L		M x S		M x L		S x L		M x S x L			
SEd	0.329		0.285		0.285		0.571		0.571		0.494		0.988			
LSD (P=0.05)	0.657		0.569		0.569		1.138		1.138		0.985		1.97			

DAS – Days After Sowing; RHA – Rice Husk Ash; SSB – Silicate Solubilizing Bacteria; ZSB – Zinc Solubilizing Bacteria;  
M – Microbes; S – Sources; L – Levels; SEd – Standard Error of difference; LSD – Least Significant Difference;  
SSB4 - *Bacillus altitudinis*; SSB7 - *Pseudomonas gessardii*; ZSB - *Bacillus aryabhatai*

**Table 3. Silicon Sources and bacterial Inoculants on Leaf Silicon (%) of Coriander**

Leaf Silicon (%) at 25 DAS																
Si Sources Si Levels (mg Si kg <sup>-1</sup> )	No Microbes				SSB4				SSB7				ZSB			
	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean
<b>0</b>	1.352	1.377	1.389	1.373	1.773	1.784	1.793	1.783	1.543	1.528	1.536	1.536	1.401	1.392	1.399	1.397
<b>50</b>	1.689	1.521	1.887	1.699	2.223	1.854	2.908	2.328	2.011	1.612	2.541	2.055	1.733	1.559	1.918	1.737
<b>100</b>	1.742	1.592	1.992	1.775	2.612	2.344	3.384	2.780	2.403	1.666	2.772	2.280	1.934	1.623	2.063	1.873
<b>Mean</b>	1.594	1.497	1.756	1.616	2.203	1.994	2.695	2.297	1.986	1.602	2.283	1.957	1.689	1.525	1.793	1.669
<b>Factors</b>	<b>M</b>		<b>S</b>		<b>L</b>		<b>M x S</b>		<b>M x L</b>		<b>S x L</b>		<b>M x S x L</b>			
<b>SEd</b>	0.032		0.027		0.027		0.055		0.055		0.048		0.095			
<b>LSD (P=0.05)</b>	0.063		0.055		0.055		0.109		0.109		0.095		0.189			
Leaf Silicon (%) at 50 DAS																
Si Sources Si Levels (mg Si kg <sup>-1</sup> )	No Microbes				SSB4				SSB7				ZSB			
	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean
<b>0</b>	2.026	2.012	2.030	2.023	2.591	2.608	2.620	2.606	2.255	2.233	2.243	2.244	2.046	2.030	2.044	2.040
<b>50</b>	2.469	2.223	2.756	2.483	3.247	2.708	3.610	3.189	2.938	2.356	3.296	2.863	2.533	2.278	2.803	2.538
<b>100</b>	2.545	2.326	2.911	2.594	3.817	3.424	4.112	3.784	3.358	2.435	3.561	3.118	2.827	2.372	3.014	2.738
<b>Mean</b>	2.347	2.187	2.566	2.367	3.219	2.913	3.447	3.193	2.850	2.341	3.033	2.742	2.469	2.227	2.620	2.439
<b>Factors</b>	<b>M</b>		<b>S</b>		<b>L</b>		<b>M x S</b>		<b>M x L</b>		<b>S x L</b>		<b>M x S x L</b>			
<b>SEd</b>	0.047		0.040		0.040		0.081		0.081		0.070		0.140			
<b>LSD (P=0.05)</b>	0.093		0.081		0.081		0.161		0.161		0.140		0.279			

DAS – Days After Sowing; RHA – Rice Husk Ash; SSB – Silicate Solubilizing Bacteria; ZSB – Zinc Solubilizing Bacteria;  
M – Microbes; S – Sources; L – Levels; SEd – Standard Error of difference; LSD – Least Significant Difference;  
SSB4 - *Bacillus altitudinis*; SSB7 - *Pseudomonas gessardii*; ZSB - *Bacillus aryabhatai*

**Table 4. Silicon Sources and bacterial Inoculants on Leaf Yield (g plant<sup>-1</sup>) of Coriander**

Si Sources	No Microbes				SSB4				SSB7				ZSB			
	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean
<b>0</b>	4.351	4.321	4.360	4.344	5.565	5.600	5.626	5.597	4.844	4.796	4.818	4.819	4.394	4.360	4.390	4.381
<b>50</b>	5.302	4.775	5.920	5.332	6.974	5.816	7.669	6.820	6.309	5.060	7.129	6.166	5.440	4.891	6.019	5.450
<b>100</b>	5.466	4.995	6.252	5.571	7.563	6.512	8.237	7.437	6.841	5.228	7.392	6.487	6.071	5.094	6.473	5.879
<b>Mean</b>	5.040	4.697	5.511	5.082	6.701	5.976	7.177	6.618	5.998	5.028	6.446	5.824	5.302	4.782	5.627	5.237
<b>Factors</b>	<b>M</b>		<b>S</b>		<b>L</b>		<b>M x S</b>		<b>M x L</b>		<b>S x L</b>		<b>M x S x L</b>			
<b>SEd</b>	0.041		0.035		0.035		0.071		0.071		0.061		0.123			
<b>LSD (P=0.05)</b>	0.081		0.071		0.071		0.141		0.141		0.122		0.244			

*DAS – Days After Sowing; RHA – Rice Husk Ash; SSB – Silicate Solubilizing Bacteria; ZSB – Zinc Solubilizing Bacteria; M – Microbes; S – Sources; L – Levels; SEd – Standard Error of difference; LSD – Least Significant Difference; SSB4 - Bacillus altitudinis; SSB7 - Pseudomonas gessardii; ZSB - Bacillus aryabhatai*

### 3.5 Adsorbed Silicon in Soil

Adsorbed silicon in soil ranged from 12.162 to 27.097 mg kg<sup>-1</sup> at 25 DAS while at 50 DAS, it ranged from 17.789 to 39.634 mg kg<sup>-1</sup> (Table 5). The maximum Ad-Si value was shown at both stages by the application of Rice Husk Ash @ 100 mg Si kg<sup>-1</sup> with SSB4, whereas the control treatment without silicon source and without microbes displayed a minimum value. The mean Ad-Si at 25 DAS ranged from 13.220 to 19.484, 14.185 to 19.766, and 15.111 to 22.050 mg kg<sup>-1</sup> while at 50 DAS it was 19.337 to 25.166, 20.506 to 28.679, and 22.410 to 32.251 mg kg<sup>-1</sup> for MgSiO<sub>3</sub>, CaSiO<sub>3</sub>, and RHA, respectively. As the amount of silicon fertilizer increased, the Ad-Si continuously increased. The Ad-Si is raised significantly by the microbial strains in the sequence of without microbes < ZSB < SSB7 < SSB4 at both stages. The interaction effects were found to be significant at both the stages *i.e.*, 25 & 50 DAS. The higher Si uptake with the use of Si fertilizer might be attributed to the increased availability of Si in the soil and the improved root system, both of which might have encouraged the plant to absorb more Si from the soil solution [38]. The application of silicon improves plant growth by increasing soil silicon content and other nutrients as well [67,72]. SSB improves the rice biomass by improving the soil silicon content [12].

### 3.6 TSS

Total Soluble Sugar content of Coriander dry leaves ranged from 21.012 to 38.227 mg g<sup>-1</sup> DW (Table 6). Application of Rice Husk Ash @ 100 mg Si kg<sup>-1</sup> with SSB4 showed a maximum soluble sugar content while control or without microbes and without Si source displayed a minimum value. Silicon sources increased the total soluble sugar of coriander leaves which is in the sequence of MgSiO<sub>3</sub> < CaSiO<sub>3</sub> < RHA. TSS showed linear improvement with the increased Si levels. The mean TSS values ranged from 22.840 to 26.853 mg g<sup>-1</sup> DW, 23.907 to 30.370 mg g<sup>-1</sup> DW, and 25.486 to 32.273 mg g<sup>-1</sup> DW for MgSiO<sub>3</sub>, CaSiO<sub>3</sub>, and RHA respectively. TSS content was increased by the microbial strains in the sequence of without microbes < ZSB < SSB7 < SSB4. Afshari M *et al.*, 2021 [57] observed that the foliar application of silicon sources increased the total soluble sugar content in coriander leaves. Many findings showed that in sorghum [73], soybean [74], rice [75], cucumber [76], sugarcane [77], maize [78] pepper [79], and barley [80], an improvement in TSS content with the help of Si fertilizers. SSB4 dissolved more Si from the soil compared to SSB7 and ZSB which might have helped in improving the TSS content in coriander leaves compared to other microbial strains.

## 4. CONCLUSION

The present investigation clearly indicated the significant effect of integration of Si sources with microbial inoculants in sandy loam soil of low soil adsorbed Si status. The application of Si sources alone and in combination with SSB/ZSB enhanced the growth parameters in terms of plant height and SPAD index values, leaf Si content, coriander leaf yield, and leaf quality besides improving the soil fertility. The plant height increased with the stage of the crop due to the applied Si sources and SSB whereas the SPAD index values were higher at 25 DAS compared to 50 DAS. The leaf silicon ranged from 1.352 to 3.384 per cent at 25 DAS, whereas it varied from 2.012 to 4.112 per cent. The Coriander leaf yield ranged from 4.321 to 8.237 g plant<sup>-1</sup>. The application of RHA @ 100 mg Si kg<sup>-1</sup> with SSB4 recorded the highest leaf yield of 8.237 g plant<sup>-1</sup> followed by RHA @ 50 mg Si kg<sup>-1</sup> with SSB4. However, the latter produced a comparable leaf yield with CaSiO<sub>3</sub> @ 100 mg Si kg<sup>-1</sup>. A similar trend was clearly evident with soil adsorbed Si. The leaf yield increase over control for RHA @ 100 mg Si kg<sup>-1</sup> + SSB4 was the highest (88.92 %) followed by RHA @ 50 mg Si kg<sup>-1</sup> with

SSB4 (76.58 %) and  $\text{CaSiO}_3$  @ 100 mg Si  $\text{kg}^{-1}$  + SSB4 (73.82 %). The results indicated that the application of Si sources and

UNDER PEER REVIEW

**Table 5. Silicon Sources and bacterial Inoculants on Adsorbed Silicon ( $\text{mg kg}^{-1}$ ) in Soil**

Ad-Si at 25 DAS																
Si Sources Si Levels ( $\text{mg Si kg}^{-1}$ )	No Microbes				SSB4				SSB7				ZSB			
	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean
0	12.247	12.162	12.271	12.227	15.665	15.762	15.835	15.754	13.634	13.500	13.561	13.565	12.369	12.271	12.357	12.332
50	14.923	13.439	16.662	15.008	19.629	16.370	23.217	19.739	17.757	14.242	20.712	17.570	15.312	13.767	16.942	15.340
100	15.385	14.059	17.598	15.681	24.003	19.484	27.097	23.528	21.223	14.716	23.521	19.820	17.088	14.339	18.219	16.548
Mean	14.185	13.220	15.511	14.305	19.766	17.205	22.050	19.674	17.538	14.153	19.265	16.985	14.923	13.459	15.839	14.740
Factors	M		S		L		M x S		M x L		S x L		M x S x L			
SEd	0.119		0.103		0.103		0.206		0.206		0.178		0.356			
LSD (P=0.05)	0.237		0.205		0.205		0.410		0.410		0.355		0.710			
Ad-Si at 50 DAS																
Si Sources Si Levels ( $\text{mg Si kg}^{-1}$ )	No Microbes				SSB4				SSB7				ZSB			
	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean
0	17.914	17.789	17.949	17.884	22.912	23.055	23.161	23.043	19.941	19.746	19.835	19.841	17.896	17.949	18.074	17.973
50	21.102	19.657	23.541	21.433	28.711	23.944	33.959	28.872	25.972	20.831	30.295	25.699	22.396	20.137	24.780	22.438
100	22.503	20.564	25.741	22.936	34.412	28.498	39.634	34.181	31.042	21.525	34.404	28.990	24.994	20.973	26.648	24.205
Mean	20.506	19.337	22.410	20.751	28.679	25.166	32.251	28.699	25.652	20.700	28.178	24.843	21.762	19.686	23.167	21.539
Factors	M		S		L		M x S		M x L		S x L		M x S x L			
SEd	0.144		0.125		0.125		0.249		0.249		0.216		0.432			
LSD (P=0.05)	0.287		0.248		0.248		0.497		0.497		0.430		0.860			

DAS – Days After Sowing; RHA – Rice Husk Ash; SSB – Silicate Solubilizing Bacteria; ZSB – Zinc Solubilizing Bacteria;  
M – Microbes; S – Sources; L – Levels; SEd – Standard Error of difference; LSD – Least Significant Difference;  
SSB4 - *Bacillus altitudinis*; SSB7 - *Pseudomonas gessardii*; ZSB - *Bacillus aryabhatai*

**Table 6. Silicon Sources and bacterial Inoculants on Total Soluble Sugar (mg g<sup>-1</sup> DW) of Coriander**

Si Sources Si Levels (mg Si kg <sup>-1</sup> )	No Microbes				SSB4				SSB7				ZSB			
	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean	CaSiO <sub>3</sub>	MgSiO <sub>3</sub>	RHA	Mean
<b>0</b>	21.159	21.012	21.201	21.124	24.226	24.301	24.381	24.303	22.912	22.997	23.003	22.971	21.369	21.201	21.348	21.159
<b>50</b>	23.982	23.218	24.854	24.018	30.781	25.971	34.210	30.321	28.971	24.605	31.831	28.469	25.016	23.786	25.556	23.982
<b>100</b>	26.580	24.290	30.404	27.091	36.103	30.287	38.227	34.872	32.009	25.425	35.023	30.819	29.522	24.773	31.476	26.580
<b>Mean</b>	23.907	22.840	25.486	24.078	30.370	26.853	32.273	29.832	27.964	24.342	29.952	27.420	25.302	23.253	26.127	23.907
<b>Factors</b>	<b>M</b>		<b>S</b>		<b>L</b>		<b>M x S</b>		<b>M x L</b>		<b>S x L</b>		<b>M x S x L</b>			
<b>SEd</b>	0.181		0.157		0.157		0.313		0.313		0.271		0.542			
<b>LSD (P=0.05)</b>	0.361		0.312		0.312		0.624		0.624		0.541		1.082			

*DAS – Days After Sowing; DW – Dry Weight; RHA – Rice Husk Ash; SSB – Silicate Solubilizing Bacteria; ZSB – Zinc Solubilizing Bacteria; M – Microbes; S – Sources; L – Levels; SEd – Standard Error of difference; LSD – Least Significant Difference; SSB4 - Bacillus altitudinis; SSB7 - Pseudomonas gessardii; ZSB - Bacillus aryabhatai*

SSB was found to be effective in enriching the Si content of coriander leaf, besides increasing the yield.

## REFERENCES

1. Ma JF, Takahashi E. Soil, fertilizer, and plant silicon research in Japan. Elsevier; 2002 Aug 9.
2. Epstein E. Silicon. Annual review of plant biology. 1999;50:641.
3. Epstein E. The anomaly of silicon in plant biology. Proceedings of the National Academy of Sciences. 1994 Jan 4;91(1):11-7.
4. Mateos-Naranjo E, Gallé A, Florez-Sarasa I, Perdomo JA, Galmés J, Ribas-Carbó M, Flexas J. Assessment of the role of silicon in the Cu-tolerance of the C4 grass *Spartina densiflora*. Journal of Plant Physiology. 2015 Apr 15;178:74-83.
5. Kim YH, Khan AL, Waqas M, Lee IJ. Silicon regulates antioxidant activities of crop plants under abiotic-induced oxidative stress: a review. Frontiers in Plant Science. 2017 Apr 6;8:510.
6. Todorova D, Sergiev I, Moskova I, Katerova Z, Georgieva N, Alexieva V, Brambilla I, Mapelli S. Biochemical responses of triticale plants treated with UV-B irradiation and nutrient solution enriched with humic acids. Turkish Journal of Botany. 2014;38(4):747-53.
7. Abdel-Haliem ME, Hegazy HS, Hassan NS, Naguib DM. Effect of silica ions and nano silica on rice plants under salinity stress. Ecological Engineering. 2017 Feb 1;99:282-9.
8. Ma JF, Yamaji N. Functions and transport of silicon in plants. Cellular and molecular life sciences. 2008 Oct;65(19):3049-57.
9. Vatanserver R, Ozyigit II, Filiz E. Essential and beneficial trace elements in plants, and their transport in roots: a review. Applied Biochemistry and Biotechnology. 2017 Jan;181(1):464-82.
10. Schaller J, Puppe D, Kaczorek D, Ellerbrock R, Sommer M. Silicon Cycling in Soils Revisited. Plants 2021, 10, 295.
11. Asgari F, Majd A, Jonoubi P, Najafi F. Effects of silicon nanoparticles on molecular, chemical, structural and ultrastructural characteristics of oat (*Avena sativa* L.). Plant Physiology and Biochemistry. 2018 Jun 1;127:152-60.
12. Meena VD, Dotaniya ML, Coumar V, Rajendiran S, Kundu S, Subba Rao A. A case for silicon fertilization to improve crop yields in tropical soils. Proceedings of the National Academy of Sciences, India Section B: Biological Sciences. 2014 Sep;84(3):505-18.
13. Sattar A, Cheema MA, Sher A, Ijaz M, Ul-Allah S, Nawaz A, Abbas T, Ali Q. Physiological and biochemical attributes of bread wheat (*Triticum aestivum* L.) seedlings are influenced by foliar application of silicon and selenium under water deficit. Acta Physiologiae Plantarum. 2019 Aug;41(8):1-1.
14. Farouk S, Omar MM. Sweet basil growth, physiological and ultrastructural modification, and oxidative defense system under water deficit and silicon forms treatment. Journal of Plant Growth Regulation. 2020 Sep;39(3):1307-31.
15. Tubana BS, Babu T, Datnoff LE. A review of silicon in soils and plants and its role in US agriculture: history and future perspectives. Soil science. 2016 Sep 1;181(9/10):393-411.
16. Raghavan S. Handbook of spices, seasonings, and flavorings. CRC press; 2006 Oct 23.

17. Sachan AK, Kumar S, Kumari K, Singh D. Medicinal uses of spices used in our traditional culture: Worldwide. *Journal of Medicinal Plants Studies*. 2018;6(3):116-22.
18. Delaquis PJ, Stanich K, Girard B, Mazza G. Antimicrobial activity of individual and mixed fractions of dill, cilantro, coriander and eucalyptus essential oils. *International journal of food microbiology*. 2002 Mar 25;74(1-2):101-9.
19. Kubo I, Fujita KI, Kubo A, Nihei KI, Ogura T. Antibacterial activity of coriander volatile compounds against *Salmonella choleraesuis*. *Journal of agricultural and food chemistry*. 2004 Jun 2;52(11):3329-32.
20. Mohanalakshmi M, Boomiga M, Gowtham T. Effect of the season and growing condition on yield and quality parameters of coriander (*Coriandrum sativum* L.). *International Journal of Chemical Studies* 2019; 7(3): 2989-2993
21. Verma A, Pandeya SN, Yadav SK, Singh S, Soni P. A review on *Coriandrum sativum* (Linn.): An ayurvedic medicinal herb of happiness. *J Adv Pharm Healthc Res*. 2011;1(3):28-48.
22. Spices Board, India & Ministry of Agriculture and Farmers Welfare, Govt. of India, 2021; <http://www.indianspices.com/>
23. Bhuiyan MN, Begum J, Sultana M. Chemical composition of leaf and seed essential oil of *Coriandrum sativum* L. from Bangladesh. *||| Bangladesh Journal of Pharmacology|||*. 2009 Dec 17;4(2):150-3.
24. Neffati M, Marzouk B. Changes in essential oil and fatty acid composition in coriander (*Coriandrum sativum* L.) leaves under saline conditions. *Industrial crops and products*. 2008 Sep 1;28(2):137-42.
25. Kalembe DA, Kunicka A. Antibacterial and antifungal properties of essential oils. *Current medicinal chemistry*. 2003 May 1;10(10):813-29.
26. Dimri BP, Khan MN, Narayana MR. Some promising selections of Bulgarian coriander (*Coriandrum sativum* L.) for seed and essential oil with a note on cultivation and distillation of oil. *Indian Perfumer*. 1976;20(1):13-21.
27. Maleva M, Borisova G, Koshcheeva O, Sinenko O. Biofertilizer based on silicate solubilizing bacteria improves photosynthetic function of *Brassica juncea*. *AGROFOR International Journal*. 2017;2(3):13-9.
28. Goteti PK, Emmanuel LD, Desai S, Shaik MH. Prospective zinc solubilising bacteria for enhanced nutrient uptake and growth promotion in maize (*Zea mays* L.). *International journal of microbiology*. 2013 Oct;2013.
29. Sunithakumari K, Devi SP, Vasandha S. Zinc solubilizing bacterial isolates from the agricultural fields of Coimbatore, Tamil Nadu, India. *Current Science*. 2016 Jan 25:196-205.
30. Rajkumar M, Freitas H. Effects of inoculation of plant-growth promoting bacteria on Ni uptake by Indian mustard. *Bioresource Technology*. 2008 Jun 1;99(9):3491-8.
31. Narayanaswamy C, Prakash NB. Calibration and categorization of plant available silicon in rice soils of South India. *Journal of plant nutrition*. 2009 Jul 22;32(8):1237-54.
32. Ning D, Song A, Fan F, Li Z, Liang Y. Effects of slag-based silicon fertilizer on rice growth and brown-spot resistance. *Plos one*. 2014 Jul 18;9(7):e102681
33. Ma JF, Goto S, Tamai K, Ichii M. Role of root hairs and lateral roots in silicon uptake by rice. *Plant physiology*. 2001 Dec 1;127(4):1773-80.
34. Korndörfer GH, Lepesch I. Effect of silicon on plant growth and crop yield. In *Studies in plant science* 2001 Jan 1 (Vol. 8, pp. 133-147). Elsevier.
35. Yemm EW, Willis A. The estimation of carbohydrates in plant extracts by anthrone. *Biochemical journal*. 1954 Jul;57(3):508.
36. Gomez KA, Gomez AA. *Statistical procedures for agricultural research*. John Wiley & sons; 1984 Feb 17.

37. Ahmad A, Afzal M, Ahmad AU, Tahir M. Effect of foliar application of silicon on yield and quality of rice (*Oryza sativa* L.). *Cercetari agronomice in Moldova*. 2013;46(3):21-8.
38. Pati S, Pal B, Badole S, Hazra GC, Mandal B. Effect of silicon fertilization on growth, yield, and nutrient uptake of rice. *Communications in Soil Science and Plant Analysis*. 2016 Feb 4;47(3):284-90.
39. Yoshida S, Navasero SA, Ramirez EA. Effects of silica and nitrogen supply on some leaf characters of the rice plant. *Plant and soil*. 1969 Aug;31(1):48-56
40. Elawad SH, Street JJ, Gascho GJ. Response of Sugarcane to Silicate Source and Rate. II. Leaf Freckling and Nutrient Content 1. *Agronomy Journal*. 1982 May;74(3):484-7
41. Gascho GJ. Silicon sources for agriculture. In *Studies in plant science* 2001 Jan 1 (Vol. 8, pp. 197-207). Elsevier.
42. Xie Z, Song F, Xu H, Shao H, Song R. Effects of silicon on photosynthetic characteristics of maize (*Zea mays* L.) on alluvial soil. *The Scientific World Journal*. 2014 Oct;2014.
43. Bokhtiar SM, Huang HR, Li YR, Dalvi VA. Effects of silicon on yield contributing parameters and its accumulation in abaxial epidermis of sugarcane leaf blades using energy dispersive x-ray analysis. *Journal of plant nutrition*. 2012 May 15;35(8):1255-75.
44. Soundararajan P, Sivanesan I, Jana S, Jeong BR. Influence of silicon supplementation on the growth and tolerance to high temperature in *Salvia splendens*. *Horticulture, Environment, and Biotechnology*. 2014 Aug;55(4):271-9.
45. Jones LH, Milne AA, Wadham SM. Studies of Silica in the Oat Plant: II. Distribution of the Silica in the Plant. *Plant and Soil*. 1963 Jun 1:358-71.
46. Sarto MV, do Carmo Lana M, Rampim L, Rosset JS, Sarto JR, Bassegio D. Effects of calcium and magnesium silicate on the absorption of silicon and nutrients in wheat. *Semina: Ciências Agrárias*. 2019;40(1):67-80.
47. Sistani KR, Savant NK, Reddy KC. Effect of rice hull ash silicon on rice seedling growth. *Journal of Plant Nutrition*. 1997 Jan 1;20(1):195-201.
48. Ma J, Takahashi E. Availability of rice straw Si to rice plants. *Soil Science and Plant Nutrition*. 1991 Mar 1;37(1):111-6.
49. Sumida H, Ohyama N. The effects of application of organic matters and calcium silicate on silica uptake by rice plant [*Oryza sativa*]. *Japanese Journal of Soil Science and Plant Nutrition (Japan)*. 1991.
50. Yu X, Bell PF. Nutrient deficiency symptoms and boron uptake mechanisms of rice. *Journal of plant nutrition*. 1998 Oct 1;21(10):2077-88
51. Qin J, Impa SM, Tang Q, Yang S, Yang J, Tao Y, Jagadish KS. Integrated nutrient, water and other agronomic options to enhance rice grain yield and N use efficiency in double-season rice crop. *Field Crops Research*. 2013 Jul 1;148:15-23.
52. Yanai J, Taniguchi H, Nakao A. Evaluation of available silicon content and its determining factors of agricultural soils in Japan. *Soil Science and Plant Nutrition*. 2016 Nov 1;62(5-6):511-8.
53. Peera SP, Balasubramaniam P, Mahendran PP. Effect of silicate solubilizing bacteria and fly ash on silicon uptake and yield of rice under lowland ecosystem. *Journal of Applied and Natural Science*. 2016 Mar 1;8(1):55-9.
54. Etesami H, Jeong BR. Biodissolution of silica by rhizospheric silicate-solubilizing bacteria. In *Silicon and Nano-silicon in Environmental Stress Management and Crop Quality Improvement* 2022 Jan 1 (pp. 265-276). Academic Press.
55. Deren CW, Datnoff LE, Snyder GH, Martin FG. Silicon concentration, disease response, and yield components of rice genotypes grown on flooded organic histosols. *Crop Science*. 1994 May;34(3):733-7

56. Vasanthi N, Saleena LM, Raj SA. Silicon in day today life. *World Appl Sci J*. 2012;17(11):1425-40
57. Afshari M, Pazoki A, Sadeghipour O. Foliar-applied Silicon and its Nanoparticles Stimulate Physio-chemical Changes to Improve Growth, Yield and Active Constituents of Coriander (*Coriandrum Sativum* L.) Essential Oil Under Different Irrigation Regimes. *Silicon*. 2021 Nov;13(11):4177-88.
58. Hassanein RA, Hussein OS, Abdelkader AF, Farag IA, Hassan YE, Ibrahim M. Metabolic activities and molecular investigations of the ameliorative impact of some growth biostimulators on chilling-stressed coriander (*Coriandrum sativum* L.) plant. *BMC plant biology*. 2021 Dec;21(1):1-23.
59. Lavinsky AO, Detmann KC, Reis JV, Ávila RT, Sanglard ML, Pereira LF, Sanglard LM, Rodrigues FA, Araújo WL, DaMatta FM. Silicon improves rice grain yield and photosynthesis specifically when supplied during the reproductive growth stage. *Journal of Plant Physiology*. 2016 Nov 1;206:125-32.
60. Jawahar S, Vijayakumar D, Bommera R, Jain N. Effect of silixol granules on growth and yield of rice. *Int J Curr Res Aca Rev*. 2015;3(5):74-80.
61. Liang YC, Ma TS, Li FJ, Feng YJ. Silicon availability and response of rice and wheat to silicon in calcareous soils. *Communications in Soil Science and Plant Analysis*. 1994 Aug 1;25(13-14):2285-97.
62. Gerami M, Fallah A, Moghadam MR. Study of potassium and sodium silicate on the morphological and chlorophyll content on the rice plant in pot experiment (*Oryza sativa* L.). *International Journal of Agriculture and Crop Sciences (IJACS)*. 2012;4(10):658-61.
63. Miyake Y, Takahashi E. Effect of silicon on the growth of solution-cultured cucumber plant. *Soil Science and Plant Nutrition*. 1983 Mar 1;29(1):71-83.
64. Savvas D, Giotis D, Chatzieustratiou E, Bakea M, Patakioutas G. Silicon supply in soilless cultivations of zucchini alleviates stress induced by salinity and powdery mildew infections. *Environmental and experimental botany*. 2009 Jan 1;65(1):11-7.
65. Zuccarini P. Effects of silicon on photosynthesis, water relations and nutrient uptake of *Phaseolus vulgaris* under NaCl stress. *Biologia Plantarum*. 2008 Mar;52(1):157-60.
66. Toresano-Sánchez F, Valverde-García A, Camacho-Ferre F. Effect of the application of silicon hydroxide on yield and quality of cherry tomato. *Journal of plant nutrition*. 2012 Mar 1;35(4):567-90.
67. Singh KK, Singh K, Singh R, Singh Y, Singh CS. Response of nitrogen and silicon levels on growth, yield and nutrient uptake of rice (*Oryza sativa* L.). *Oryza*. 2006;43(3):220.
68. Ma J, Nishimura K, Takahashi E. Effect of silicon on the growth of rice plant at different growth stages. *Soil Science and Plant Nutrition*. 1989 Sep 1;35(3):347-56.
69. Prakash NB, Chandrashekar N, Mahendra C, Patil SU, Thippeshappa GN, Laane HM. Effect of foliar spray of soluble silicic acid on growth and yield parameters of wetland rice in hilly and coastal zone soils of Karnataka, South India. *Journal of plant nutrition*. 2011 Sep 1;34(12):1883-93.
70. Ross L, Nababsing P, Cheong YW. Residual effect of calcium silicate applied to sugarcane soils. In *Proceedings of the... congress 1974*.
71. Rosas SB, Avanzini G, Carlier E, Pasluosta C, Pastor N, Rovera M. Root colonization and growth promotion of wheat and maize by *Pseudomonas aurantiaca* SR1. *Soil Biology and Biochemistry*. 2009 Sep 1;41(9):1802-6
72. Patil AA, Durgude AG, Pharande AL. Effect of silicon application along with chemical fertilizers on nutrient uptake and nutrient availability for rice plants. *IJCS*. 2018;6(1):260-6.

73. Sonobe K, Hattori T, An P, Tsuji W, Eneji AE, Kobayashi S, Kawamura Y, Tanaka K, Inanaga S. Effect of silicon application on sorghum root responses to water stress. *Journal of Plant Nutrition*. 2010 Dec 1;34(1):71-82
74. Hussain S, Mumtaz M, Manzoor S, Shuxian L, Ahmed I, Skalicky M, Brestic M, Rastogi A, Ulhassan Z, Shafiq I, Allakhverdiev SI. Foliar application of silicon improves growth of soybean by enhancing carbon metabolism under shading conditions. *Plant Physiology and Biochemistry*. 2021 Feb 1;159:43-52.
75. Meharg C, Meharg AA. Silicon, the silver bullet for mitigating biotic and abiotic stress, and improving grain quality, in rice?. *Environmental and Experimental Botany*. 2015 Dec 1;120:8-17.
76. Zhu YX, Xu XB, Hu YH, Han WH, Yin JL, Li HL, Gong HJ. Silicon improves salt tolerance by increasing root water uptake in *Cucumis sativus* L. *Plant Cell Reports*. 2015 Sep;34(9):1629-46.
77. Oliva KM, da Silva FB, Araújo PR, de Oliveira EC, do Nascimento CW. Amorphous silica-based fertilizer increases stalks and sugar yield and resistance to stalk borer in sugarcane grown under field conditions. *Journal of Soil Science and Plant Nutrition*. 2021 Sep;21(3):2518-29.
78. Abdel Latef AA, Tran LS. Impacts of priming with silicon on the growth and tolerance of maize plants to alkaline stress. *Frontiers in Plant Science*. 2016 Mar 10;7:243.
79. Trejo-Téllez LI, García-Jiménez A, Escobar-Sepúlveda HF, Ramírez-Olvera SM, Bello-Bello JJ, Gómez-Merino FC. Silicon induces hormetic dose-response effects on growth and concentrations of chlorophylls, amino acids and sugars in pepper plants during the early developmental stage. *PeerJ*. 2020 Jun 9;8:e9224.
80. Benslimma W, Zorrig W, Bagues M, Abdelly C, Hafsi C. Silicon mitigates potassium deficiency in *Hordeum vulgare* by improving growth and photosynthetic activity but not through polyphenol accumulation and the related antioxidant potential. *Plant and Soil*. 2021 Oct 26: