

**Comparative Efficacy of Two Antibiotics for the Management of Secondary Bacterial Infection in Goats Clinically Affected by *Peste des Petits Ruminants***

---

**ABSTRACT**

**Aims:** *Peste des petits Ruminants* (PPR) is a major viral disease that poses a challenge to small ruminant farming. Its natural occurrence has been complicated by secondary bacterial infection which has led to increase in morbidity and mortality rates. This study reports the management outcome of natural PPR-infected goats using two types of antibiotics in Nsukka metropolis of Enugu State Nigeria.

**Methodology:** Goats (N=24) were confirmed to be suffering from PPR based on clinical signs and using polymerase chain reaction (PCR). The animals were grouped into two. Group A was treated with 20% oxytetracycline (N= 10) and group B with procaine penicillin and streptomycin combination (penstrep) (N= 14) injection. Clinical signs, recovery and survivability, temperature, haematology [Packed cell volume (PCV); haemoglobin concentration (Hb conc); red blood cell (RBC) and white blood cell (WBC) counts] and some serum biochemical profiles [alanine aminotransferase (ALT); aspartate transaminase (AST); total protein (TP); Albumin; urea; creatinine; potassium and sodium] were used to assess the efficacy of the antibiotics using standard techniques.

**Results:** The mean temperature, RBC, WBC and urea values of the PPR-infected goats were above their reference ranges, mean albumin values were below the reference range while mean Hb conc, PCV, AST, ALT, TP and creatinine values were within their reference range before the commencement of treatment. Following treatment, clinical signs cleared in 20% and 35.7% of the goats treated with oxytetracycline and penstrep respectively. Death was recorded in 20% and 15% of goats treated with oxytetracycline and penstrep respectively before the end of treatment. Penstrep-treated group showed improvement in their haematological profile.

**Conclusion:** Based on our findings, the use of penstrep in the management of PPR in goats gave a better result.

*Keywords: Goats; Haemo-biochemical profile; Oxytetracycline; Peste des petits ruminants; Procaine penicillin and streptomycin combination*

**1. INTRODUCTION**

*Peste des petits ruminants* (PPR) has been identified as one of the major diseases that pose a serious challenge to small ruminant farming [1]. This infection is acute, highly contagious, trans-boundary viral disease of sheep and goats, causing devastating disease that poses a threat to livestock production in many developing countries. The disease is enzootic in several countries of West Africa and produces variable rates of morbidity and mortality that

can reach 100% and 90%, respectively [2]. For example, due to PPR outbreaks, the economic loss in Nigeria has been estimated to be 6.8 billion of Naira (NGN) [3]. The aetiologic agent of PPR- *Small Ruminant Morbillivirus* (SRM), is a single-stranded ribosomal nucleic acid (RNA) virus which is a member of the *Morbillivirus* genus of the Paramyxoviridae family [4]. Infected animals present clinical signs such as fever, conjunctivitis, ocular discharges, encrustation in the media canthus, mucopurulent nasal discharges, ulcerative stomatitis, salivation, pneumonia, coughing, sneezing, profuse diarrhoea, dehydration with sunken eyes, emaciation and inflammation of the mucous membrane of the respiratory and digestive tracts. This infection of small ruminants can only be managed symptomatically as it has no specific treatment. Management using broad-spectrum antibiotics, antihistaminic, fluid therapy, intestinal sedative and multivitamin have been reported [5]. The disease in its natural occurrence has been complicated by secondary bacterial infection, which has led to increase in morbidity and mortality rates [2,6] with the main pathologic lesions observed in the digestive and respiratory systems. This paper, therefore, compares the efficacy of two antibiotics in the management of secondary bacterial infection in goats clinically affected by PPR in Nsukka metropolis, Enugu state Nigeria.

## **2. MATERIAL AND METHODS**

### **2.1 Experimental Animals**

A total of 74 goats showing clinical signs suggestive of PPR (e.g. ocular and nasal discharges, encrustation in the media canthus, ulcerative stomatitis, coughing, difficulty/laboured breathing, diarrhoea) were identified (by clinical examination as described by [6,7]) in households in Nsukka metropolis of Enugu State, Nigeria from November 2019 - March 2020. A total of 58 goats (comprising 22 males and 36 females) were purchased from the owners (convenience sampling based on willingness to sell their sick goat) and used for this study. The infected goats were not vaccinated as per the information obtained from the owner.

The goats were kept in a fly-proof well ventilated house, fed cut and carry grasses [elephant grass (*Pennisetum purpureum*), guinea grass (*Panicum maximum*), and palm font] as is the usual practice in this eco-zone and clean water provided ad libitum. Each goat was identified using neck tag (numbered 1-58). They were handled in compliance with the guidelines for the humane treatments of animals during experimentation in the University of Nigeria. The faecal dropping was removed on a daily basis and the floor disinfected with common disinfectants (phenol and detergents) and exposed to 2% sodium hydroxide to kill the virus present in the faeces [8]. The goats were grouped systematically as those with odd numbered neck tags were grouped as A (treated with Long acting oxytetracycline injection) while those with even numbered neck tags were grouped as B (treated with Procaine Penicillin and Dihydrostreptomycin combination)

Blood (5 ml) through venipuncture and nasal swab samples were collected from each goat before the commencement of treatment and were properly labelled. Treatment was done using 20% oxytetracycline injection (Kepro, Netherland) (20mg/kg body weight intramuscularly, 2<sup>nd</sup> dose given after 48 hours) and Procaine Penicillin and Dihydrostreptomycin combination injection (Penstrep) (Kepro, Netherland) (1ml/25kg body weight intramuscularly for 3 days). On days 3 and 7 post-commencement of treatment, blood samples (5ml) were again collected from the jugular vein; temperature and clinical observation were taken and documented.

Blood samples (4ml) for biochemistry were allowed to clot and then centrifuged at 1500 x g for 10 minutes. Sera harvested were stored at -20°C, while blood samples (1ml) for haematology were put into sample tubes containing ethylenediamine tetraacetic acid (EDTA) as an anticoagulant. All the samples were taken to the Department of Veterinary Medicine Laboratory, University of Nigeria Nsukka in plastic coolers for processing and analysis. The nasal swabs and sera samples were packed in dry ice and transported to Biotechnology Centre and the Biochemical Division, National Veterinary Research (NVRI) Institute, Vom, Jos Plateau State, Nigeria respectively for PPRV detection and biochemical analysis.

## **2.2 PPR Virus Detection**

The nasal swabs were evaluated for PPRV by RT-PCR assay. RNA extraction using QIAamp Viral RNA Mini kit (Qiagen, Hilden, Germany), RT-PCR using QIAGEN® OneStep Ahead RT-PCR Kit (Qiagen, Hilden, Germany) and analysis of amplicon were done using standard procedures as described by Chukwudi *et al.* [9].

## **2.3 Parameters for Assessing the Efficacy of the Drugs**

Rectal temperature, clinical signs and survivability were documented. Packed cell volume (PCV), Haemoglobin concentration, Red blood cell (RBC) count, and White blood cell (WBC) count as described by Coles [10] and Schalm *et al.* [11] were determined using standard laboratory techniques. Total protein, albumin, urea, creatinine, alanine aminotransferase (ALT), aspartate transaminase (AST) and serum electrolytes such as Potassium (K) and sodium (Na) were determined using standard Randox® test kit according to manufacturer's recommendation.

## **2.4 Statistical Analysis**

Data generated from temperature, haematology and serum biochemistry were analysed using One-way analysis of variance (ANOVA) (monitoring the progress of treatment) using SPSS version 16 software package. The means were separated using Duncan's multiple range test [12] at 5% level of probability.

# **3. RESULTS AND DISCUSSION**

## **3.1 PCR**

Out of the 58 treated goats that were showing clinical signs suggestive of PPR, only the nasal swabs of 24 goats were positive for PPRV by RT-PCR assay (Fig. 1).




Fig 1. Representative gel picture of samples analysed by RT-PCR.

Lane M= 100bp DNA molecular weight maker, Lane EC, NC and PC are extraction, negative and positive control respectively, Lane 1-25 are PCR product for representative samples.

### 3.2 Clinical Signs, Recovery and Survivability

The varying clinical signs displayed by the 24 goats before and after treatment was any or a combination of the following: coughing, mucopurulent nasal discharges, ocular discharges, rough hair coat, stomatitis, weak, dull, emaciation and others (Table 1).

Following treatment, only 7 goats [20% (n =2) for Oxytet treated group; 35.7% (n = 5) for Penstrep treated group] had their clinical signs completely cleared. Death was recorded in 4 goats [20% (n =2) for Oxytet treated group; 15% (n =2) for Penstrep treated group] before the end of the treatment.

**Table 1. Details of clinical signs displayed by the treated 24 goats before and after commencement of treatment**

<b>Oxytet Group (A)</b>	<b>Goat S/NO</b>	<b>Clinical signs before treatment</b>	<b>Clinical signs after treatment</b>	<b>Response</b>
	5	Dia, Cou, Od, MpNd,	Nd,	
	9	Cou, Nd,	No obvious signs	Cleared
	13	Nd, Cou. Dia, Du, RHC, Wk, Us	Nd	
	19	Nd, Od, Cou. Du, Ema, RHC, Wk	Nd, Ema,	
	17	Mpnd, Od, Cou	No obvious sign	Cleared
	27	Nd, Od, Dia, Us	Nd, Dia,	
	29	Nd, Od, Cou. Dia, Du, RHC	Nd, Dia,	
	33	Nd, Cou, Du, Dia	Death	
	35	MpNd, Dia, Cou	Nd, Dia wk	
	51	Nd, Od, Cou. Du, Wk	Death	
<b>Pentrep group (B)</b>				
	4	Severe MpNd, Du	Nd,	
	14	Nd, Dia, Us, Cou. Du, RHC	Nd, Dia (reduced)	
	28	MpNd, Cou. Dia, Du, RHC, Ema, Wk, Se	Nd, Dia (reduced), Ema	
	30	Nd, Cou. Du, RHC, Ema	RHC, Ema	

32	Nd, Od, Cou. Du, RHC,	No obvious sign	Cleared
36	MpNd, Cou. Dia, Us, Du, RHC, Ema	Nd, Dia (reduced)	
38	Nd, Od, Cou. Du,	Nd	
42	MpNd, Us, Cou, Du	Death	
44	Severe MpNd, Dia, Du	Death	
46	Nd, Cou, Du	No obvious sign	Cleared
48	Nd, Du, Cou	No obvious sign	Cleared
50	MpNd, Cou, Dia	Nd	
52	MpNd, Cou	No obvious sign	Cleared
56	Cou, Nd, Du	No obvious sign	Cleared

---

Clinical signs: Nd- nasal discharge; Od- ocular discharge; Cou- coughing; Ema- emaciated; RHC- rough hair coat; Wk- weak; Du- dull; Dia- diarrhea; MpNd- mucopurulent nasal discharge; Us- Ulcerative stomatitis

### 3.3 Temperature

There was an increase in the mean temperature of both groups (oxytetracycline-treated and penstrep-treated groups) before the commencement of treatment (above the reference value) (Table 2). By day 3 post-commencement, a significant decrease ( $P < .05$ ) was observed in both groups. On day 7 post-commencement of treatment, a significant increase ( $P < .05$ ) was recorded in the oxytetracycline-treated group compared to that of day 3 post commencement of treatment.

**Table 2. Temperature (°C) of goats naturally infected with PPR and treated with oxytetracycline and penstrep**

Groups	Day 0	Day 3 post-treatment	Day 7 post-treatment
A- Oxytet treated	40.95 ± 0.55 <sup>b</sup>	38.60 ± 0.10 <sup>a</sup>	39.70 ± 0.40 <sup>b</sup>
B- Penstrep treated	40.70 ± 0.20 <sup>b</sup>	38.20 ± 0.25 <sup>a</sup>	38.8 ± 1.70 <sup>a</sup>
Reference value	38.5-39.7°C [13]		

Superscript a,b,c indicates significance row-wise

### 3.4 Haematological Analysis

#### 3.4.1 Packed cell volume (PCV)

There were no significant differences ( $P > .05$ ) in the mean PCV of oxytet-treated group all through the duration of treatment, while there were significant decreases ( $P < .05$ ) on days 3 and 7 post-commencement of treatment in the penstrep-treated group when compared to their initial value on day 0 before the commencement of treatment (Table 3). The mean PCV values of both groups all through the experiment/treatment and monitoring were within the reference range.

#### 3.4.2 Haemoglobin concentration (Hb conc)

There were no significant differences ( $P > .05$ ) in the mean Hb conc of oxytet-treated group all through the duration of treatment, while there was a significant decrease ( $P < .05$ ) on day 3 post-commencement of treatment in the penstrep-treated group when compared to their initial values before the commencement of treatment (Table 3). The mean Hb conc values of both groups all through the experiment/treatment and monitoring were within the reference range.

#### 3.4.3 Red blood cell count (RBC)

There was an increase in the mean RBC values (above the reference value) of both groups (oxytetracycline-treated and penstrep-treated groups) before the commencement of treatment (Table 3). By day 3 post-commencement of treatment, a significant decrease ( $P < .05$ ) in RBC values was observed in both groups.

#### 3.4.4 White blood cell count (WBC)

There was a slight increase in the mean WBC values (above the reference value) of both groups (oxytetracycline-treated and penstrep-treated groups) before the commencement of treatment (Table 3). There were no significant differences ( $P > .05$ ) in the mean WBC values of both groups all through the duration of treatment.

**Table 3. Haematology of goats naturally infected with PPR and treated with oxytetracycline and penstrep**

Group/parameter	Day 0	Day 3 post-treatment	Day 7 post-treatment
<b>PCV (%)</b>			
A- Oxytet treated	29.00 ± 3.06 <sup>a</sup>	28.33 ± 4.91 <sup>a</sup>	23.5 ± 0.50 <sup>a</sup>
B- Penstrep treated	29.50 ± 2.50 <sup>a</sup>	23.00 ± 2.00 <sup>c</sup>	26.00 ± 0.15 <sup>b</sup>
Reference value	22-35% [14]		
<b>HB conc (g/dl)</b>			
A- Oxytet treated	11.30 ± 0.60 <sup>a</sup>	9.30 ± 1.48 <sup>a</sup>	11.60 ± 0.90 <sup>a</sup>
B- Penstrep treated	10.85 ± 0.95 <sup>a</sup>	7.90 ± 0.7 <sup>b</sup>	13.05 ± 2.05 <sup>a</sup>
Reference value	7-15 g/dl [14]		
<b>RBC (x 10<sup>6</sup>/mm<sup>3</sup>)</b>			
A- Oxytet treated	16.23 ± 0.54 <sup>a</sup>	8.23 ± 1.25 <sup>b</sup>	12.54 ± 2.33 <sup>c</sup>
B- Penstrep treated	14.96 ± 0.59 <sup>a</sup>	6.55 ± 0.55 <sup>b</sup>	15.38 ± 3.18 <sup>a</sup>
Reference value	9.2-13.5 x 10 <sup>6</sup> /mm <sup>3</sup> [14]		
<b>WBC (x 10<sup>3</sup>/mm<sup>3</sup>)</b>			
A- Oxytet treated	15.87 ± 7.57 <sup>a</sup>	14.10 ± 2.00 <sup>a</sup>	17.68 ± 0.48 <sup>a</sup>
B- Penstrep treated	14.45 ± 3.10 <sup>a</sup>	13.34 ± 3.99 <sup>a</sup>	15.15 ± 1.8 <sup>a</sup>
Reference value	6-13 x10 <sup>3</sup> /mm <sup>3</sup> [15]		

Superscript a,b,c indicates significance row-wise

### 3.5 Biochemical Profile

#### 3.5.1 Aspartate transaminase (AST)

There were no significant differences ( $P > .05$ ) in the mean AST values of both groups all through the duration of treatment (Table 4). The AST values all through the experimental/ treatment and monitoring period were within the reference range.

#### 3.5.2 Alanine aminotransferase (ALT)

There were no significant differences ( $P > .05$ ) in the mean ALT values of both groups all through the duration of treatment (Table 4). The ALT values all through the experimental/ treatment and monitoring period were within the reference range.

#### 3.5.3 Total protein (TP)

There were no significant differences ( $P > .05$ ) in the mean TP values of both groups all through the duration of treatment (Table 4). The TP values all through the experimental/ treatment and monitoring period were within the reference range.

#### 3.5.4 Albumin

There was a significant decrease ( $P < .05$ ) in the mean albumin values of both groups from day 3 post-commencement of treatment when compared to their initial values before the commencement of treatment (Table 4). The albumin values from day 3 post-commencement of treatment were below the reference range.

#### 3.5.5 Urea

There was an increase in the mean urea value (above the reference value) of both groups (oxytetracycline-treated and penstrep-treated groups) before the commencement of treatment. By day 3 post-commencement of treatment, there was a significant decrease ( $P < .05$ ) in the mean urea values of the penstrep-treated group when compared to their initial values on day 0 (Table 4).

#### 3.5.6 Creatinine

There were no significant differences ( $P > .05$ ) in the mean creatinine values of both groups all through the duration of treatment (Table 4). The creatinine values all through the experimental/ treatment and monitoring period were within the reference range.

#### 3.5.7 Potassium (K)

There was an increase in the mean potassium value (above the reference range) of penstrep-treated group before the commencement of treatment. Following treatment by day 3, it decreased to nearly normal value, and by day 7 post-commencement of treatment the potassium value increased above the reference value although not significant ( $P > .05$ ). There were no significant differences ( $P > .05$ ) in the mean potassium levels of oxytet-treated group all through the duration of treatment (Table 4).

#### 3.5.8 Sodium (Na)

There was an increase in the mean sodium level (above the reference range) of oxytet-treated group before the commencement of treatment. Following treatment by day 3, it continued to increase, and by day 7 post-commencement of treatment the sodium level decreased and was within the reference value although not significant. There were no

significant differences ( $P > .05$ ) in the mean sodium levels of penstrep-treated group all through the duration of treatment (Table 4).

**Table 4. Biochemistry of goats naturally infected with PPR and treated with oxytetracycline and penstrep**

Group/parameter	Day 0	Day 3 post-treatment	Day 7 post-treatment
<b>AST (IU/L)</b>			
A- Oxytet treated	201.34 ± 21.29 <sup>a</sup>	229.58 ± 25.84 <sup>a</sup>	243.46 ± 10.59 <sup>a</sup>
B- Penstrep treated	287.44 ± 15.05 <sup>a</sup>	196.05 ± 16.12 <sup>a</sup>	224.99 ± 12.24 <sup>a</sup>
Reference value	0-300 IU/L [16]		
<b>ALT (IU/L)</b>			
A- Oxytet treated	9.13 ± 3.23 <sup>a</sup>	13.33 ± 2.93 <sup>a</sup>	18.90 ± 2.63 <sup>a</sup>
B- Penstrep treated	12.70 ± 4.88 <sup>a</sup>	9.61 ± 0.10 <sup>a</sup>	14.77 ± 1.32 <sup>a</sup>
Reference value	6-19 IU/L [15]		
<b>Total protein (mg/gl)</b>			
A- Oxytet treated	6.28 ± 1.21 <sup>a</sup>	7.65 ± 0.69 <sup>a</sup>	6.36 ± 0.51 <sup>a</sup>
B- Penstrep treated	6.84 ± 1.83 <sup>a</sup>	6.66 ± 0.74 <sup>a</sup>	7.99 ± 0.34 <sup>a</sup>
Reference value	6.3-8.5 g/dl [14]; 6.2-7.9 g/dl [15]		
<b>Albumin (mg/dl)</b>			
A- Oxytet treated	2.87 ± 0.32 <sup>a</sup>	2.16 ± 0.33 <sup>b</sup>	1.84 ± 0.22 <sup>b</sup>
B- Penstrep treated	3.34 ± 0.50 <sup>a</sup>	1.89 ± 0.35 <sup>b</sup>	2.47 ± 0.30 <sup>b</sup>
Reference value	2.9-4.3 g/dl [14]; 2.8-4.3 g/dl [15]		
<b>Urea (mg/dl)</b>			
A- Oxytet treated	36.28 ± 2.55 <sup>a</sup>	34.78 ± 4.20 <sup>a</sup>	34.42 ± 8.06 <sup>a</sup>
B- Penstrep treated	37.06 ± 0.78 <sup>a</sup>	27.29 ± 1.71 <sup>a</sup>	23.34 ± 8.00 <sup>a</sup>
Reference value	10-26 mg/dl [15]		
<b>Creatinine (mg/dl)</b>			

A- Oxytet treated	1.29 ± 0.16 <sup>a</sup>	1.65 ± 0.28 <sup>a</sup>	1.53 ± 0.03 <sup>a</sup>
B- Penstrep treated	1.23 ± 0.06 <sup>a</sup>	1.21 ± 0.14 <sup>a</sup>	0.90 ± 0.38 <sup>a</sup>
Reference value	0.6-1.6 mg/dl [15]		
<b>Potassium (mmol/L)</b>			
A- Oxytet treated	4.56 ± 1.19 <sup>a</sup>	4.34 ± 1.34 <sup>a</sup>	6.27 ± 1.75 <sup>a</sup>
B- Penstrep treated	7.02 ± 2.50 <sup>a</sup>	6.07 ± 1.47 <sup>a</sup>	7.81 ± 2.81 <sup>a</sup>
Reference value	3.0-6.0 mmol/L [14]; 3.4-6.1 mmol/L [15]		
<b>Sodium (mEq/L)</b>			
A- Oxytet treated	147.7 ± 14.74 <sup>a</sup>	154 ± 6.34 <sup>a</sup>	134.4 ± 15.00 <sup>a</sup>
B- Penstrep treated	139.5 ± 10.00 <sup>a</sup>	151 ± 5.84 <sup>a</sup>	138.305 ± 8.03 <sup>a</sup>
Reference value	124-146 mEq/L [14]		

Superscript a,b,c indicates significance row-wise

#### 4. DISCUSSION

The clinical manifestations of *Peste des petits ruminants* (PPR) observed in goats used in this study agree with the findings in previous studies [6,7,17]. The goats showing clinical signs suggestive of PPR but were negative for PPRV as seen in this study were suggested to be suffering from other respiratory and digestive diseases such as contagious caprine pleuropneumonia, pneumonic pasteurellosis, salmonellosis and colibacillosis, as this diseases share same clinical symptoms and thus were considered as differential diagnosis of PPR [18].

Following therapeutic intervention, only 20% of Oxytet-treated group and 35.7% of Penstrep-treated group had their clinical signs completely cleared indicating that the antibiotics used (oxytetracycline and penstrep) were not efficient in handling the possible secondary bacterial complications of PPR in the goats treated. This observation is in contrast with the report of Islam *et al.* [19] and Islam *et al.* [20] who reported 55% and 64% recovery rate respectively. This could be attributed to the symptomatic treatment carried along with oxytetracycline administration. Wosu, [5] had earlier reported a recovery rate of 58.8% using chloramphenicol, penicillin and streptomycin in addition to anthelmintic, fluid therapy, intestinal sedative and scrubbing of the labial scab with lemon fruit. However, Anene *et al.* [21] recorded a recovery rate of 14.29% in PPR infected goats using oxytetracycline, chloramphenicol 25% solution and metamerazine in different groups at recommended doses. Both antibiotics used in this study were indicated for the treatment of respiratory infections. Thus, the low recovery rate obtained may be attributed to treatment without supportive therapy. Supportive therapy is of great importance in the successful management of PPR.

Mortality rate of 41.67% recorded in this study could also be attributed to treatment without supportive therapy and presence of an underlying disease conditions such as helminthosis complicating the situation. The work of Sharma and Tarunpreet-Joshi [22] recorded a lower mortality of 18.38% using antibiotic (ceftriazone), anti-inflammatory, antihistamine, levamisole, vitamin C, B-complex and vitamin A, D and E injections in the therapeutic management of PPR infected goats.

Pyrexia observed before the commencement of treatment is a feature of PPR, and it subsided following treatment. This finding is consistent with that of Islam *et al.* [19] who found that increased temperature of goats/sheep infected with PPR virus was reduced after antibiotic therapy.

The increased red blood cell (RBC) count observed before the commencement of treatment is attributed to the diarrhoeic nature of the disease and this agrees with the findings of Islam *et al.* [23] and Kataria *et al.* [24]. Diarrhoea is a symptom of PPR which leads to dehydration in the host resulting in haemoconcentration in the blood profile of PPR infected animals [24]. The non-significant difference observed in the total WBC despite treatment administered could be attributed to the activity of the virus. PPR virus is known to have lymphotropic activity as it has an affinity for lymphoid organs. Antibiotic treatment has no effect on the virus.

The Aspartate transaminase (AST) and Alanine aminotransferase (ALT) activities and total protein, albumin, creatinine, potassium and sodium levels which were within reference ranges during the viral infection may suggest that less or no damage has been done to the liver and kidney, or it could be that the reference values might not be true for our indigenous breed of goats. It has been reported that breed, physiological status, nutrition, season [25,26] and geographical location [27] affect biochemical profile. PPR infection has been reported to be associated with liver and kidney damage and was further confirmed by a marked increase in the level of ALT and AST, and marked decrease in total protein and albumin when compared with normal healthy small ruminants [14]. The penstrep-treated group had a significant increase in their albumin level by day 7 post commencement of treatment thus showing better recovery trait.

The increased urea level of both groups agrees with the findings of Balogun *et al.* [28] and Sharma and Tarunpreet-Joshi [22]. Balogun *et al.* [28] observed a high level of urea in the serum of WAD goats post PPR infection which continued to increase as the infection progressed from the acute stage to death. Following treatment, penstrep-treated group had their urea level returned to within the reference value thus showing a better recovery trait.

## 5. CONCLUSION

Based on our finding, the use of penstrep in the management of PPR cases gave a better result. Supportive therapy is of great importance in the improvement of survivability in PPR cases and thus should be included. Antibiotic sensitivity test is also advised for the most appropriate antibiotic of choice to be used in the management of PPR

## ETHICAL APPROVAL

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee.

## REFERENCES

1. Chukwudi IC, Ogbu KI, Nwabueze AL, Olaolu OS, Ugochukwu EI, Chah KF. Update on *Peste des petits ruminants* status in South East Nigeria: Serological and farmers awareness investigation, and potential risk factors. *Trop Anim Health Prod.* 2020a;52(6) 328 –3291. doi: 10.1007/s11250-020-02359-7.
2. Kumar KS, Babu A, Sundarapandian G, Roy P, Thangavelu A, Arumugam R, et al. Molecular characterisation of lineage IV *Peste des Petits Ruminants* virus using multigene sequencing data. *Vet Microbiol.* 2014;174(1-2):39-49. doi:10.1016/j.vetmic.2014.08.031.
3. Fadiga M, Jost C, Ihedioha J. Financial costs of disease burden, morbidity and mortality from priority livestock diseases in Nigeria. Disease burden and cost–benefit analysis of targeted interventions. ILRI Research Report 33. Nairobi, Kenya: ILRI; 2013.
4. Adams MJ, Lefkowitz EJ, King AMQ, Harrach B, Harrison RL, Knowles NJ, et al. Changes to taxonomy and the International code of virus classification and nomenclature ratified by the International Committee on Taxonomy of Virus. *Arch\_Virol.* 2017;162(8):2505-2538. doi: 10.1007/s00705-017-3358-5.
5. Wosu LO. Management of clinical cases of *Peste des Petits Ruminants* (PPR) disease in goats. *Beitr Trop Landwirtsch Veterinarmed.* 1989;27(3):357-361. PMID: 2619704.
6. Balamurugan V, Hemadri D, Gajendragad MR, Singh RK, Rahman H. Diagnosis and control of *Peste des Petits Ruminants*: a comprehensive review. *Virus Disease.* 2014;25(1): 39–56. doi: 10.1007/s13337-013-0188-2
7. Pope RA, Parida S, Bailey D, Brownlie J, Barrett T, Banyard AC. Early events following experimental infection with *Peste des Petits Ruminants* virus suggest immune cell targeting. *PLoS One.* 2013;8(2):e55830. <https://doi.org/10.1371/journal.pone.0055830>
8. OIE (Office International des Epizooties/ World Organization of animal health). *Peste des petits ruminants*. Technical disease card database; 2002.

9. Chukwudi IC, Ogbu KI, Luka PD, Malesa RP, Heath LE, Ugochukwu EI, et al. Comparison of colorimetric loop-mediated isothermal amplification kit and reverse transcription-PCR in the diagnosis of *Peste des Petits Ruminants* in sheep and goats in Southeast Nigeria. *Vet World*. 2020b;13(11): 2358-2363. doi:10.14202/vetworld.2020.2358-2363
10. Coles EH.. *Veterinary clinical pathology*, (4th edn), W.B Saunders Co., Philadelphia; 1986.
11. Schalm OW, Jain NC, Carrol EJ. *Veterinary haematology*, 3rd edn. Lea and Febiger, Philadelphia; 1975.
12. Duncan OD. Pathanalysis: sociological examples. *Am. J. Sociol*. 1966;72:1-16.
13. Robertshaw D. Temperature Regulation and Thermal Environment, In *Dukes' Physiology of Domestic Animals*, (12th ed), Reece WO, Ed. Copyright 2004 by Cornell University; 2004.
14. Daramola JO, Adeloye AA, Fatoba TA, Soladoye AO. Hematological and biochemical parameters of West African Dwarf goats. *Livest Res Rural Dev*. 2005;17(8):21-27. <http://www.lrrd.org/lrrd17/8/dara17095.htm>
15. Kaneko JJ, Harvey JW, Bruss ML. *Clinical Biochemistry of Domestic Animals*, 6th edition, 2008;551-553.
16. Jackson PGG, Cockcroft PD. *Clinical examination of farm animals*. Oxford, UK: Blackwell Science; 2002.
17. Das S, Nath R, Balamurugan V, Choudhury R, Devi M. Haematobiochemical analysis of goats naturally infected with *Peste des Petits Ruminants*. *Int J Res Emerg Sci Technol*. 2015;2(9):19-24. <https://ijrest.net/downloads/volume-2/issue-9/pid-ijrest-29201506.pdf>
18. Food and Agriculture Organisation (FAO). *Recognizing Peste des Petits Ruminants. A Field Manual*. Animal Production and Health Division. Viale della Teme di Caracalla: Rome Italy. 1999;Pp: 3-16, <http://www.fao.org/docrep/003/x1703e/x1703e00.HTM>.
19. Islam K, Ahad A, Mahmood A, Rahman MM, Islam MZ, Kabir MHB, et al. Prevalence and clinico-pathological features of *Peste des Petits Ruminants* in different breeds of Goats and their response to antimicrobial. *J Infect Mol Biol*. 2014;2(3):43-48. <https://doi.org/10.14737/jimb.2307-5465/2.3.43.48>
20. Islam MS, Khan MSI, Kader HA, Begum MR, Asgar MA. Prevalence of *Peste des Petits Ruminants* (PPR) of Goat and their response to antibiotic treatment at Mirzaganj Upazila of Patuakhali District. *J Environ Sci Nat Resour*. 2012;5(2):181-184. <https://doi.org/10.3329/jesnr.v5i2.14811>
21. Anene BM, Ugochukwu EI, Omamegbe JO. An appraisal of three treatment regimens for pneumoenteritis complex (PPR) in goats. *Bull Anim Health Prod Afr*. 1987;35:1-3.
22. Sharma SK, Tarunpreet-Joshi M. Clinico-haemato-biochemical characterization of *Peste des Petits Ruminants* in Sirohi goats and its management. *Int J Environ Sci Technol*. 2016;5:2200-2204

23. Islam M, Pathak DC, Das S, Rahman T, Sarma S, Borgohain R. Hematological and biochemical alterations in goats due to *Peste des Petits Ruminants* viral infection. *J Entomol Zool Stud.* 2018;6(3):710-713
24. Kataria AK, Kataria N, Gehlot AK.. Large scale outbreak of *Peste des Petits Ruminants* in sheep and goats in Thar Desert of India. *Slov Vet Res.* 2007;44(4):123–132. UDC 636.3.09:616.98-07:616-036.22:551.585.53
25. Arfuso F, Fazio F, Rizzo M, Marafioti S, Zanghi E, Piccione G. Factors affecting the haematological parameters in different goat breeds from Italy. *Ann Anim Sci.* 2016;16:743–757. doi: 10.1515/aoas-2015-0094.
26. Antunović Z, Novoselec J, Klir Ž. Hematological parameters in ewes during lactation in organic farming. *Poljoprivreda.* 2017;23:46–52. <https://doi.org/10.18047/poljo.23.2.7>
27. Antunović Z, Maric I, Klir Ž, Seric, V, Mioc B, Novoselec J. Haemato-biochemical profile and acid–base status of Croatian spotted goats of different ages. *Arch Anim Breed.* 2019;62:455–463. <https://doi.org/10.5194/aab-62-455-2019>
28. Balogun FA, Fasanmi OG, Oladipo TA, Popoola MA, Olona JF, Adeoye YD. Field evaluation and confirmation of acute *Peste des Petits Ruminants* outbreak in a flock of West African dwarf goats in Ibadan, Nigeria. *Int J Vet Sci Med.* 2017;5:175–180. doi: 10.1016/j.ijvsm.2017.08.004