

## Original Research Article

### **FUNGAL BIOFILMS IN DRINKING WATER STORAGE CONTAINERS: FLIPPING THOUGH THEIR HEALTH RISK POTENTIALS. A CASE-STUDY OF HOUSEHOLDS AT DIOBU, PORT HARCOURT, NIGERIA.**

#### **ABSTRACT**

The formation of fungal biofilm in drinking water storage containers has not been giving considerable attention over the years despite the challenges posed by bacterial biofilms. Hence, the study aimed to determine the fungal occurrence and the associated biofilm formation capacity from domestic drinking water storage containers in Diobu homes of Port Harcourt. Fifty water samples were obtained from domestic storage containers in different house units, and the sample was subjected to the standard microbiological procedure. The spread and streak techniques were applied on antibiotics treated Sabroaud Dextrose and Congo red growth media. The result shows that the mean fungi count of  $6.2 \times 10^2$  and  $6.1 \times 10^2$  CFU/ml in household sizes of 4 and 5 persons respectively were less than that of the household population of 2, 6, and 8 persons with the fungal count of  $8.7 \times 10^2$ ,  $8.3 \times 10^2$  and  $8.0 \times 10^2$  CFU/ml respectively. Positive, negative and zero relationships were established between the household sizes and the fungal recovered. The analysis further revealed a total of 25 fungal isolates of which three genera were distinguished morphologically, namely, *Aspergillus flavus*, *Candida albicans* and *Penicillium notatum*. All isolates were able to form a heavy mass of biofilm; except *Candida albicans* which had a delayed formation after 24hours with less biofilm development potential, and thus signifies its inability to persist, despite having the highest count or load. The success of fungi biofilm is based on the constant growth rate of *Aspergillus flavus* and *Penicillium notatum* biofilms for which, the household size has reasonably be attributed. The study recommends that storage facilities in the homes be regularly cleaned and always disinfected to inhibit the successes of fungal biofilm.

**Keywords:** Fungal Biofilm, Drinking Water, Storage Containers, Household Size

#### **Introduction**

Fungal contaminated drinking water has become essential to researchers and health practitioners alike due to the severe implications of fungal infection on immune-suppressed patients (Tekere *et al.*,2019). Fungi infection has also been linked to allergic diseases, including worsening asthmatic symptoms and conditions, hypersensitivity, and skin irritation (Tekere *et al.*,2019). The role of fungal contaminated water in the transmission of diseases was not given the required attention (Babic *et al.*,2017). Fungi and other microbes such as bacteria, viruses and protozoa were reported as biological pollutants in drinking water (Hamid *et al.*, 2014). Several years earlier, Arvanitidou *et al.* (2000) analysed the presence of fungi in drinking water samples obtained from different sources such as spring water, shower water, tap water, groundwater,

storage tank as well as water consumed in hospitals and schools from which the following fungi genera: *Fusarium*, *Aspergillus*, *Cladosporium*, *Epicoccum* and *Trichoderma* were isolated and identified. According to Grabinska-Lonesiska (2007), water can be contaminated by fungi in distribution pipes, thus adversely affecting the water's odour and taste. Pipe-borne water, when consumed, as reported in studies carried out by Hamid *et al.* (2014), is known to be strongly allergenic or successful in immune-suppressed persons of cancer, asthma and AIDS patients. Like *Fusarium* spp and *Aspergillus* spp, Fungi isolates were reported by Anaissie *et al.* (2003) to occur in water storage containers of some households where the fungi build up biofilm, which encourages its persistence.

Perhaps, a deep look into the definition of biofilms will help elucidate the subject matter of discuss. According to the American Environmental Protection Agency (2002), “ biofilm is a complex mixture of microbes, organic and inorganic material accumulated amidst a microbially produced organic polymer matrix attached to the inner surface of the water distribution system or storage containers” (EPA 2002). This definition of biofilm, points to the fact they protect microbes from disinfection and allow microbes injured by environmental stress and disinfectants to recover and grow afterwards; causing deterioration of water quality, generation of bad tastes and odors, and proliferation of macroinvertebrates. Such contamination and the materials in the biofilm may subsequently be slowly released into the flowing water under various circumstances for prolonged constant contamination of the slowing water systems or storage water. This underpinned the importance of biofilms and their significance to public health.

Meanwhile, over three decades ago, fungal biofilm was reported significant in distribution pipes. The water-borne in the underground piping line is contaminated (LeChavallier & Babcock, 1987), affecting the water storage containers. The risk of fungal infection resulting from water consumption calls for the investigation of the fungal biofilm success in the containers of stored drinking water. Fungi are unevenly distributed in water because they are held in biofilms, capable of establishing and multiplying, and have a long residence time in the water. The significance of this study is that the occurrence of fungal biofilm in water storage containers accounts for some related fungal infections in humans. Moreso, it is reported that fungal biofilm occurrence increases the risk of fungal transmission in food handling facilities and the risk of human illnesses (Eduar *et al.*, 2011). Fungal biofilm has also been noted by Bravo *et al.* (2010) to have increased resistance to disinfectants, antibiotics and environmental stressors such as salt, pH, which thus makes it challenging to address problems that arise from their formation. Therefore, the study focuses on the determination of fungal density in stored drinking water with respect to household size of homes, which per se is a critical factor in the quality of drinking water stored in homes.

## **Materials and Methods**

### **Study Area**

The study area for this study is Diobu, Port Harcourt, Nigeria. Diobu was chosen due to its several squatter settlement homes with rooms that are averagely overcrowded. Broadly, the

houses have a structure that gives it an average calibrated indoor temperature of 37 degrees celsius, and such home condition encourages fungal growth with implications for fungal biofilm formation and associated health risk

### Sample Collection

Sources of water to these settlements are private borehole service providers. Based on the number of inhabitants or household sizes, 50 water samples were collected aseptically. Ten samples were collected from various housing units/settlements and grouped into five (5) household sizes(2,4,5,6 and 8).

### Fungal Bioload Evaluation

Total heterotrophic fungi biomass were determined on abroad dextrose medium as prepared and described by Wemedo *et al.* (2016) using the spread plate technique, where a 0.2ml volume potion of the water samples was inoculated into a freshly prepared medium using a1 ml pipette and a glass spreader to spread the sample over the Saboraud destrose medium evenly. It was then followed by incubation at 25°C for five days. Biomass /fungal density results were obtained and reported as fungal forming unit per mill.

### Pearson's Correlation Coefficient

The population size or household size of the homes were correlated against the density of mould recovered in the study as carried out by Mbah & Oselebe (2004)

### Morphological Identification

The taxonomical identity of the fungal was determined based on Ellis and Ellis, 1988. The identified isolates were then sub-cultured to get pure forms for further study.

### Biofilm Test

Biofilm analysis was achieved by preparing the biofilm agar and inoculating the test fungal organisms into it by streaking, as Amadi-Ikpa et al. (2020) described.

## Result

### Enumeration of Fungi Biomass in the Home

Table 1 shows the mean fungi count of  $6.2 \times 10^2$  and  $6.1 \times 10^2$  CFU/ml in the household sizes of 4 and 5 persons respectively were less than that of the household population of 2, 6, and 8 with the fungal count of  $8.7 \times 10^2$ ,  $8.3 \times 10^2$  and  $8.0 \times 10^2$  CFU/ml respectively. Although statistically, the mean count of fungi in household sizes of 4 and 5 persons was not significant to 2,6 and 8 persons at a  $P > 0.05$ . Household sizes of 2, 5, 6, and 8 persons had higher yeast counts than mould counts at a significant difference of  $P < 0.05$ .

**Table 1: Mean Enumeration of Fungi Biomass in the Home**

S/No	Household Size	Yeast (CFU/ml)	Mould (CFU/ml)	Total Mean Count (CFU/ml)
------	----------------	----------------	----------------	---------------------------

I	2	$4.4 \times 10^2$	$4.3 \times 10^2$	$8.7 \times 10^2$
II	4	$1.4 \times 10^2$	$4.8 \times 10^2$	$6.2 \times 10^2$
III	5	$5.9 \times 10^2$	$2.0 \times 10^1$	$6.1 \times 10^2$
IV	6	$8.1 \times 10^2$	$2.0 \times 10^1$	$8.3 \times 10^2$
V	8	$7.9 \times 10^2$	$1.0 \times 10^2$	$8.0 \times 10^2$

### Pearson's Correlation Coefficient Between Household Sizes and Fungi Isolates from the Water Sample

Figure 1A below shows a positive relationship between the household population size and the density of yeast recovered from the water sample. The correlation coefficient shows a positive  $R^2$  value of 0.4819. whereas, Figure 1B shows a negative relationship between the household population size and the density of mould recovered from the water sample. The correlation coefficient shows an  $R^2$  value of 0.6379. Figure 1C shows no relationship between the household population size and the density of fungi recovered from the water sample. The correlation coefficient shows an  $R^2$  value of zero (0).




Figure 1A

Figure 1B

Figure 1C

**Figure 1 Pearson's Correlation Coefficient Between Household Sizes and Fungal Isolates from the Water Samples.**

### Morphological Identification of the Fungal Isolates

In Table 2 below, A total of 25 fungal isolates were obtained in the study, from which two fungal classifications (mould and yeast) were derived, making up three genera, namely; *Aspergillus flavus*, *Candida albicans* and *Penicillium notatum* with the yellow, milky and blue cob web appearance on culture media respectively.

**Table 2 Morphological Identification and Occurrence of Fungal Population**

Morphological Features	Fungal Occurrence	Percentage Fungal Occurrence (%)	Fungal Isolates
Yellow pigmentation with white background surface circular in shape with an elevated centre	7	28	<i>Aspergillus flavus</i>
Milky colonies with mainly surface growth growing singly	15	60	<i>Candida</i> spp
Blue cobweb or hair-like appearance growing to cover plate	3	12	<i>Penicillium notatum</i>

### **Biofilm Forming Ability**

Plate 1: bellow shows a biofilm test of the fungal isolates obtained in the investigation. *Aspergillus flavus* and *Penicillium notatum* isolates were able to form a biofilm, while *Candida albicans* did not constitute a biofilm



*Penicillium notatum*

*Aspergillus flavus*

*Candida albicans*

### **Plate 1; Fungal Biofilm Formation**

#### **Discussion**

Biofilms are the predominant mode of microbial growth within the drinking water distribution systems, constituting a key public health challenge as consumption of contaminated water with pathogenic biofilms has been linked to human infections and waterborne outbreak in recent past (Angles et al., 2007; Prest et al., 2016; Douterelo et al., 2016). And the major biofilm producing bacteria in drinking water are *P. aeruginosa*, *Campylobacter jejuni*, *Legionella pneumophila*, *Mycobacteria*, *Aeromonas hydrophila*, and *Klebsiella pneumoniae* (Chan et al., 2019). However, from this study, household sizes of 2, 5, 6 and 8 persons had higher counts of yeast than mould counts obtained. Also, it suggests the impact of seasonal variation, specifically, the summer season, as reported by Urooj *et al.* (2018) and observed in the present circumstance. The mean counts of Fungi in the household sizes of 4 and 5 persons were less than that of the household population of 2, 6, and 8 persons. Statistically, the mean count of fungi in household units of 4 and 5 persons was not significant at a  $P > 0.05$ . Thus, a high microbial count in a densely polluted city, as Amin (2008) reported, indicates an unhygienic condition of a food processing area. Following this, Household sizes of 2, 5, 6 and 8 persons had higher yeast counts than mould counts. Thus, counts of yeast were far more significant than mould. A positive relationship exists between the household population size and the density of yeast recovered from the water sample, which signifies that the yeast count increases as the household population size increases. Perhaps, this may have resulted from favourable growth factors such as optimum temperature, nutrient and hydrogen ion concentration (pH) (Mbah & Oselebe, 2004). The negative relationship between the household population size and the density of mould recovered from the water sample signifies that the household population size increases. Hence, the mould counts decrease, thus collaborating with studies carried out by Mbah & Oselebe (2004), where they listed temperature, pH, nutrients and light as factors that could inhibit mould growth if not harnessed. However, the household population sizes and the density of fungi recovered from the

water sample showed no relationship exist; that is to say, the size of a household does not affect fungi growth (Hall & Noverr, 2017). A total of 43 fungal isolates were obtained in the study, thus disagrees with investigations carried out by Mhlongo *et al.* (2019), who got more than 43 isolates. Similarly, Goussou *et al.* (2009) recovered eight (8) indigenous fungal genera, in which this study obtained only three genera, namely, *Aspergillus* spp, *Candida* spp and *Penicillium* spp. The high prevalence of *Candida albicans* in this study suggests why it is the most common fungi isolated from water (Goussou *et al.*, 2009). *Aspergillus flavus* has been isolated from drinking water in many studies. Thus, this study agrees with Annaise *et al.* (2003), who submitted that water is increasingly recognised as an environmental source of *Aspergillus* spp with augmenting that suggests water is considered an important route of transmission pathogenic *Aspergillus* spp. The presence of *Penicillium notatum* in the drinking water sample signifies it as a source of exposure. However, the source of *Penicillium* infection is not yet known (Rodrignes & Nosanchuk, 2020). *Aspergillus flavus* and *Penicillium notatum* isolate formed a heavy biofilm mass, while *Candida albicans* developed a low biofilm density. This study is inconsonant with the one carried out by Siqueira (2013), who evaluated the capacity of some fungi isolates to form biofilms and reported in their findings that *Aspergillus flavus* and *Penicillium notatum* showed better ability in terms of higher biomass to form biofilm when compared to other fungi isolates, in which case *Candida albicans*. Similarly, the inability of *Candida albicans* to form a biofilm of heavy biomass has also been reported by Pathak *et al.* (2012).

Despite the fact that fungi have been detected in drinking water distribution systems and biofilms, they have yet to be definitively linked to waterborne disease. The fungal pathogens found in the distribution system, on the other hand, are opportunistic and seldomly cause any sickness, regardless of the source of infection. *A. flavus* and several other *Aspergillus* species found in drinking water, on the other hand, create powerful toxins (mycotoxin). Aflatoxins, cyclopiazonic acid, ochratoxin A, and sterigmatocystin are examples of mycotoxins produced by *Aspergillus* species. Citrinine, patulin, and penicillic acid, which are likewise produced by *Penicillium* species, and several *Aspergillus* species (CDC, 2020). Aflatoxin, a species-specific toxin generated by *Aspergillus flavus*, a specie implicated in this research, is the most important *Aspergillus* mycotoxin. According to Bennett and Kich (2003) and Gülhan (2020), acute aflatoxicosis outbreaks have been documented in a number of locations around the world, despite the fact that acute poisoning in humans is uncommon. Acute aflatoxicosis in humans causes vomiting, stomach pain, jaundice, pulmonary edema, unconsciousness, convulsions, and death. Aflatoxins are carcinogens that cause DNA damage. Aflatoxin exposure over time has been linked to liver disorders such as cancer, cirrhosis, hepatitis, and jaundice. The *Penicillium* mycotoxins that affect liver or kidney function, by acute or chronic exposure, are usually asymptomatic in humans or animals (Gülhan, 2020; Bennett and Kich, 2003). Consequently, the American Environmental Protection Agency (2002) recommended consistent monitoring of standards. However, in the Nigerian context, standards monitoring of drinking water supplies in households or neighbourhoods is a herculian task, if not non-existent.

*Candida albicans* (candidiasis), a harmful yeast, implicated in this research, can also colonize the GI tract. (Schaechter et al., 1998) and thus could potentially be spread by the fecal-oral route. It has been found in seawater and in sand at marine beaches. *C. albicans* and other important fungal pathogens are also associated with soil. Thus, the potential for waterborne disease caused by fungi in the biofilm exists, but the significance is unknown as was the opinion of Azuonwu et al., (2019) which underpinned that water storage containers and underground water distribution systems could potentially be a source of harmful pathogens to humans. Thus, suggesting the need for consistent monitoring, surveillance and proactive water quality evaluation; in their words, "keeping a careful watch at all times from the public health point of view, over safety and acceptability of drinking water supply" standards (Azuonwu et al., 2019). Perhaps with proper monitoring and evaluation of drinking water quality, prevention of the effects of biofilm forming microbes that could lead to life-threatening infections and diseases in humans such as cystic fibrosis (CF), otitis media, periodontitis, infective endocarditis (IE), chronic wounds, and osteomyelitis would be achieved (Southey-Pillig et al., 2005; Akyildiz et al., 2013; Masters et al., 2019). Similarly, U.S. Food and Drug Administration (FDA 2020) and the European Food Safety Authority standards guidelines could be used as the spring board for strategic monitoring of these standards.

### **Conclusion**

The Isolates, *Aspergillus flavus* and *Penicillium notatum* created a high mass of biofilm for their survival. *Candida albicans* made a low mass of biofilm, despite *Candida albicans* high recovery density or biomass count. Thus, the success of fungal in this study is based on the persistence rate of biofilm formed by *Aspergillus flavus* and *Penicillium notatum* which are notable toxin producers and potential public health risks. The household size has somewhat been implicated. The study recommends that storage facilities in the homes be cleaned and disinfected always, and regular turn-over of water in storage cans should be adhered to religiously. Therefore, Standards monitoring, evaluation and enforcement, should be strengthened more especially in low income settings like this. Nonetheless, it is firmly advocated that pipe-borne water in storage containers from these affected areas, irrespective of household size, should be treated, boiled, and filtered before use for drinking and washing fresh fruits to avert the possibility of an outbreak of water-borne epidemic in the area.

### **Reference**

- Aaron U. U. & Onoja H. (2017). Incidence of Dermatophyte Infections among Primary School Pupils In Three Selected Primary Schools In Obio-Akpo Local Government Area Of Rivers State, Nigeria. *Current Studies in Comparative Education, Science and Technology*, 4(1), 40-49.
- Amadi-Ikpa, C. N., Akani, N. P. & Wemedo, S.A. (2020). Biofilm Formation and Virulent Properties of Bacterial Isolates in Stored Drinking Water of Some Homes.

*International Journal of Research and Innovation in Applied Science* 5 (8), 5454 – 6194.

- Amin, M.A. (2018). Microbiological Quality Analysis of Commercial Fruit Juice in Dhaka City. *Bangladesh Journal Online*.
- Arvanitidou, M., Spain, S., Velegraki, A., Pazarloglou, M., Kenetidis, D. & Pangidis, P. (2000). High Level of Recovery of Fungi from Water and Dialysate in Haemodialysis Unit. *Journal of Hospital Infection* 129: 74-77.
- Anaisie, E.J., Stratton, S. L., Dignani, M.C., Lee, C. K., Summerbell, R.C. (2003). Pathogenic Mold in Hospital Water Distribution System: a 3-Year Prospective Study & Clinical Implications for Patients with Hematologic Malignancies. *Blood*, 101, 2542-2546.
- Azuonwu, O., Aaron, U, U. and Ekong, I. U. (2019). Investigation of Prevalence of Escherichia coli in Public Drinking Water sources randomly Collected in and around Doibu Residential Area of Port Harcourt, Niger Delta. *International Journal of Research Studies in Microbiology and Biotechnology (IJRSMB)*, 5(4), 6-11.
- Babic, M. N., Gunde-Cimerman, N. & Brandao J. (2017). Fungal Contaminants in Drinking Water Regulation/ A Tale of Ecology Exposure, Purification and Clinical Relevance, *International Journal of Environmental Research and Public Health*, 4 (8), 362-371.
- Bennett, J. W. and Klich, M. (2003). "Mycotoxins." *Clinical Microbiology Review*, 16(3), 497–516.
- Bravo, E.A., Zagarra, A.J & Piscocoya, A. (2010). Chronic Diarrhea and Pancolitis Caused by Paracoccidioidomycosis: a Case Report; *Journal of Medical Case Reports*, Article ID 140505.
- CDC. 2020. "Toxins." Centers for Disease Control and Prevention. <https://www.cdc.gov/biomonitoring/toxins.html>.
- Eduar, A.B., Arturo, I.Z., Alejandro, P., Jose, I.P., Raul, T.(2011). Dimorphic Fungal Coinfection as a Cause of Chronic Diarrhea and Pancolitis. *Case Report in Medicine*, 2011, 960638.
- Ellis, M.B. & Ellis, J.P. (1988). *Microfungi on Miscellaneous Substrates. An Identification Handbook*. Croom Helm London, Timber Press Portland, Oregon.
- EFSA. 2020. "Mycotoxins." European Food Safety Authority. <https://www.efsa.europa.eu/en/topics/topic/mycotoxins>.

- FAO and WHO. 2018. "Codex Alimentarius: Understanding Codex." 5th ed. Food and Agriculture Organization of the United Nations and World Health Organization. Rome.
- FDA. 2020. "Natural Toxins and Mycotoxins." U.S. Food and Drug Administration. <https://www.fda.gov/food/chemicals-metals-pesticides-food/natural-toxins-and-mycotoxins>.
- Gülhan (2020). Food Safety and Quality. Mycotoxins and Their Health Impacts. *International Food Technology Magazine*, 78(8) online. <https://www.ift.org/news-and-publications/food-technology-magazine/issues/2020/august/columns/food-safety-and-quality-mycotoxins-and-their-health-impacts>.
- Goussou, S.J., Hameed, K.M. & Saadoun, I.(2009). Isolation and Evaluation of Indigenous Fungal and Bacterial Isolates as Potential Bio agents Against. *Broom Rape in Jordan* 8(3), 98-105.
- Grabinska-Loniewska, A., Konillowicz, T.K., Wardzynska,G. & Boryn, K. (2007). Occurrence of Fungi in Water Distribution System. *Polish Journal of Environmental Studies* 16 (4), 539-547.
- Hall, R. A. & Noverr, M. C. (2017). Fungal Interacts with the Human Host Exploring the Spectrum of Symbiosis, *Current Opinion in Microbiology* 40, 58-64
- Hamid, M.A., Tianling, X. & Xin, Y. (2014). Fungi Contamination of Drinking Water of Environmental Contamination and Toxicology. 228; 121
- LeChavallier, M.W. & Babcock, T.M. (1987). Examination and Characterisation of Distribution System Biofilm. *Applied Environmental Microbiology*, 53, 2714- 2724.
- Mbah, M. C. & Oselebe H. (2004). *General Biology* (Third Edition). Abakiliki. Ebonyi State: Speed and Skill Publication Company.
- Mhlongo, N.T., Tekere, M. and Sibanda, T. (2019). Prevalence and Public Health Implications of Mycotoxigenic Fungi in Treated Drinking Water System. *Journal of Water and Health*, 17(4) 517-531
- Pathak, A.K., Sharma, S. & Shrivastva, P. (2012) Multi-Species Biofilm of *Candida albicans* and Non-Candida spp on Acrylic Substrate, *Journal of Applied Oral Science*.1678-7757.

- Rodrigues, M. L. & Nosanchuk, J.D. (2020). Fungal as a Neglected Pathogen: A Wake-Up Call to Public Health Officials *PLoS Neglected Tropical Disease* 14 (2), 71-64.
- Siqueira, V. M. (2013). Biofilm Formation by Filamentous Fungi Recovered from a Water System, *Journal of Mycology*. 10; 11-55
- Tekere, M., Sibanda, T. & Mhlongo, Ntombie T. M. (2019). Prevalence and Public Health Implication of Mycotoxigenic Fungi in Treated Drinking Water System, *Journal of Water System* 17, (4): 517-531.
- Urooj, S., Mirani, Z. A. & Naz, S. (2018). Impact of Seasonal Variation on Bacterial, Yeast, Mold Counts in Drinking Water Collected from Karachi. Pakistan, *PSM Microbiology* 37-42.
- Wemedo, S.A, Amadi-Ikpa, C. N. & Essien , J.P. (2016). Population and Virulent Attributes of bacteria in Sachet Water Sold in a Port Harcourt Subhub (Rumuepirikom) *Journal of Biology and Genetic Research*,2 (3), 2545-5710.
- Akyıldız, I., Take, G. U., Uygur, K., Kızıl, Y. and Aydil, U. (2013). Bacterial Biofilm Formation in the Middle-Ear Mucosa of Chronic Otitis Media Patients. *Indian Journal of Otolaryngology . Head Neck Surgery*, 65(3), S557–S561; DOI 10.1007/s12070-012-0513