

## Case study

### Scrub Typhus presenting as severe thrombocytopenia

#### Abstract:

Scrub typhus is a zoonotic disease caused by *Orientia tsutsugamushi*. A rare finding of Scrub typhus with thrombocytopenia has been reported in very few cases. Thrombocytopenia is usually complicated with multiorgan dysfunction and high mortality rate. Here we report a case of a diagnosed scrub typhus infection with severe thrombocytopenia and management of such a case.

Keywords: Scrub typhus, thrombocytopenia

#### Introduction:

Scrub typhus is a zoonotic illness caused by *Orientia tsutsugamushi* and spread by mites<sup>1</sup>. It is a threat to public health and mostly affects Asia-Pacific regions; every year it burdens the economy and health by affecting one billion people and causes illness around the world<sup>2</sup>. Scrub typhus is characterised by a sudden high fever, headache which is severe in intensity, lymphadenopathy, generalised myalgia, rash and eschar<sup>3</sup>. Due to the lack of characteristic symptoms, accurate and effective methods for diagnosing scrub typhus are lacking, which frequently leads to a difficulty in diagnosis. Disseminated vasculitis is a pathologic feature of scrub typhus, and it can affect one or more organs, including the skin, liver, lung, kidney, brain and meninges<sup>4</sup>. Scrub typhus, on the other hand, is most commonly manifested by a scab at the site of the mite's bite. Approximately one third of the patients suffer from multiorgan dysfunction during the course of the illness<sup>5</sup>. One of the differential diagnoses for fever with thrombocytopenia or hemorrhagic fever is scrub typhus<sup>6</sup>. Scrub typhus mortality ranges from 0% to 30% in untreated individuals, and it varies according to age and infection location<sup>7</sup>.

Scrub typhus is a rare disease which is difficult to diagnose and hence makes treatment challenging. In this report, we have discussed a case of scrub typhus manifested as thrombocytopenia posing a challenge in diagnosis and treatment. Also the course of the disease in such case is discussed.

#### HISTORY

A 19 year old female presented to us with complaints of fever, vomiting and abdominal pain since 5 days. She took some treatment outside but treatment particulars were not available. There was no history of travel to another place.

On examination, her blood pressure was 100/60 mm hg, pulse rate was 80/min, axillary temperature of 101 F was recorded, there was no skin rash, per abdominal examination was suggestive of mild

splenomegaly. Routine blood investigations revealed haemoglobin 14 gm%, TLC 3100 per cu mm, platelet count 16000 per cu mm, MCV 83.8, SGPT-218, SGOT-49, Alkaline Phosphatase-254, Total bilirubin 3.4, conjugated 2.3, unconjugated 1.1, aPTt- control 30.0, aPTt- patient 30.7, prothrombin time-control 12.5, prothrombin time-patient 13.4, malaria strip test was negative. Serology for Scrub Typhus- Immunofluorescence assay was positive (light bands were seen). Other investigations were all within normal range.

On the basis of history and laboratory findings, diagnosis was made as Scrub typhus presenting as severe thrombocytopenia.

Initial treatment was started in the form of antibiotics, Inj Ceftriaxone 1 gm IV 12 hourly, Inj Metronidazole 500 mg IV 8 hourly, Cap Doxycycline 100 mg 12 hourly. 2 units of Single donor platelets were transfused. Supportive treatment was given in the form of IV fluids, multivitamin supplements and antacids.

After 3 days of admission, platelet counts started showing an increasing trend with a count of 80000 per cu mm, liver enzymes showed a declining trend.

On day 5, platelet count was 1,36,000 per cu mm and liver function improved furthermore.

Patient was discharged on day 5 in a hemodynamically stable condition and advised to follow up on a later date with fresh platelet count and liver function test reports. On follow up after 7 days, platelet counts and liver enzymes were within normal range.

#### Discussion:

Scrub typhus diagnosis is difficult and when multiple organs are damaged, the rate of diagnosis error occurs. One of the most common causes of scrub typhus misdiagnosis is doctors' lack of understanding of the disease's clinical symptoms<sup>8,9</sup>. Till 10 days after the onset of fever, the detection of nucleic acid like HtrA (periplasmic serine protease), type-specific antigen rrs (16S rRNA), and groEL (the heat shock protein Hsp60) is accurate. After that, serology plays a more specific role<sup>10</sup>. Furthermore, recent investigations have indicated that Raman scattering-based lateral flow test approaches are both quick and accurate in the diagnosis of the disease<sup>11</sup>. Scrub typhus biomarkers include *O. tsutsugamushi* immunoglobulin G (IgG) in patient blood, and an immunofluorescence assay is the best and most widely accepted approach for diagnosis (IFA). Improved anti-Orientia immunoglobulin M (IgM) and IgG-based rapid diagnostic tests and ELISAs have recently superseded subjective IFA due to their excellent sensitivity, specificity, and repeatability<sup>11,12,13,14</sup>. Scrub typhus with rapid advancement should be given special attention since it might cause hypothermia, tachycardia, crepitation, a low lymphocyte count, hypoalbuminemia, high aspartate aminotransferase, raised serum creatinine, and positive urine albumin<sup>14</sup>.

Thrombocytopenia and MODS are possible side effects of scrub typhus. Previous research has shown that thrombotic microangiopathic syndrome (TMAS) is the cause of thrombocytopenia-related MODS<sup>15,16</sup>. Thrombotic thrombocytopenic purpura (TTP), secondary thrombotic microangiopathy (TMA), and DIC are some of the pathological alterations that can arise<sup>15</sup>. To our knowledge, however, only a single case of scrub typhus presenting thrombocytopenia and MODS has been

documented thus far<sup>17</sup>. Medications should be used as soon as possible to avoid complicated scrub typhus, and practitioners usually have access to a variety of effective antibiotics<sup>18</sup>. Doxycycline, chloramphenicol, tetracycline, and azithromycin are all effective drugs against scrub typhus. Furthermore, with the same effect, azithromycin may be more tolerated than doxycycline<sup>19</sup>.

#### Conclusion:

We conclude that Scrub typhus being a rare disease might be complicated in the form of thrombocytopenia and needs more research into this aspect. It is associated with higher morbidity and mortality thus necessitating the need for aggressive treatment for better outcomes.

#### References:

- 1) Lee BJ, Chen CY, Hu SY, Tsan YT, Lin TC, Wang LM. Otaglia and eschar in the external auditory canal in scrub typhus complicated by acute respiratory distress syndrome and multiple organ failure. *BMC infectious diseases*. 2011 Dec;11(1):1-4.
- 2) Kelly DJ, Fuerst PA, Ching WM, Richards AL. Scrub typhus: the geographic distribution of phenotypic and genotypic variants of *Orientia tsutsugamushi*. *Clinical Infectious Diseases*. 2009 Mar 15;48(Supplement\_3):S203-30.
- 3) Rajapakse S, Weeratunga P, Sivayoganathan S, Fernando SD. Clinical manifestations of scrub typhus. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2017 Feb 1;111(2):43-54.
- 4) Dogra S. Recent advances in understanding pathophysiology of scrub typhus. *JK science*. 2010 Apr 1;12(2):70.
- 5) Peter JV, Sudarsan TI, Prakash JA, Varghese GM. Severe scrub typhus infection: clinical features, diagnostic challenges and management. *World journal of critical care medicine*. 2015 Aug 4;4(3):244.
- 6) Kothari VM, Karnad DR, Bichile LS. Tropical infections in the ICU. *The Journal of the Association of Physicians of India*. 2006 Apr 1;54:291-8.
- 7) Watt G, Parola P. Scrub typhus and tropical rickettsioses. *Current opinion in infectious diseases*. 2003 Oct 1;16(5):429-36.
- 8) Jamil MD, Hussain M, Lyngdoh M, Sharma S, Barman B, Bhattacharya PK. Scrub typhus meningoencephalitis, a diagnostic challenge for clinicians: a hospital based study from North-East India. *J Neurosci Rural Pract*. 2015;6(4):488–493. doi: 10.4103/0976-3147.169769
- 9) Sharma SR, Masaraf H, Lynrah KG, Lyngdoh M. Tsutsugamushi disease (scrub typhus) meningoencephalitis in north eastern India: a prospective study. *Ann Med Health Sci Res*. 2015;5(3):163–167. doi: 10.4103/2141-9248.157486
- 10) Paris DH, Dumler JS. State of the art of diagnosis of rickettsial diseases: the use of blood specimens for diagnosis of scrub typhus, spotted fever group rickettsiosis, and murine typhus. *Curr Opin Infect Dis*. 2016;29(5):433–439. doi: 10.1097/QCO.0000000000000298.
- 11) Blacksell SD, Tanganuchitcharnchai A, Nawtaisong P, Kantipong P, Laongnualpanich A, Day NP, Paris DH. Diagnostic accuracy of the InBios scrub typhus detect enzyme-linked

- immunoassay for the detection of IgM antibodies in Northern Thailand. *Clin Vaccine Immunol.* 2016;**23**(2):148–154. doi: 10.1128/CVI.00553-15
- 12) Rodkvamtook W, Zhang Z, Chao CC, Huber E, Bodhidatta D, Gaywee J, Grieco J, Sirisopana N, Kityapan M, Lewis M, et al. Dot-ELISA rapid test using recombinant 56-kDa protein antigens for serodiagnosis of scrub typhus. *Am J Trop Med Hyg.* 2015;**92**(5):967–971. doi: 10.4269/ajtmh.14-0627.
  - 13) Silpasakorn S, Srisamut N, Ekpo P, Zhang Z, Chao CC, Ching WM, Suputtamongkol Y. Development of new, broadly reactive, rapid IgG and IgM lateral flow assays for diagnosis of scrub typhus. *Am J Trop Med Hyg.* 2012;**87**(1):148–152. doi: 10.4269/ajtmh.2012.11-0583.
  - 14) Kingston HW, Blacksell SD, Tanganuchitcharnchai A, Laongnualpanich A, Basnyat B, Day NP, Paris DH. Comparative accuracy of the InBios scrub typhus detect IgM rapid test for the detection of IgM antibodies by using conventional serology. *Clin Vaccine Immunol.* 2015;**22**(10):1130–1132. doi: 10.1128/CVI.00390-15.
  - 15) Nguyen TC, Carcillo JA. Bench-to-bedside review: thrombocytopenia-associated multiple organ failure--a newly appreciated syndrome in the critically ill. *Crit Care.* 2006;**10**(6):235. doi: 10.1186/cc5064.
  - 16) Nguyen TC, Cruz MA, Carcillo JA. Thrombocytopenia-associated multiple organ failure and acute kidney injury. *Crit Care Clin.* 2015;**31**(4):661–674. doi: 10.1016/j.ccc.2015.06.004.
  - 17) Ittyachen AM, Abraham SP, Krishnamoorthy S, Vijayan A, Kokkat J. Immune thrombocytopenia with multi-organ dysfunction syndrome as a rare presentation of scrub typhus: a case report. *BMC Res Notes.* 2017;**10**(1):496. doi: 10.1186/s13104-017-2826-z.
  - 18) Wang CC, Liu SF, Liu JW, Chung YH, Su MC, Lin MC. Acute respiratory distress syndrome in scrub typhus. *Am J Trop Med Hyg.* 2007;**76**(6):1148–1152. doi: 10.4269/ajtmh.2007.76.1148.
  - 19) Wee I, Lo A, Rodrigo C. Drug treatment of scrub typhus: a systematic review and meta-analysis of controlled clinical trials. *Trans R Soc Trop Med Hyg.* 2017;**111**(8):336–344. doi: 10.1093/trstmh/trx066.