

## **Original Research Article**

# **PREVALENCE AND OUTCOME OF TOTAL ABDOMINAL HYSTERECTOMY PERFORMED FOR CERVICAL INTRAEPITHELIAL NEOPLASIA IN BENIN CITY**

---

### **ABSTRACT**

**Background:** Cervical cancer is the commonest malignancy of the female genital tract in developing countries. A preventable disease with a precursor; cervical intraepithelial neoplasia (CIN), can be detected via screening methods and can be managed by employing ablative or excisional method. However, the incidence and death have continued to rise in developing countries due to a lack of screening programs/follow up.

**Aims:** The study looked at the recourse to total abdominal hysterectomy (TAH) with CIN as the primary indication at the University of Benin Teaching Hospital as well as the outcome of surgery and observed complications.

**Methodology:** This was a 4-year retrospective cross-sectional analytical study that was conducted on 145 women who underwent total abdominal hysterectomy (TAH) for benign gynaecological conditions. Relevant data were extracted from case notes and hospital records. Analysis was performed using SPSS version 22.

**Results:** A total of 145 cases were analysed and the prevalence of TAH for CIN (n = 25) was 17.2%. Uterine fibroids contributed the highest prevalence for TAH at 51% followed by endometrial hyperplasia at 22.8%. The mean age distribution was 47.2 years and 96.6% were parous. The commonest adverse surgical outcome was haemorrhage, followed by a prolonged hospital stay.

**Conclusion:** Hysterectomy remains the most effective treatment for CIN. With the advent of effective ablative and excision procedures, it is considered an overtreatment in developed countries. However, the dearth of skills and equipment for these procedures and loss to follow up in developing countries make hysterectomy a viable treatment modality for CIN.

**Keywords:** Cervical intraepithelial Neoplasm, Total Abdominal Hysterectomy, Benign Gynaecological Conditions, ablative and excision procedures.

## 1. Introduction:

Cancer of the cervix is the second most common cause of cancer-related death in women worldwide [1] and the principal cause of cancer death in sub-Saharan Africa [2] where it places a significant social and economic burden on the health facilities [3]. An estimated 85% of global death from cervical cervix occurs in low- to middle-income countries compared to developed countries [2].

The reasons ascribed for these disproportions are the near-absence of effective prevention through early detection and prompt treatment programmes, and the lack of equal access to such programmes [2]. Without these interventions, patients with cervical cancer usually present at an advanced stage when it is too late for effective treatment, resulting in high mortality [2,4]. Even the few that may present in the premalignant stage, are often lost to follow-up [3,4]. Other reasons that have been adduced include the ageing and growth of the population, inadequate health facilities and access to it, poor nutrition, late presentation, and insufficient facilities for treatment [2-5].

The incidence and deaths from this malignancy in developing countries have persistently remained on the increase<sup>2</sup>. An estimated 604,127 new cases were diagnosed worldwide in 2020 with associated 341,831 deaths in the same year [6]. This is even though, it is a preventable and curable disease if detected early [3]. The malignant phase often evolves from a long precancerous state that can be detected by screening and treated appropriately [3,4]. Follow up assessment is easy as the cervix is easily accessible [7].

Evidence has shown that HPV infection and health-related lifestyles influence the progression of CIN [8,9]. Analyses of cervical neoplasia lesions showed the presence of HPV in more than 80% of all CIN lesions and 99.7% of all invasive cervical cancers [10].

Risk factors for CIN include early coitarche, multiple sexual partners, history of sexually transmitted infections, cigarette smoking, immunosuppressive conditions, multiparity and long-term use of oral contraceptive pills [10-13]. Bacterial vaginosis and trichomonas vaginalis infections are significantly associated with persistent HPV infection and the development of CIN [14].

The diligent implementation of wide-scale secondary prevention interventions, which comprises effective screening, timely diagnosis and prompt treatment for pre-cancer and early cancer, has engendered a reduction in the incidence of cervical cancer and its associated mortality in developed countries of the world [2,12]. The reverse is the case in developing countries and this is due to financial constraints, lack of political will and an unwillingness to surmount barriers in the implementation of policies aimed at cervical cancer reduction [7,15]. Thus, cervical intraepithelial neoplasia progressing to cervical cancer has continued to rise in these sub-regions [15].

Other factors responsible for the high cervical cancer burden include lack of awareness, low socioeconomic status, poor health-seeking behaviour, inadequate and overburdened health facilities, insufficient supplies, cultural and religious beliefs and most importantly, loss to follow up in patients with premalignant lesions [7,16]. A major factor in the loss to follow-up of patients is that as reported by Nketim [17] who opined that “about 60–75% of women in sub-Saharan Africa who develop cervical cancer live in rural areas. Many of these women go untreated, mostly due to lack of access (financial and geographical) to a health facility” [17].

The employment of Colposcopy in the 1970s revolutionized the management of cervical intraepithelial neoplasia (CIN), with the introduction of conservative treatment modalities [7,15,16]. These include cryotherapy, electro-diathermy, carbon dioxide laser and cold coagulation. Excisional methods include cone biopsies (knife, loop and laser), large loop excision of the transformation zone (LLETZ), and needle/straight-wire excision of the transformation zone [8,18,19]. However, these

treatment options are associated with the risk of recurrence of the disease [8,18,19]. Consequently, a robust and diligent follow-up strategy must be in place.

While much can be achieved in the developed countries, Hysterectomy has remained the mainstay treatment modality for CIN in developing countries due to inadequate screening programs as well as fear of loss of the patient to follow-up for the reasons stated above [17,20].

Organized population-based cervical cancer screening linked to prompt treatment of abnormal results has reduced morbidity and mortality from the disease in developed countries [16,17]. The ability to reliably follow up with patients after treatment has allowed less invasive treatment approaches such as ablative and excisional techniques on the cervix and thus avoiding TAH [17].

In developing countries like Nigeria, factors militating against organized screening programs and challenges with patient follow up have led to the use of TAH in the management of cervical intraepithelial neoplasia (CIN) [20,21].

The study aimed to evaluate the prevalence and other related factors of CIN as a primary indication for TAH performed at the University of Benin Teaching Hospital and compare it with TAH done for benign gynaecological conditions.

## 2. Materials and methods

A retrospective, case-controlled analytical study was carried out at the University of Benin Teaching Hospital (UBTH), Benin City, Nigeria. Information was retrieved from case records of patients who had TAH for benign gynaecological conditions including CIN from January 2015 to December 2019.

### 2.1 Data management:

A database was generated containing information on socio-demographic characteristics and relevant clinical information of the study population. Subgroup analysis was performed comparing patients who had Hysterectomy for CIN and those who had Hysterectomy for other benign conditions. The outcomes of interest include the sociodemographic profile of the patients and the outcome of surgery. The social classification was done as outlined by Oyedeji [22].

The statistical analysis was done using Statistical Package for Social Sciences (SPSS) version 22 and results were presented as whole numbers and percentages for categorical variables. While continuous variables are expressed as mean values with standard deviations. Categorical variables were compared using Chi-square or Fisher's exact test and continuous variables were compared with the t-test. Statistical significance p-value less than 0.05.

## 3. RESULTS

A total of 145 patients underwent total abdominal hysterectomy for benign gynaecological conditions from January 2015 to December 2019. The majority of the patients (54.5%) were above 40 years with a mean age of 47.2±5.2 years. 96.6% of the women were parous and a total of 5 (3.4%) women were nulliparous comprising 4 celibates and 1 married woman. 37.9% of the women were in social class 2 while 7.6% were in social class 5. (Table 1).

The commonest indications for TAH were uterine fibroids (51%), endometrial hyperplasia (22.8%) and cervical intraepithelial lesion contributed 17.2%. Of the 25 women that had a total abdominal hysterectomy for cervical intraepithelial neoplasia, all were parous (100%) and 60% were above 50 years. (Table 2). 83.4% of all the women who had TAH reside in the urban area. (Table 3) and 52% of the patients that had TAH for CIN had blood loss of half a litre and more. (Table 4)

**Table 1: Socio-demographic distribution**

Variable	Frequency (145)	Percent (%)
<b>Age group</b>		
30-49	79	54.5


50-69	64	44.1
>70	2	1.4
Mean age (S.D) = 47.2±5.2		
<b>Parity</b>		
0	5	3.4
1-4	93	64.2
>4	47	32.4
<b>Marital Status</b>		
Single	4	2.8
Married	136	93.8
Divorced	5	3.4
<b>Level of education</b>		
No formal education	2	1.4
Primary	18	12.4
Secondary	46	31.7
Tertiary	79	54.5
<b>Occupation</b>		
Skilled	35	24.1
Semi-skilled	49	33.8
Unskilled	61	42.1
<b>Social class</b>		
1	30	20.7
2	55	37.9
3	34	23.5
4	15	10.3
5	11	7.6
<b>Residence</b>		
Urban	121	83.4
Rural	24	16.6

*Social class by Oyedeji\*<sup>22</sup>*

**Table 2: Indications for TAH**

Variable	Frequency (145)	Per cent (%)
Uterine fibroid	74	51.0
Endometrial hyperplasia	33	22.8
CIN	25	17.2
Endometrial polyp	6	4.1
Adenomyosis	3	2.1
Cervical polyp	2	1.4
Molar gestation	2	1.4

**FIG. 1: pie chart showing the type of CIN among the study population**



The majority of the patient (40.5%) had CIN 3

**Table 3: Comparing socio-demographic characteristics**

Variable	TAH for CIN 25 (%)	TAH for Benign lesions 120 (%)	Test of Statistics
<b>Age Group</b>			
30-49	10 (40.0)	69 (57.5)	F.E = 0.188
50-69	15 (60.0)	49 (40.8)	
>70	0 (0.0)	2 (1.7)	
<b>Parity</b>			
0	0 (0.0)	5 (4.1)	F.E = 0.172
1-4	13 (52.0)	80 (66.7)	
>4	12 (48.0)	35 (29.2)	
<b>Marital status</b>			
Single	0 (0.0)	4 (3.3)	F.E = 0.317
Married	23 (92.0)	113 (94.2)	
Divorced	2 (8.0)	3 (2.5)	
<b>Level of education</b>			
No formal education	0 (0.0)	2 (1.7)	F.E = 0.173
Primary	5 (20.0)	13 (10.8)	
Secondary	4 (16.0)	42 (35.0)	
Tertiary	16 (64.0)	63 (52.5)	
<b>Occupation</b>			
Skilled	3 (12.0)	58 (48.3)	P = .001
Semi-skilled	10 (40.0)	39 (32.5)	
Unskilled	12 (48.0)	23 (19.2)	
<b>Residence</b>			
Urban	20 (80.0)	101 (84.2)	P = .610
Rural	5 (20.0)	19 (15.8)	

F. E= fisher's exact, P = P-value.

**Table 4: Comparison of the complications associated with both study group**

Complications	TAH		Test of statistics
	CIN 25 (%)	Other benign lesions 120 (%)	
Blood loss			
<500ml	12 (48.0)	64 (53.3)	<i>P</i> = .471
500 –1000ml	11 (44.0)	39 (32.5)	
>1000ml	2 (8.0)	17 (14.2)	
Injury to GUT/GIT			
Yes	0 (0.0)	3 (2.5)	F.E = 1.000
No	25 (100.0)	117 (97.5)	
Wound breakdown/sepsis			
Yes	1(4.0)	3 (2.5)	F.E = 0.535
No	24 (96.0)	117 (97.5)	
Anaesthetic complications			
Yes	1 (4.0)	0 (0.0)	F.E = 0.172
No	24 (100.0)	120 (99.2)	
Hospital stay >7days			
Yes	1 (4.0)	12 (10.0)	<i>p</i> = .339
No	24 (96.0)	108 (90.0)	

#### 4. Discussion

The prevalence of total abdominal hysterectomy for cervical intraepithelial neoplasia was 17.2% and this is comparable to the report by Oseki et al [23] but higher than that reported by other researchers [21,24,25]. Anzaku et al [20] reported a prevalence of 27%. The high prevalence in this study reflects the fact that there was a lack of equipment for conservative treatment methods for CIN. This is against the backdrop that the hospital is a major referral centre and a large proportion of the women are older with a completed family size.

The majority of the patients were in their fifth decade of life with a mean age of  $47.2 \pm 5.2$  years and 96.4% of the women were multiparous. These findings are corroborated by other similar studies [23,26, 27]. Completion of family size tends to be a key consideration during counselling, considering the high premium placed on childbearing, especially among the reproductive age group in our contemporary society [21,28]. Most of the women in this study were menopausal or perimenopausal, parous, and have completed their family size. Thus, counselling these patients for total abdominal hysterectomy for cervical premalignant lesions didn't pose many challenges for both the women and the gynaecologists. However, only 5 women (3.4%) were nulliparous. 4 were clergies and only one (1) was married.

When compared to other benign gynaecological conditions, hysterectomy for CIN ranked as the second most common indication after uterine fibroids. A similar find was reported by other studies [20, 21, 23, 28]. The choice of hysterectomy in the treatment of CIN has been considered an overtreatment since the advent of ablative and local excision techniques [20,21,25]. Hence the adoption of the conservative approach in the developed countries such as ablative and excisional methods, especially for women in the reproductive age bracket who desire pregnancy in the future [28,29]. In this case, a

robust follow-up plan is put in place to prevent recurrence and the development of invasive cervical cancer. Such practice remained elusive in most parts of sub-Saharan Africa due to the paucity of the required skills and lack of an organized screening program [21,23,24]. Furthermore, the risk of losing patients to follow-up review/examination may heighten the resultant morbidity and mortality associated with the malignant form of the disease [20,28]. As a result, the practice of total abdominal hysterectomy for cervical intraepithelial neoplasia has remained the mainstay treatment modality for patients who consents, in poor resource settings [20,25].

The procedure is not completely innocuous as complications may arise from the procedure itself or the anaesthesia. Similar to other studies [18,19,24], notable complications amongst women that had a hysterectomy for CIN were haemorrhage (8.0%) and sepsis/wound breakdown (4.0 %) leading to a prolonged hospital stay. While one patient (4.0 %) had prolonged recovery from anaesthetic complications.

Conversely, a higher number of women had complications associated with hysterectomy for other benign gynaecological conditions. In consonance with other studies [20,22,24,26], 17(14.2%) women suffered haemorrhages and 12(10.0%) had prolonged hospital stays. Wound breakdown was recorded in 3(2.5%) women and another 3(2.5%) women had injuries to the gut. These findings tend to suggest that hysterectomy for CIN may not be associated with an increased risk of surgical complications. However, over 50% of women who had TAH for CIN had >500ml blood loss at the surgery. This is contrary to the application of ablative and local excision methods which are associated with minimal blood loss, shorter operation time and short duration of hospital stay [30].

## **5. CONCLUSION**

Hysterectomy is the gold standard in the management of benign gynaecological conditions. Moreover, with the advent of effective and efficient ablative and local excisional procedures, it is now considered an overtreatment for women with premalignant lesions of the cervix, especially in developed countries of the world. However, in poor and/or low resource countries, with scarce facilities and skills for local excision and ablative procedures, hysterectomy is still paramount as a treatment option for women with cervical intraepithelial neoplasia. Consequently, with the attendant risk and complications following TAH for cervical dysplasia, there is a need for the provision of accessible infrastructural facilities and requisite skills for minimally invasive procedures for the treatment of cervical dysplasia in developing countries.

## **ETHICAL APPROVAL (WHERE EVER APPLICABLE)**

The approval for the study was obtained from the Research and Ethics Committee of the University of Benin Teaching Hospital, Benin City.

## **REFERENCES**

1. Chirenje ZM. HIV and cancer of the cervix. *Best Pract Res Clin Obstet Gynaecol.* 2005;19 :269–276.

2. World Health Organization. Guideline for screening and treatment of cervical pre-cancer lesions for cervical cancer prevention. In: 2nd ed. Geneva; 2021.
3. Utoo BT, Utoo PM, Ngwan SD, Anzaku SA, Daniel MA. Cervical intraepithelial neoplasia: prevalence, risk factors, and utilization of screening services among an urban population in Nigeria. *Trop J Obstet Gynaecol.* 2016; 33: 279.
4. Balogun MR, Odukoya OO, Oyediran MA, Ujomu PI. Knowledge of cervical cancer and its risk factors among women residing in two urban slums in Lagos, Nigeria. *Niger Med Pract.* 2012;61: 74–81.
5. Anorlu RI. Cervical cancer: the sub-Saharan African perspective. *Reprod Health Matters.* 2008;16: 41– 49.
6. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin.* 2021; 71(3):209 – 49.
7. Johnson LG, Armstrong A, Joyce CM, Teitelman AM, Buttenheim AM. Implementation strategies to improve cervical cancer prevention in sub-Saharan Africa: a systematic review. *Implement Sci.* 2018;13: 1 – 18.
8. Tierney B, Westin SN, Schlumbrecht MP, Ramirez PT. Early cervical neoplasia: advances in screening and treatment modalities. *Clin Adv Hematol Oncol HO.* 2010;8(8):547–55.
9. WHO guideline for screening and treatment of cervical pre-cancer lesions for cervical cancer prevention, 2021. [cited 2022 Apr 10]. Available from: <https://www.who.int/publications-detail-redirect/9789240030824>
10. Holschneider CH. Premalignant & Malignant Disorders of the Uterine Cervix. In: DeCherney AH, Nathan L, Laufer N, Roman AS, editors. *Current Diagnosis & Treatment: Obstetrics & Gynaecology*, 11th ed. New York, NY: The McGraw-Hill Companies; 2013. p. 807–831.
11. Daru PH, Pam IC, Musa J, Daniyan MG, Silas OI, Adesina OA, et al. Cervical epithelial changes in a tertiary hospital in Northern Nigeria. *Trop J Obstet Gynaecol.* 2013;30(1).
12. Adisa J, Tukur S, Bukar M, Egbujo E. Prevalence of cervical intraepithelial neoplasia in relation to knowledge, attitudes/beliefs, and practices among university students in North-Eastern Nigeria. *Ann Trop Med Public Health.* 2014;6(4).
13. Teixeira NCP, Araújo ACL, Correa CM, Lodi CT da C, Lima MIM, Carvalho N de O, et al. Prevalence and risk factors for cervical intraepithelial neoplasia among HIV-infected women. *Braz J Infect Dis Off Publ Braz Soc Infect Dis.* 2012;16: 164–169.
14. Tao L, Han L, Li X, Gao Q, Pan L, Wu L, et al. Prevalence and risk factors for cervical neoplasia: a cervical cancer screening program in Beijing. *BMC Pub Health.* 2014;14: 1–9.
15. Wang Z, Wang J, Fan J, Zhao W, Yang X, Wu L, et al. Risk factors for cervical intraepithelial neoplasia and cervical cancer in Chinese women: large study in Jiexiu, Shanxi Province, China. *J Cancer.* 2017; 8: 924 – 932.
16. Adepoju EG, Ilori T, Olowookere SA, Idowu A. Targeting women with free cervical cancer screening: challenges and lessons learnt from Osun state, southwest Nigeria. *Pan Afr Med J* 2016;24:319.
17. Ntekim A. Cervical Cancer in Sub Sahara Africa. In: Rajkumar R, editor. *Topics on cervical cancer With an Advocacy for Prevention.* Rijeka: IntechOpen; 2012.
18. Khan MJ, Smith-McCune KK. Treatment of cervical precancers: back to basics. *Obstet Gynecol.* 2014; 123: 1339 - 1343. doi:10.1097/AOG.0000000000000287
19. Shafi MI. Premalignant and malignant disease of the cervix In: Dewhurst's Textbook of Obstetrics and Gynaecology. D. K Edmonds (ed). Blackwell Publishing Ltd. 2012. Pg 747-759
20. Anzaku AS, Musa J. Total abdominal hysterectomy for benign gynaecological conditions at a University Teaching Hospital in Nigeria. *Niger J Med.* 2012; 21: 326–330.

21. Samaila Modupeola OA, Adesiyun AG, Agunbiade OA, Mohammed-Duro A. Clinico-pathological assessment of hysterectomies in Zaria. *Eur J Gen Med.* 2009; 150–153.
22. Oyedeji GA. Socio-economic and cultural background of hospitalized children in Ilesa. *Niger J Paediatr.* 1985; 12: 111–117.
23. Oseki C, Osaikhuwomwan JA. A Review of indications and outcome of total abdominal hysterectomy at a tertiary public health facility in Southern Nigeria. *New Niger J Clin Res.* 2018; 7: 21.
24. Patel DA, Saraiya M, Copeland G, Cote ML, Datta SD, Sawaya GF. Treatment patterns for cervical carcinoma in situ in Michigan, 1998-2003. *J Regist Manag.* 2013; 40: 84–92.
25. Abbas F, Birara M, Bekele D. Review of Hysterectomies done for Cervical Intraepithelial Neoplasias (CIN) in a teaching referral Hospital. *Ethiop J Reprod Health.* 2014; 7; 41-47.
26. Bukar M, Audu BM, Yahaya UR. Hysterectomy for benign gynaecological conditions at Gombe, north eastern Nigeria. *Niger Med J.* 2010; 51: 35.
27. Ajah LO, Ezeonu PO, Ozonu NC, Iyoke CA, Nkwo PO, Ajah MI. A five year review of cervical cytology in Abakaliki, Nigeria. *Am J Cancer Prev.* 2015; 3 :23–6.
28. Ahmed ZD, Taiwo N. Indications and Outcome of Gynaecological Hysterectomy at Aminu Kano Teaching Hospital, Kano: A 5-Year Review. *Open J Obstet Gynecol.* 2015; 05: 298–304.
29. Papadopoulos MS, Tolikas AC, Miliaras DE. Hysterectomy—Current Methods and Alternatives for Benign Indications. Mettler L, editor. *Obstet Gynecol Int.* 2010; 28; 2010:356740. Available from: <https://doi.org/10.1155/2010/356740>.
30. WHO Guidelines for Treatment of Cervical Intraepithelial Neoplasia 2–3 and Adenocarcinoma in situ Cryotherapy, Large Loop Excision of the Transformation Zone, and Cold Knife Conization Geneva: World Health Organization; 2014. ISBN-13: 978-92-4-150677-9